

CATALOG OF THE LACERTIDAE



Harold De Lisle

Cover: *Gallotia galloti galloti* (Oudart, 1839)
Photo by David Modrý
Southern Tenerife, Canary Islands

Family Lacertidae

A CATALOG OF RECENT SPECIES

Harold F De Lisle

© Harold F. De Lisle, 2016
Email: gambelia@hughes.net

Lacertidae

The **Lacertidae** is a family which are native to Europe, Africa, and Asia. The group includes the genus *Lacerta*, which contains some of the most commonly seen lizard species in Europe. It is a diverse family with some 323 species in 43 genera.

The European and Mediterranean species live mainly in forest and scrub habitat, the Asian species inhabit grasslands and deserts. African species usually live in rocky, arid areas. One genus is arboreal.

Lacertids are small or medium-sized lizards. Most species are less than 9 cm long, excluding the tail, although the largest living species, *Gallotia stehlini*, reaches 46 cm, and some extinct forms were larger still. They are primarily insectivorous. Lacertids are remarkably similar in form, with slender bodies and long tails, but have highly varied patterns and colours, even within the same species. Their scales are large on the head, which often also has osteoderms, small and granular on the back, and rectangular on the underside. Most species are sexually dimorphic.

At least eight species are parthenogenetic, and three species give birth to live young.

Pyron & Burbrink (2013) have the best phylogenetic survey (see below). It shows that most of the generic arrangements are correct, with the exception of *Ichnotropis* and *Nucras* which might be better contained within other genera.


The systematic arrangement of the catalog is intuitive once a few principles are stated. Subspecific entries for each species are indented. Dates in brackets indicate that these have been subsequently determined from sources external to the original document. Misspellings are generally excluded except where they have been widely used or misinterpreted. Junior subjective and objective synonyms are listed under each valid name. Alternate taxonomic arrangements are included in a taxonomic note in many cases.

In order to enhance the usefulness of the present list, the type species of each genus has been added keeping in mind the famous quote of Harold Cogger, "Taxonomy is a matter of opinion" (Golay et al., 1993) based, of course, on the evaluation of the best evidence. The taxonomy used herein is our evaluation and we accept full responsibility for its use.

We have used alphabetical ordering for utility of use, ignoring the ever-changing families. Junior objective and subjective synonyms follow each senior synonym. We have deliberately omitted most lapsus calami, nomina nuda, nomina dubia, nomina oblita, nomina substituta, etc., from the synonymy because these names are not available for use. The synonymy herein is limited to new descriptions. The synonymy used herein is for the most part based in Kluge (2001). The Catalog of Gekkota covers more than 1800 species and subspecies, almost 100 genera, and includes almost 1000 bibliographic references. Entries for species consist of species or subspecies name, genus, author, year of description, publication page, type locality, location of primary (name-bearing) type specimen(s), if known, and known distribution. Type localities are those in the original description, except those in obvious error or those amended subsequently by

authoritative sources. Acronyms employed in the catalogue follow those used by American Museum of Natural History.

The definitions of the type categories Holotype, Syntype, Paratype, Lectotype and Neotype follow those given in the glossary and articles of the ICZN Code (ICZN, 1999). In addition, we also use the following term: Iconotype. Some early 18th century species descriptions were done from paintings of live animals. More recently, photographs of some rare animals for which no museum specimens exist are termed iconotypes. The synonymy in this paper is our own. We have deliberately omitted all lapsus, nomina nuda, etc., from the synonymy because these names are not available for use. The synonymy herein is limited to new descriptions - junior subjective synonyms.



Abbreviations of museums holding the types are as follows

- AMNH** - American Museum of Natural History, Division of Vertebrate Zoology (Herpetology), 79th Street and Central Park West, New York, New York 10024, USA.
- ANSP** - Academy of Natural Sciences, Department of Herpetology, 19th and the Benjamin Franklin Parkway, Philadelphia, Pennsylvania 19103, USA.
- BMNH** - The Natural History Museum, Department of Zoology, Cromwell Road, London SW7 5BD, United Kingdom.
- CA** - Universidad Autònoma de Barcelona, Plaça Cívica, s/n, 08193 Bellaterra, Barcelona, Spain.
- CAS** - California Academy of Sciences, Department of Herpetology, Golden Gate Park, San Francisco, California 94118, USA.
- CNR** - Caucasus Nature Reserve Russia, Krasnodar Krai, Sochi, Russia
- DBULPGC** - Departamento de Biología de la Universidad de Las Palmas, Calle Juan de Quesada, 35001 Las Palmas de Gran Canaria, Las Palmas, Spain
- DC** - private collection of Domenico Capolongo, Via Roma 8, 80030 Roccarainola, Napoli, Italia
- DZLU** - Departamento de Zoología, Universidad de La Laguna, Calle Heraclio Sánchez, 43, 38204 San Cristóbal de La Laguna, Santa Cruz de Tenerife, Spain
- EBD** - Doñana Biological Station, Lugar Urbanización Playa de Matalascañas, 0 S/N, 21730 Almonte, Huelva, Spain
- FMNH** - Field Museum, Division of Amphibians and Reptiles, Roosevelt Road at Lake Shore Drive, Chicago, Illinois 60605, USA.
- G** - Institute for Biological Research "Siniša Stanković", 142 Bulevar despota Stefana, Belgrade, Serbia
- GNM (MC)** - Georgian National Museum, 3 Shota Rustaveli Ave, Tbilisi, Georgia
- HLMD** - Hessisches Landesmuseum, Darmstadt, Hessen, Friedenspl. 1, 64283 Darmstadt, Germany
- HNHM** - Hungarian Natural History Museum, 1083 Budapest, Ludovika tér 2-6, Hungary
- HUJ** - Hebrew University, Department of Zoology, Jerusalem, Israel
- IPE** - Instituto Pirenaico de Ecología, Av. Ntra. Sra. de la Victoria, S/N, 22700 Jaca, Huesca, Huesca, Spain .
- IRSNB** - Institut Royal des Sciences Naturelles de Belgique, 29 Rue Vautier, 1000 Bruxelles, Belgium.
- KUZ** - Department of Zoology, Graduate School of Science, Kyoto University, Sakyo, Kyoto 606-8502, Japan
- MBL** - Museu Bocage, Universidade de Lisboa, Lisbon, Portugal [destroyed by fire in March 1978]
- MCVR** - Museo Civico di Storia Naturale, Verona, Porta Vittoria 9 - 37129 Verona, Italy
- MCZ** - Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138, USA
- MHNCF** - Musée d'histoire naturelle, Avenue Léopold-Robert 63, 2300 La Chaux-de-Fonds, Switzerland
- MD** - Museu Regional do Dundo, Dundo, Luanda-Norte, Angola.
- MISN** - Museo Insular de la Palma, Santa Cruz de la Palma, Plaza San Francisco, 3, 38700 Santa Cruz de la Palma, Santa Cruz de Tenerife, Spain
- ML** - Slovenian Museum of Natural History, Prešernova cesta 20, Ljubljana, Slovenija
- MLSU** - St. Petersburg State University, Zoological Museum, 199034, Universitetskaya Emb., 7/9 St. Petersburg, Russia
- MM** - Museum für Naturkunde Magdeburg, Magdeburg, Saxony Otto-von-Guericke-Straße 68, 39104 Magdeburg, Germany
- MMK** - McGregor Museum, Kimberley, North Cape Province, Egerton Rd, Kimberley, 8301, South Africa
- MMTT** - National Museum of Natural History, Niavaran Square, Darabad Street, Tehran 11369, Iran
- MNCN** - Museo Nacional de Ciencias Naturales, Calle de José Gutiérrez Abascal, 2, 28006 Madrid, Spain
- MNHG** - Museum d'Histoire Naturelle, Geneva, Switzerland.
- MNHNP** - Museum National d'histoire Naturelle, Laboratoire des Amphibiens et Reptiles, 25 Rue Cuvier, 75005 Paris, France
- MNKNU** - Museum of Nature, V. N. Karazin Kharkiv National University, Kharkiv, Ukraine
- MRSN** - Museo Storia Naturale di Torino, Italy

MSNG - Museo Civico di Storia Naturale di Genova Giacomo Doria, Via Brigata Liguria 9, 16121 Genova, Italy
MTKD - Staatliche Naturhistorische Sammlungen Dresden, Museum für Tierkunde, Königsbrücker Landstrasse 159, 01109 Dresden, Germany.
MZB - Parc del Fòrum, Plaça Leonardo da Vinci, 4-5, 08019 Barcelona, Spain
MZUF - Università di Firenze, Museo Zoologico La Specola, Via Romana 17, 50125 Firenze, Italy
MZUP - Museo Zoologico dell'Università di Palermo, Via Archirafi, 16, 90100 Palermo, Italy
MZUT - Museo ed Istituto di Zoologia Sistemática, via Giolitti 36, 10123 Torino, Italy
NAS KR - Museum of the Institute of Biology, National Academy of Sciences, Bishkek, Kyrgyz Republic
NCBS - National Centre for Biological Sciences, Bangalore, India
NHMC - Natural History Museum of Crete, Leof. Sofokli Venizelou, Iraklio 712 02, Greece
NHMG - Naturhistorisches Museum, Box 11049, Slottskogen, 40030 Göteborg, Sweden
NHMW - Naturhistorisches Museum, Zoologische Abteilung, Postfach 417, Burgring 7, 1014 Wien, Austria
NMNW - State Museum Windhoek, Robert Mugabe Ave City Centre, Windhoek, Namibia
NHRM - Naturhistoriska Riksmuseet, Section for Vertebrate Zoology, Roslagsvägen 120, Box 50007, 104 05 Stockholm 50, Sweden.
NMB - National Museum, Bloemfontein, 36 Aliwal St, Bloemfontein Central, Bloemfontein, 9301, South Africa
NMNH - Museum of the National Academy of Sciences of Ukraine, Kiev
NMNS - National Museum of Natural Sciences, Guanqian Rd, North District, Taichung City, Taiwan 404
NMP6V - Naturhistorisches Nationalmuseum, Prague, Czech Republic
NMSL - Sri Lanka National Museum, PO Box 854, Colombo 07, Sri Lanka
NMZB - National History Museum of Zimbabwe, Bulawayo, Zimbabwe
NZMS - National Museum of Natural History, Bulgarian Academy of Sciences, 1000 Sofia Center, Sofia, Bulgaria
OA - private collection of Oscar J Arribas Amo
PEM - Port Elizabeth Museum, Port Elizabeth, Republic of South Africa
PMNH - Pakistan Museum of Natural History, Islamabad, Pakistan
RMCA - Royal Museum for Central Africa, Leuvensesteenweg 13 3080 Tervuren, Belgium
RUZM - Zoology Department Museum, Razi University, Kermanshah, Iran
SAM - South African Museum, P.O. Box 61, Cape Town 8000, Western Cape Province, South Africa.
SMF - Forschungsinstitut und Natur-Museum Senckenberg, Senckenberg-Anlage 25, 6000 Frankfurt-am-Main 1, Germany.
SMNS - Staatliches Museum für Naturkunde, Stuttgart, Germany
SNP - Herpetological collection, Sochi National Park, Russia
SUHC -
SZE - Zoology Department, Aegean University, Bornova-Izmir, Turkey.
TAU - Tel Aviv University, Tel Aviv, Israel.
THUP - Biology Department, Tunghai University, Taichung, Taiwan
TMP - Transvaal Museum, P.O. Box 413, Pretoria 0001, Gauteng, South Africa.
TNZ - Tianjin Natural History Museum, 206 Machang Rd, Hexi Qu, Tianjin Shi, China, 300074
USNM - National Museum of Natural History, Division of Amphibians and Reptiles, Washington, D.C. 20560, USA.
UTEP - University of Texas at El Paso, Department of Biological Sciences and Centennial Museum, Laboratory for Environmental Biology, 500 W University Ave, El Paso, TX 79968, USA
UUZM - Uppsala University Museum of Evolution, Zoology section, Uppsala, Sweden
VPM - private collection of Valentin Pérez Mellado
ZCBS/M - Dipartimento di Zoologia e Museo di Storia Naturale della Baviera, Monaco
ZDEU - Zoology Department, Ege University, Izmir, Turkey
ZFMK - Zoologisches Forschungsinstitut und Museum Alexander Koenig, Herpetologische Abteilung, Adenauerallee 150-164, 5300 Bonn-1, Germany.
ZIA - Institute of Zoology, 7, Sevak Str., Yerevan 0014, Republic of Armenia
ZIK - Ukrainian Academy of Sciences, Zoological Institute, 252.150 Kiev: 30, Ukraine.
ZISP - Zoological Institute, Russian Academy of Sciences, St. Petersburg 199034, Russia.
ZIT - Georgian Academy of Sciences, Tbilisi, Georgia 380077

ZIUG - Zoologisches Museum der Georg-August Universität Göttingen, Göttingen, Niedersachsen Germany

ZMA - Universiteit van Amsterdam, Zoologisch Museum, Plantage Middenlaan 53, Amsterdam: C, Netherlands
(now incorporated into NCB Naturalis, Leiden).

ZMB - Universität Humboldt, Zoologisches Museum, Invalidenstrasse 43, 10115 Berlin, Germany.

ZMH - Zoologisches Institut und Museum, Universität Hamburg, Martin-Luther-King Platz 3, D-2000 Hamburg 13,
Germany

ZMMU - M. V. Lomonosov Moscow State University, Zoological Museum, Ul. Herzena 6, 103009 Moscow K-9,
Russia.

ZSM - Zoologisches Staatssammlung, Münchhausenstrasse 21, 8000 München 21, Germany.

ZSI - Zoological Survey of India, Herpetology Section, Fire-Proof Spirit Building, 27 JL Nehru Road, Kolkata 700
016, India.

Lacertidae Opperl, 1811 Opperl, M. 1811. *Die Ordnungen, Familien und Gattungen der Reptilien, als Prodrum einer Naturgeschichte derselben*. Munchen: Joseph Lindauer, xii+87 pp. Type genus: *Lacerta* Linnaeus, 1758

***Acanthodactylus* Wiegmann, 1834** Wiegmann, A.F.A. 1834. *Herpetologica Mexicana, seu descriptio*



amphibiorum Novae Hispaniae. Pars Prima, Saurorum Species, Berlin: C.G. Lüderitz. vi + 54 pp. + Plates I-X. (*Acanthodactylus*, p. 10). Type species: *Lacerta boskiana* Daudin, 1802

Acanthodactylus schmidti Seyed B Mousavi

Acanthodactylus aegyptius Baha El Din, 2007

[Acanthodactylus aegyptius Baha El Din, 2007: ?](#)

Type locality: El Maghra Oasis (30°15'N, 28°55'E), Egypt

Holotype: FMNH 269501

Distribution: western Egypt to northern Sinai; southern Israel.

Acanthodactylus ahmaddisii Werner, 2004

[Acanthodactylus ahmaddisii Werner, 2004: 40](#)

Type locality: 36 km south of Amman, Jordan

Holotype: HUI R1296

Distribution: northwestern Jordan Last seen 1936

Acanthodactylus arabicus Boulenger, 1918

[Acanthodactylus cantoris arabicus Boulenger, 1918: 154](#)

Type locality: southern Arabia (Aden)

Syntypes: BMNH 1946.8.4.33-42; MCZ 28685

Distribution: southern Yemen

Acanthodactylus aureus Günther, 1903

[Acanthodactylus inornatus aureus Günther, 1903: 298](#)

Type locality: Villa Cisneros [= Dakhla], Western Sahara

Lectotype: MNHNP 1918.11

Distribution: coastal southwestern Morocco; Western Sahara; western Mauritania; northern Senegal.

Acanthodactylus bedriagai Lataste, 1881

[Acanthodactylus bedriagai Lataste, 1881:357](#)

Type locality: El Guernah, Hauts Plateaux, Algeria

Syntypes: BMNH 1946.9.3.51, 1920.1.20.1324

Distribution: northern Algeria and adjoining Tunisia

Acanthodactylus beershebensis Moravec, El Din, Seligmann, Sivan & Werner, 1999

[Acanthodactylus beershebensis Moravec, El Din, Seligmann, Sivan & Werner, 1999: 39](#)

Type locality: Be'er Sheva, subdistrict , 5 km E of 'Omer-Hebron-' Arad crossroads (Israel Grid 14350770), i.e. around the Nahal Yattir crossing of the Zomet Shoqet - 'Arad road, Israel.

Holotype: HUI R 7621

Distribution: Israel: Negev Desert

Acanthodactylus blanci Doumergue, 1901

Acanthodactylus blanci Doumergue, 1901: 184

Type locality: Hammam-El-Lif, near Tunis

Syntypes: BMNH 1946.9.3.3-5

Distribution: northern Tunisia; northern Algeria

Acanthodactylus blanfordii Boulenger, 1918

Acanthodactylus cantoris blanfordii Boulenger, 1918: 154

Type locality: Balochistan, Iran

Syntypes: BMNH 1946.9.3.54-55

Distribution: southeastern Iran; southern Afghanistan; western Pakistan; northeastern Oman.

Acanthodactylus boskianus (Daudin, 1802)

Acanthodactylus boskianus boskianus (Daudin, 1802)

Lacerta boskiana Daudin, 1802: 188

Type locality: unknown

Holotype: MNHNP0.2762

Distribution: Egypt: Nile Delta area, northern Sinai

Acanthodactylus boskianus asper (Audouin, 1829)

Lacerta asper Audouin, 1829: 121

Type locality: Egypt Holotype:

not located

Synonyms:

Scapteira inaequalis Gray, 1838 Type locality: Egypt Holotype: BMNH 1849.3.29.15

Acanthodactylus boskianus khattensis Trape & Trape, 2012 Type locality: the bushes of the bed of a "khatt" located at the foot of the butte Mohammed Moûloûd (19 ° 47 '46' N, 14 ° 25 '27' W), 8 km northwest of Akjoujt in Mauritania Holotype: MNHNP 2011.0223

Acanthodactylus boskianus nigeriensis Trape, Chirio & Geniez, 2012 Type locality: near Birmin-Gaouré (13° 05' N, 02° 55' E), Niger Holotype: MNHNP 2011.0273

Distribution: Morocco, Algeria, Tunisia, Libya, Egypt, Mauritania, Mali, Niger, northern Nigeria, Chad, Sudan, Eritrea, the whole Arabian Peninsula, Israel, Lebanon, Jordan, Palenstine, Syria, Iraq, southwestern Iran

Acanthodactylus boskianus euphraticus Boulenger, 1919

Acanthodactylus boskianus euphraticus Boulenger, 1919: 549

Type locality: Euphrates valley, Iraq

Holotype: BMNH, not located

Distribution: Iraq, southeastern Turkey

Acanthodactylus boueti Chabanaud, 1917

Acanthodactylus boueti Chabanaud, 1917: 87

Type locality: Agouagou, Dahomey

Syntypes: MNHNP 1917.50-2, 1917.54-6; BMNH 1946.8.4.68

Distribution: northern Ghana, northern Benin, northern Dahomey, western Nigeria, Togo

Acanthodactylus busacki Salvador, 1982

Acanthodactylus busacki Salvador, 1982: 88

Type locality: 30 km south of Goulimine, Morocco

Holotype: BMNH 1970.250

Distribution: southwestern Morocco; northern Western Sahara.

Acanthodactylus cantoris Günther, 1864

Acanthodactylus cantoris Günther, 1864: 73

Type locality: Ramnagar, India

Syntypes: BMNH 1946.8.4.15-20.

Distribution: eastern Afghanistan; Pakistan; India: Punjab, Uttar Pradesh, Rajasthan, Gujarat States

Acanthodactylus dumerilii (Milne-Edwards, 1829)

Lacerta dumerilii Milne-Edwards, 1829: 76

Type locality: unknown

Lectotype: MNHNP 0.2759

Synonyms:

Scapteira inornata Gray, 1838 Type locality: Tripoli, Libya Holotype: BMNH 1946.9.3.76

Acanthodactylus scutellatus exiguus Lataste, 1885 Type localities: Algeria: Biskra, Mraier, Tougourt, Bled

Ahmar, Hadjira, N'Gouça, Tilremt, Laghouat, Ain el Hel et Bou-Saada. En Tunisie, à l'oued el Kreil, Kebili,

Tozeur, Nefta Syntypes: BMNH 1920.1.20.1349c (Bled Ahmar), 1920.1.20.1349 (Hadjira),

1920.1.20.1349d (Tilremt), 1920.1.20.1349e (Laghouat), 1920.1.20.1349i (Mraier), 1920.1.20.1349g

(Biskra), 1920.1.20.3006 (Oued el Kreil)

Distribution: western Senegal, western Mauritania, Western Sahara, southern Morocco, southern Algeria, southern Tunisia, northwestern Libya, Mali

Acanthodactylus erythrurus (Schinz, 1833)

Acanthodactylus erythrurus erythrurus (Schinz, 1833)

Lacerta erythrura Schinz, 1833: 102

Type locality: Spain

Holotype: not located

Synonyms:

Acanthodactylus vulgaris Duméril & Bibron, 1839 Type locality: southern Spain Holotype: not located

Acanthodactylus vulgaris bocagei Ferreira, 1892. Type locality: Portugal. Holotype,

MBL. Presumably destroyed during the 1978 fire.

Distribution: Portugal, Spain, Gibraltar

Acanthodactylus erythrurus atlanticus Boulenger, 1918

Acanthodactylus erythrurus atlanticus Boulenger, 1918: 149

Type locality: Tamaruth Valley, Morocco

Syntypes: BMNH 1946.9.2.72-73

Distribution: Morocco: Middle Atlas including the Central Plateau, northern slopes of the High Atlas, plains to the north and west of the Atlases

Acanthodactylus erythrurus belli Gray, 1845

Acanthodactylus bellii Gray, 1845: 36

Type locality: Algiers, Algeria

Syntypes: BMNH 1946.9.3.8-11

Synonyms:

Acanthodactylus erythrurus ksourensis Doumergue, 1901 Type locality: Stitten, El Bayadh, Algeria Holotype: not located

Acanthodactylus vulgaris mauritanicus Doumergue, 1901 Type locality: Oran, Algeria Holotype: not located

Distribution: northeastern Morocco and the coastal regions and high plateaux in Algeria.

Acanthodactylus erythrurus lineomaculatus Duméril & Bibron, 1839

Acanthodactylus lineomaculatus Duméril & Bibron, 1839: 276

Type locality: Mogador [=Essaouira], Morocco

Lectotype: MNHNP0.5912

Synonym:

Acanthodactylus vulgaris tingitanus Doumergue, 1901 Type locality: Tangier, Morocco Holotype: not located

Acanthodactylus felicitis Arnold, 1980

Acanthodactylus felicitis Arnold, 1980: 300

Type locality: Hadhramaut', Yemen

Holotype: BMNH 1897.3.11.65

Distribution: southern Yemen; western Oman

Acanthodactylus gongrorhynchatus Leviton & Anderson, 1967

[Acanthodactylus gongrorhynchatus Leviton & Anderson, 1967: 171](#)

Type locality: Beda Azan (23°41'N, 53°28'E), Abu Dhabi

Holotype: CAS 97803

Distribution: eastern Saudi Arabia; western United Arab Emirates

Acanthodactylus grandis Boulenger, 1909

[Acanthodactylus grandis Boulenger, 1909: 189](#)

Type localities: Jerud and Ataibe, east of Damascus, and Khan Agach, between Damascus and Kutaife, Iraq

Syntypes: BMNH 1909.4.20.29 (Jerud), MNHNP 23.8-11, BMNH 1946.9.2.69-70 (Khan Agach).

Synonym:

Acanthodactylus fraseri Boulenger, 1918 Type locality: Az Zubayr (30°23'N, 47°43'E), Shariba, Iraq

Holotype: BMNH 1946.8.7.40.

Distribution: eastern Lebanon; Syria; Jordan; Iraq; northern Saudi Arabia; southwestern Iran

Acanthodactylus guineensis (Boulenger, 1887)

[Eremias guineensis Boulenger, 1887: 51](#)

Type locality: Brass, mouth of the Niger, southern Nigeria

Holotype: BMNH 1946.8.6.31

Synonym:

Eremias benuensis Monard, 1949 Type locality: Ngaouyanga, Cameroon Holotype: MNHG 1055.62

Distribution: Mali, Burkina Faso, northeastern Togo, Ghana, southwestern Niger, Nigeria, northern Cameroon, southwestern Chad and Central African Republic

Acanthodactylus haasi Leviton & Anderson, 1967

[Acanthodactylus haasi Leviton & Anderson, 1967: 177](#)

Type locality: Dhahran (Az Zahran), Saudi Arabia

Holotype: CAS 84596

Distribution: northern and eastern Saudi Arabia; United Arab Emirates: Abu Dhabi Emirate; Oman: Wahiba sands

Acanthodactylus hardyi Haas, 1957

[Acanthodactylus scutellatus hardyi Haas, 1957: 72](#)

Type locality: Hirmas Station, Saudi Arabia

Holotype: HUI 2682

Distribution: Jordan, Iraq, Saudi Arabia, Kuwait

Acanthodactylus harranensis Baran, Kumlutas, Lanza, Sindaco, Avci & Crucitti, 2005

[Acanthodactylus harranensis Baran, Kumlutas, Lanza, Sindaco, Avci & Crucitti, 2005: 324](#)

Type locality: at Harran (36°51' N, 39°00' E), near the ruins of the ancient university, ca.75 km as the crow flies from Sanliurfa, Sanliurfa province, southeastern Turkey

Holotype: ZDEU 65/2001:1

Distribution: southeastern Turkey

Acanthodactylus khamirensis Heidardi, Rastegar-Pouyani, Rastegar-Pouyani & Rajabizadeh, 2013

[Acanthodactylus khamirensis Heidardi, Rastegar-Pouyani, Rastegar-Pouyani & Rajabizadeh, 2013: 335](#)

Type locality: 7 km east of Khamir port, from coastal line of the Persian Gulf, 1km to sea shore (26° 30' 47"N, 055° 58' 44.2"E), Hormozgan province, southern Iran.

Holotype: RUM 146

Distribution: southern Iran: Hormozgan Province

Acanthodactylus longipes Boulenger, 1918

[Acanthodactylus scutellatus longipes Boulenger, 1918: 154](#)

Type locality: Wargla [= Ouargla], Algeria

Lectotype: BMNH 1946.8.31

Synonym:

Acanthodactylus longipes panousei Bons & Girot, 1964 Type locality: Edge of the Erg Chebbi at the level of Merzouga, southeastern Morocco Syntypes: MNHNP 1963.1013-1014
Distribution: Algeria: Sashara; northern Chad; northern Egypt; Libya; northern Mali; Mauritania; southern Morocco, Western Sahara; northern Niger; Tunisia

Acanthodactylus maculatus (Gray, 1838)

Scapteira maculata Gray, 1838: 281

Type locality: Tripoli, Libya

Holotype: BMNH 1946.9.3.53

Synonyms:

Zootoca deserti Günther, 1859 Type locality: Ngoussa Oasis, Algeria Holotype: BMNH 1946.9.3.52

Acanthodactylus pardalis intermedius Doumergue, 1901 Type locality: from the northern limit of the Haut-Plateaux to the Sahara Holotype: not located

Acanthodactylus pardalis latastei Boulenger, 1918 Type locality: southern Tunisia Holotype: not located

Acanthodactylus mechriguensis Nouria & Blanc, 1998 Type locality: Sidi-Mechrig, northern Tunisia
Holotype: not located

Distribution: northern Algeria, northwestern Libya, northeastern Morocco, Tunisia

Acanthodactylus margaritae Tamar, Geniez, Brito & Crochet, 2017

Acanthodactylus margaritae Tamar, Geniez, Brito & Crochet, 2017: 368

Type locality: 500 m north-east of Sidi Binzarne, approximately 3 km south-east of Sidi R'bat (Parc national de Souss-Massa (30.0580°N, 9.6456°W), Morocco

Holotype: MNHNP 2016.105

Distribution: southwestern Morocco

Acanthodactylus masirae Arnold, 1980

Acanthodactylus masirae Arnold, 1980: 303

Type locality: North point, Masirah Island, Sultanate of Oman

Holotype: BMNH 1975.1226

Distribution: Oman: Dhufar, Masira Island

Acanthodactylus micropholis Blanford, 1874

Acanthodactylus micropholis Blanford, 1874:33

Type locality: Gedrosia, Baluchistan, Pakistan

Syntypes: BMNH 74.11.23.79 (Rigan), BMNH 1946.9.3.71-72 (Magas), BMNH 74.11.23.78 (Bampur); ZSI 5301

Distribution: southeastern Iran; western Pakistan; Afghanistan

Acanthodactylus nilsoni Rastegar-Pouyani, 1998

Acanthodactylus nilsoni Rastegar-Pouyani, 1998: 257

Type locality: 34°30'N, 45°33'E, 5 km S Qasr-e Shirin, Kermanshah Province, Iran, about 7 km from Iran-Iraq border

Holotype: NHMG 5145

Distribution: Iran, Kermanshah Province

Acanthodactylus ophiodurus Arnold, 1980

Acanthodactylus ophiodurus Arnold, 1980: 296

Type locality: Jazir coast (18°40' N, 16°40' E), Sultanate of Oman

Holotype: BMNH 1969.314

Distribution: Saudi Arabia, Kuwait, Israel, Jordan, Iraq, Yemen, Oman, United Arab Emirates (UAE), Qatar

Acanthodactylus orientalis Angel, 1936

Acanthodactylus tristrami orientalis Angel, 1936: 109

Type localities: Palmyre, Tell Abiad, Ain Zahra, Deir Ez Zor, Iraq

Syntypes: MNHNP 1935.231-35 (Palmyre), 1935.236 (Tell Abiad), 1935.237 (Ain Zahra), 1935.238 (Deir Ez Zor)

Synonym:

Acanthodactylus tristrami iracensis Schmidt, 1939 Type locality: Haditha, Iraq Holotype: FMNH 21679

Distribution: eastern Syria; western & central Iraq; northern Saudi Arabia

Acanthodactylus pardalis (Lichtenstein, 1823)

Lacerta pardalis Lichtenstein, 1823: 99

Type locality: Egypt

Lectotype: ZMB 1077, designated by Moravec, 1999

Synonym:

Lacerta deserti Milne-Edwards, 1829 Type locality: Near East Syntypes, MNHNP 5322(2).

Distribution: northeastern Mediterranean coast from the Gulf of Sirte, Libya, to the Nile Valley, Egypt

Acanthodactylus robustus Werner, 1929

Acanthodactylus robustus Werner, 1929: 240

Type locality: western Syria, near Bir Molusi (Ka'ra)

Holotype: NHMW 23362

Distribution: southwestern Iraq, Syria, Jordan, northwestern Saudi Arabia.

Acanthodactylus savignyi (Audouin, 1827)

Lacerta savignyi Audouin, 1827: 118

Type locality: Egypt

Holotype: MNHNP, not located

Synonyms:

Acanthodactylus vaillanti Lataste, 1885 Type locality: Comalis country, Algeria Holotype: MNHNP 0.6484

Acanthodactylus savignyi oranensis Doumergue, 1901 Type locality: Oran, on the sands of the Spanish Battery Syntypes: BMHN 1946.9.3.58-70; MNHNP 0.8956

Distribution: coastal Algeria

Acanthodactylus schmidti Haas, 1957

Acanthodactylus cantoris schmidti Haas, 1957: 72

Type locality: Dhahran, Saudi Arabia

Holotype: CAS 84599

Distribution: Saudi Arabia, Jordan, southeastern Iraq, southwestern Iran, United Arab Emirates, Kuwait, Oman, Qatar

Acanthodactylus schreiberi Boulenger, 1878

Acanthodactylus savignyi schreiberii Boulenger, 1878: 188

Type locality: Cyprus

Syntypes: IRSNB 2350

Synonyms:

Acanthodactylus schreiberi atauri Yalçinkaya & Güçmen, 2012 Type locality: Botaş-Adana, Turkey Holotype: ZDEU 6/1999-2.

Acanthodactylus boskianus syriacus Boettger, 1878 Type locality: Haifa, Israel

Lectotype: SMF 13633

Distribution: Mediterranean coasts: Lebanon & Israel

Acanthodactylus scutellatus (Audouin, 1827)

Acanthodactylus scutellatus audouini Boulenger 1918

Type locality: Wed El Kreil, Tunisia

Lectotype: BMNH 1920.1.20.3006, designated by Crochet et al., 2003

Distribution: northern Mali, Niger, northern Chad, southern Algeria, southern Tunisia, Libya, northern Sudan, southern Egypt

Acanthodactylus scutellatus scutellatus (Audouin, 1827)

Lacerta scutellata Audouin, 1827: 117

Type locality: Egypt

Holotype: lost

Distribution: Egypt (including Sinai); southern Israel

Acanthodactylus senegalensis Chabanaud, 1918

[Acanthodactylus senegalensis Chabanaud, 1918: 162](#)

Type locality: near Rufisque, Senegal

Holotype: MNHNP 1918.43

Distribution: Mauritania, Western Sahara, northern Senegal, Mali, southern Algeria

Acanthodactylus spinicauda Doumergue, 1901

[Acanthodactylus pardalis spinicauda Doumergue, 1901: 162](#)

Type localities: El Abiod-Sidi-Cheikh & Arba Tahtani, Algeria

Syntypes: BMNH 1946.8.5.34-43 (Arba Tahtani); MNHNP 0.8958 (El Abiod-Sidi-Cheikh)

Distribution: northwestern Algeria: southern base of Sahara-Atlas

Acanthodactylus taghitensis Geniez & Foucart, 1995

[Acanthodactylus taghitensis Geniez & Foucart, 1995: 7](#)

Type locality: 36 km SW Taghit (30°41' N, 2°07' W), Beni Abbès region, Algeria

Holotype: MNHNP 1995.1201

Distribution: along the border between Mauritania and Western Sahara and between Algeria and Morocco

Acanthodactylus tilburyi Arnold, 1986

[Acanthodactylus tilburyi Arnold, 1986: 378](#)

Type locality: Nafud as Sirr [26°11' N, 44°19'E], near Riyadh, Saudi Arabia

Holotype: BMNH 1985.1273

Distribution: central and northern Saudi Arabia; southern Jordan

Acanthodactylus tristrami (Günther, 1864)

[Zootoca tristrami Günther, 1864: 491](#)

Type locality: Lebanon

Holotype: BMNH 1946.9.3.73

Synonym:

Acanthodactylus dorsalis Peters, 1869 Type locality: unknown Syntypes: ZMB 1056-58

Distribution: eastern Lebanon, western Syria, western Iraq, western Jordan

Acanthodactylus yemenicus Salvador, 1982

[Acanthodactylus yemenicus Salvador, 1982: 53](#)

Type locality: Ta'izz, Yemen

Holotype: FMNH 66510 Distribution:

western Yemen

Adolfus Sternfeld, 1912 Sternfeld, R. 1912. Reptilia. pp. 197–279 + Plates VI-IX. In: Schubotz, H. (Editor).



1913. *Wissenschaftliche Ergebnisse der Deutschen Zentral-Afrika-Expedition 1907-1908 unter Führung Adolf Friedrichs, Herzogs zu Mecklenburg*, Band IV [Volume 4], Zoologie II. Leipzig: Klinkhardt & Biermann. 485 pp Type species: *Algiroides africanus* Boulenger, 1906

Adolfus jacksoni  Nicolás Lutzmann

Adolfus africanus (Boulenger, 1906)

[Algiroides africanus Boulenger, 1906: 570](#)

Type locality: Entebbe, Uganda

Holotype: BMNH 1946.9.3.13

Synonym:

Adolfus fridericianus Sternfeld, 1912 Type locality: Aruwimi,

DC Congo Holotype: ZMB 24121

Distribution: South Sudan; Kenya: Kakamega Forest; Uganda; western Rwanda; Democratic Republic of the Congo; northwestern Zambia; southern Cameroon; Central African Republic

Adolfus alleni (Barbour, 1914)

[Algyroides alleni Barbour, 1914: 97](#)

Type locality: near the tree limit, northeast slope of Mt. Kenya, Kenya

Holotype: MCZ R-9280

Distribution: Uganda: Mount Elgon; Kenya: Cherangani Hills, Mount Kenya, Aberdare Mountains

Adolfus jacksoni (Boulenger, 1899)

[Lacerta jacksoni Boulenger, 1899: 96](#)

Type locality: Ravine Station, Mau Mts., Kenya

Holotype: BMNH 1946.9.2.99

Distribution: southern Kenya; southern Uganda; western Rwanda; northwestern Burundi; Tanzania: Kilimanjaro region and west of Lake Victoria; eastern Democratic Republic of Congo

Adolfus kibonotensis (Lönnerberg, 1907)

[Lacerta jacksoni kibonotensis Lönnerberg, 1907: 5](#)

Type locality: Kibongoto, Kilimanjaro, Tanzania

Holotype: NRM, not located

Distribution: Kenya: Taita Hills, Mt. Kenya; Tanzania: West Usambara Mountains and Mt. Kilimanjaro

Adolfus masavaensis Wagner, Greenbaum & Branch, 2014

[Adolfus masavaensis Wagner, Greenbaum & Branch, 2014: 9](#)

Type locality: Mount Elgon (ca. 1.040983°N, 34.783645°E), Kenya, near Koitobos Guest House

Holotype: ZFMK 75001

Distribution: Kenya: Mount Elgon, Aberdares Mts., Cherangani Hills; Uganda: Mount Elgon

Adolfus mathewsensis Greenbaum, Beer, Wagner, Anderson, Villanueva, Malonza, 2018

[Adolfus mathewsensis Greenbaum, Beer, Wagner, Anderson, Villanueva, Malonza, 2018: 58](#)

Type locality: between Ng'eny forest and Kitich Camp (1.251408°N, 37.293328°E), Muger Base Camp, Mathews Range North, Samburu County, Rift Valley Province, Kenya.

Holotype: NMK L/3376/2

Distribution: Kenya: Mathews Range

Algyroides Bibron & Bory, 1833

Bibron, M. & J.P. Bory de Saint-Vincent. 1833. Description of *Podarcis*



peloponnesiacus and *Algyroides moreoticus* in Vertébrés a Vertébrés a sang froid. Reptiles et poissons. Reptiles. In: Geoffroy & Geoffroy, *Expédition Scientifique de Morée*, Tome III. 1re partie, Zoologie: pp. 57 Type species: *Algyroides moreoticus* Bibron & Bory, 1833

Algyroides nigropunctatus nigropunctatus 
Horst Kretzschmar

Algyroides fitzingeri (Wiegmann, 1834)

[Notopholis fitzingeri Wiegmann 1834: 10](#)

Type locality: Corsica, France

Syntypes: ZMB 910

Distribution: France Corsica; Italy: Sardinia

Algyroides hidalgoi Bosca, 1916

Type locality: Los Rasos, Cazorla, Cauce del Guadalquivir San Ildefonso, Segovia, Spain

Neotype: MNCN 32128

Synonyms:

Algyroides marchi marchi Valverde, 1958:127 Type locality: Piedra de Aguamula, Sierra de Cazorla, Jaén Province, Spain Holotype: Instituto de Aclimatación de Almería

Algyroides marchi niethammeri Buchholz, 1965: 244 Type locality: Sierra de Agua, Jaen Prov., Spain
Holotype: ZFMK 4541

Distribution: Spain: Jaén Province: Sierra de Cazorla, Sierra de Segura, Sierra de Alcaez.

Algyroides marchi niethammeri Buchholz, 1965: 244

Type locality: Sierra de Agua, Jaen Prov., Spain

Holotype: ZFMK 4541

Distribution: Spain: Sierra de Agua

Algyroides marchi marchi Valverde, 1958

Algyroides marchi Valverde, 1958: 127

Type locality: Pledra de Aguamula, Sierra de Cazorla, Jaén Province, Spain

Holotype: Instituto de Aclimatación de Almería

Distribution: Spain: Sierra de Cazorla, Sierra de Segura, Sierra de Alcaez.

Algyroides moreoticus Bibron & Bory, 1833

Algyroides moreoticus Bibron & Bory, 1833: 67

Type locality: Koubeh, Peloponnese, Greece

Holotype: BMNH 1946.9.3.12

Distribution: Greece: Peloponnese, and the three southern Ionian islands: Kephalliana, Ithaca, Zakynthos.

Algyroides nigropunctatus (Duméril & Bibron, 1839)

Algyroides nigropunctatus nigropunctatus (Duméril & Bibron, 1839)

Lacerta nigro-punctata Duméril & Bibron, 1839: 190

Type locality: Corfu Island, Greece

Holotype: MNHNP 0.2674

Distribution: northeastern Italy, southern Slovenia, coastal Croatia, river valleys in Bosnia-Herzegovina, southern Serbia, Montenegro, Albania, Macedonia, western Greece

Algyroides nigropunctatus kephallithacius Keymar, 1986

Algyroides nigropunctatus kephallithacius Keymar, 1986: ?

Type locality: Lefkada Island, Greece

Holotype: ?

Distribution: Greece: Lefkada, Kephalliana, Ithaca Islands

Anatololacerta Arnold, Arribas & Carranza, 2007

Arnold, E.N., O. Arribas & S. Carranza. 2007.



Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions

of eight new genera. *Zootaxa* 1430: 1-86

Type species: *Zootoca danfordi* Günther, 1876

Anatololacerta anatolica aegaea  Joachim Rutschke

Anatololacerta anatolica (Werner, 1900)

Anatololacerta anatolica anatolica (Werner, 1900)

Lacerta anatolica Werner, 1900: 269

Type locality: Gökçekisik, Eskişehir province, Turkey

Holotype: NHMW 15120

Distribution: Turkey: northwestern Anatolia; Greece: Samos and Ikaria Islands

Anatololacerta anatolica aegaea Eiselt & Schmidtler, 1987

[Lacerta anatolica aegaea Eiselt & Schmidtler, 1987: 312](#)

Type locality: Samsundağ, S Davultar, Aydın Province, Turkey

Holotype: ZSM 50/1983

Distribution: Turkey: Aydın Province; Greece: Samos Island

Anatololacerta anatolica oertzeni (Werner, 1904)

[Lacerta oertzeni Werner, 1904: 313](#)

Type locality: Maratrokambos, off Sambos, Greece

Syntype: ZMB 13495

Distribution: Greece: Ikaria Island

Anatololacerta budaki (Eiselt & Schmidtler, 1987)

[Lacerta oertzeni budaki Eiselt & Schmidtler, 1987: 316](#)

Type locality: 5 km N Cobanisa (= 25 km NE Elmali), Vilayet Anta-Hlya, Turkey

Holotype: NHMW 26751

Distribution: Turkey: southwestern Anatolia, from ENE of Kas and Karaman Pass to Antalya

Anatololacerta danfordi (Günther, 1876)

Anatololacerta danfordi danfordi (Günther, 1876)

[Zootoca danfordi Günther, 1876: 818](#)

Type locality: Zebil Bulgar Dagh (Cilician Taurus), Turkey

Syntypes: BMNH 1946.9.2. 34-41

Synonym:

Lacerta danfordi quanttaylori Börner 1974 Type locality: ? Holotype: ?

Distribution: Turkey: mountain ranges around the Bolkar Dag.

Anatololacerta danfordi bileki Eiselt & Schmidter, 1986

[Lacerta danfordi bileki Eiselt & Schmidter, 1986: 323](#)

Type locality: Irmansan-Pas, Vilayet, Antayai, Turkey

Holotype: NHMW 26759

Distribution: Turkey: interior south coast

Anatololacerta pelasgiana (Mertens, 1959)

Anatololacerta pelasgiana pelasgiana (Mertens, 1959)

[Lacerta oertzeni pelasgiana Mertens, 1959: 18](#)

Type locality: wharfside, Rhodes Island, Greece

Holotype: SMF 55228

Synonym:

Lacerta danfordi pentanisiensis Wettstein, 1964 Type locality: Largest island of the Pentanisos group, east of St. Nikolas at Lindos, east coast of Rhodes, Greece Holotype: NHMW 18247:1

Distribution: Turkey: western coast, between Fethiye and Milas; Greece: Rhodes Island and surrounding islands

Anatololacerta pelasgiana finikensis (Eiselt & Schmidtler, 1986)

[Lacerta oertzeni finikensis Eiselt & Schmidtler, 1987: 316](#)

Type locality: ca. 3 km SW Finike, Vilayet Antalya, at the grotto above the coastal road, Turkey

Holotype: NHMW 18355:7

Distribution: southwestern Turkey, between Kas and Finike

Anatololacerta pelasgiana ibrahimi (Eiselt & Schmidtler, 1986)

Lacreta oertzeni ibrahimi Eiselt & Schmidtler, 1986: 322

Type locality: ca. 20 km NW Anamur, Mersin Province, Turkey

Holotype: ZSM 59/83

Distribution: Turkey: central south coast

Apathya Méhely, 1907 Méhely, Lajos von. 1909. Materialien zu einer Systematik und Phylogenie der



muralis-ähnlichen Lacerten. *Annales historico-naturales Musei nationalis Hungarici*, Budapest, 7 (2): 409-621
Type species: *Lacerta cappadocica* Werner, 1902

Apathya cappadocica urmiana Omid Mozaffari

Apathya cappadocica (Werner, 1902)

Apathya cappadocica cappadocica (Werner, 1902)

Lacerta cappadocica Werner, 1902: 1086

Type locality: Erciyes Massif, near Kayseri, central Anatolia, Turkey

Syntype: ZMH R04403

Distribution: Kilikian Taurus to west of Malatya, Turkey

Apathya cappadocica muhtari (Eiselt, 1979)

Lacerta cappadocica muhtari Eiselt, 1979: 413

Type locality: 26 km SW Bitlis, Turkey

Holotype: NHMW 23208:2

Distribution: east of Malatya and the Euphrates to the south of Lake Van, Turkey; northeastern Iraq: to Sersang.

Apathya cappadocica schmidtlerorum (Eiselt, 1979)

Lacerta cappadocica schmidtlerorum Eiselt, 1979: 15

Type locality: 10 km S Diyarbakir, Abzweigung Ovabag, Turkey

Holotype: ZSM 252/77,

Distribution: Turkey: around Diyarbakir and Viransehir.

Apathya cappadocica urmiana (Lantz & Suchow, 1934)

Lacerta cappadocica urmiana Lantz & Suchow, 1934: ?

Type locality: Kherra, gorge of the Berdesur River, ca. 20 km SW Rezaiyeh, Iran

Holotype: ZISP 12657

Distribution: southeastern Turkey; northeastern Iraq; northwestern Iran

Apathya cappadocica wolteri (Bird, 1936)

Lacerta cappadocica wolteri Bird, 1936: 266

Type locality: ten miles west of Gaziantep; on limestone rocks in thick scrub country, southern Turkey

Holotype: BMNH 1935. 11. 4. 128

Distribution: southern Turkey; northwestern Syria

Apathya yassujica (Nilson, Rastegar-Pouyani, Rastegar-Pouyani & Andr n, 2003)

Lacerta yassujica Nilson, Rastegar-Pouyani, Rastegar-Pouyani & Andr n, 2003: 18

Type locality: Type locality: main Zagros Mountain Range (30°28'N; 51°31'E), 30 km SW Yassuj, Kohgiluyeh-va-Boyer-Ahmad Province, Iran

Holotype: NHMG 5612

Distribution: Iran: southern and central Zagros Mountains

Archaeolacerta Mertens, 1921 Mertens, R. 1921. Zur Kenntnis der Reptilienfauna von Malta. *Zoologischer*

Anzeiger 53: 236-240 Type species:

Lacerta reticulata Bedriaga, 1881



Archaeolacerta bedriagae bedriagae 

Wolfram Schurig

Archaeolacerta bedriagae (Camerano, 1885)

Archaeolacerta bedriagae bedriagae (Camerano, 1885)

Lacerta oxycephala bedriagae Camerano, 1885: 418

Type locality: Corsica

Syntypes: BMNH 1946.9.1.46-47

Distribution: Corsica, France

Archaeolacerta bedriagae paessleri (Mertens, 1927)

Lacerta bedriagae paessleri Mertens, 1927: 179

Type locality: Mt. Limbara, Sardinia, Italy

Holotype: SMF 22086

Distribution: Italy: Sardinia: Limbara Mountains

Archaeolacerta bedriagae ferrerae Stemmler, 1961

Lacerta bedriagae ferrerae Stemmler, 1961: 21

Type locality: Bateria Ferrera, Gallura, northern Sardinia

Holotype: NMB 15272

Distribution: Italy: northern Sardinia

Archaeolacerta bedriagae sardoa (Peracca, 1903)

Lacerta sardoa Peracca, 1905: 1

Type locality: Punta Paulinu, Monti del Gennargentu, Sardinia

Holotype: MRSN R2125

Distribution: Italy: central and southern Sardinia

Atlantolacerta Arnold, Arribas & Carranza, 2007 Arnold, E.N., O. Arribas & S. Carranza. 2007.



Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata:

Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* 1430: 1-86

Type species: *Lacerta andreanskyi*

Werner, 1929

Atlantolacerta andreanskyi Daniel Escoriza

Atlantolacerta andreanskyi (Werner, 1929)

Lacerta andreanskyi Werner, 1929: 4

Type locality: Tachdirt, Grand Atlas, Morocco

Holotype: MCZ R-27391

Distribution: Morocco: High Atlas Mountains

Australolacerta Arnold, Arribas & Carranza, 2007 Arnold, E.N., O. Arribas & S. Carranza. 2007.



Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata:

Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* 1430: 1-86

Type species: *Lacerta australis* Hewitt,

1926

Australolacerta australis A Rebelo

Australolacerta australis (Hewitt, 1926)

Lacerta australis Hewitt, 1926: 423

Type locality: Matroosberg, Cape Province, South Africa

Holotype: SAM 14741

Distribution: South Africa: southwestern Cape Fold Mountains, Western Cape Province

Congolacerta Greenbaum, Villanueva, Kusamba, Aistote, & Branch, 2011

Greenbaum, E.,



C.O. Villanueva, C. Kusamba, M.M. Aristote and W.R. Branch. 2011. A molecular phylogeny of Equatorial African Lacertidae, with the description of a new genus and species from eastern Democratic Republic of the Congo. *Zoological Journal of the Linnean Society* 163: 913–942 Type species: *Lacerta vauereselli* Tornier, 1902

Congolacerta vauereselli Michele Menegon

Congolacerta asukului Greenbaum, Villanueva, Kusamba, Aristote & Branch, 2011

Congolacerta asukului Greenbaum, Villanueva, Kusamba, Aristote & Branch, 2011: 928

Type locality: footpath south of Rurambo village (2°59'39"S, 28°52'34"E) , Itombwe Plateau, South Kivu Province, DR Congo

Holotype: UTEP 20263

Distribution: Itombwe Plateau, South Kivu Province, DR Congo

Congolacerta vauereselli (Tornier, 1902)

Lacerta vauereselli Tornier, 1902: 701

Type locality: Kagera, west of Lake Victoria, Tanzania

Holotype: ZMB 17700


Synonym:

Algiroides boulengeri Peracca, 1917 Type locality: Fort Portal, Toro, western Uganda Holotype: MRSN ?

Distribution: southern Uganda; northwestern Tanzania; Rwanda; Burundi; northeastern Democratic Republic of Congo.

Dalmatolacerta Arnold, Arribas & Carranza, 2007

Arnold, E.N., O. Arribas & S. Carranza. 2007.



Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* 1430: 1-86
Type species: *Lacerta oxycephala* Duméril & Bibron, 1839

Dalmatolacerta oxycephala  Tekla Boersma

Dalmatolacerta oxycephala (Duméril & Bibron, 1839)

Lacerta oxycephala Duméril & Bibron, 1839: 235

Type locality: Dalmatia, Croatia

Holotype: MNHNP 0.2738

Synonym:

Lacerta oxycephala tomasinii Schreiber, 1891 Type locality: Herzegovina Holotype: NHMW 16234

Distribution: southern Croatia, Bosnia-Herzegovina, Montenegro, northwestern Albania

Darevskia Arribas, 1997

Arribas, O. J. 1997. Morfología, filogenia y bibliografía de las lagartijas de alta montana de los Pirineos [Microforma].



Tesis doctoral - Universitat Autònoma de Barcelona. *Publicaciones de la Universitat Autònoma de Barcelona*, 08193 Bellaterra (Barcelona), 353 pp. Type species: *Lacerta saxicola* Eversmann, 1834

Darevskia caspica  N. Moradi

Darevskia adjarica (Darevsky & Eiselt, 1980)

Lacerta parvula adjarica Darevsky & Eiselt, 1980: 35

Type locality: Abastumani (41°44'N, 42°51'E), Samtskhe-Javakheti, Georgia.

Holotype: ZISP 17740

Distribution: Georgia; northeastern Turkey

Darevskia alpina (Darevsky, 1967)

Lacerta caucasica alpina Darevsky, 1967: 118

Type locality: Kabardino-Balkaria (43°15'N, 42°30'E), vicinity of Terskol Village, Elbrussky District, southern Russia

Holotype: ZISP 17942

Distribution: Great Caucasus Range of Georgia and Russia

Darevskia aghasyani Tuniyev and Petrova, 2019

Darevskia aghasyani Tuniyev and Petrova, 2019: 141

Type locality: Armenia, Ararat District, crest of Urts Ridge

Holotype: SNP 1770B

Distribution: Armenia: Ararat region: Khosrov Forest

Darevskia armeniaca (Méhely, 1909)

Lacerta saxicola armeniaca Méhely, 1909: 549

Type locality: coast of the Lake Sevan, Armenia

Lectotype: SMF 12066

Synonym:

Lacerta muralis chalybdea Boulenger, 1904 Type locality: Tbilissi, Georgia Type: ?

Distribution: northeastern Turkey, northwestern Armenia, western Azerbaijan, southern Georgia. Ukraine (introduced)

Darevskia bendimahiensis (Schmidtler, Eiselt & Darevesky, 1994)

Lacerta bendimahiensis Schmidtler, Eiselt & Darevesky, 1994: 62

Type locality: Bendimahi Waterfall, 9 km N Muradiye, Provi. Van, Turkey

Holotype: ZSM 169/1993

Distribution: eastern Turkey: northeast of Lake Van

Darevskia bithynica (Méhely, 1909)

Darevskia bithynica bithynica (Méhely, 1909)

Lacerta saxicola bithynica Méhely, 1909: 495

Type locality: Bithynian Olympus at Brussa, Turkey

Holotype: MZUT 2737:1

Distribution: northwestern Turkey: Ulu Dağ south of Bursa

Darevskia bithynica tristis (Lantz & Cyréen, 1936)

Lacerta saxicola tristis Lantz & Cyréen, 1936: 176

Type locality: Lafaka Déré, near Adapazar, Turkey

Holotype: ?

Distribution: Turkey: western Black Sea coast

Darevskia brauneri (Méhely, 1909)

Darevskia brauneri brauneri (Méhely, 1909)

Lacerta saxicola brauneri Méhely, 1909: 493

Type locality: Krasnaya Polyana in SW Caucasus, Russia

Lectotype: ZISP 16352

Synonyms:

Lacerta saxicola darevskii Szczerbak, 1962 Type locality: Kamennomostsky Village (44°17'N 40°11'E), Maikop Region, Republic of Adygea, Russian Federation Holotype: NMNH Re 1

Lacerta saxicola szczerbaki Lukina, 1963 Type locality: the coastal cliff at 1.5 km south-east of the Lidzava (= Ldzaa) Village (43°9'N 40°25'E), Pitsunda-Myusserky Reserve, Gudauta District, Republic Abkhazia, Russia Holotype: ?

Distribution: Russia: western Caucasus foothills; northwestern Georgia.

Darevskia brauneri myusserica Doronin, 2011

Darevskia brauneri myusserica Doronin, 2011: 248

Type locality: cliff at 1,5 km south-east of the village Lidzava (= Ldzaa) (43°9'N, 40°25'E), Pitsunda-Myussera Reserve, Gudauta District, Republic of Abkhazia, Republic of Georgia

Holotype: ZISP 25964

Distribution: Georgia Republic: Abachasia, Pitsunda-Reserve and Myussera Gagra

Darevskia caspica Ahmadzadeh, Flecks, Carretero, Mozaffari, Böhme, Harris, Freitas & Rödder, 2013

Darevskia caspica Ahmadzadeh, Flecks, Carretero, Mozaffari, Böhme, Harris, Freitas & Rödder, 2013: 9

Type locality: Beliroon (36° 23'38'' N, 52° 25'1.48''E), Amol, Mazandaran Province, Iran

Holotype: ZFMK 94109

Distribution: Iran: Mazandaran Province: Hyrcanian forest

Darevskia caucasica (Méhely, 1909)

Darevskia caucasica caucasica (Méhely, 1909)

Lacerta caucasica Méhely, 1909: 560

Type locality: Mount Kazbek (42°40' N 44°35' E), Kazbegi Municipality, Mtskheta-Mtianeti, Georgia

Lectotype: SMF 12069

Distribution: Caucasus Mountains: Russia, Georgia, northeastern Azerbaijan

Darevskia caucasica vedenica (Darevsky & Roitberg, 1999)

Lacerta caucasica vedenica Darevsky & Roitberg, 1999: 209

Type locality: 1 km south of Vedeno rock faces along the road Vedeno –Kharochoi, southeastern Chechen Republic

Holotype: ZISP 17744(1)

Distribution: Russia: southeastern Chechnya, Ingushetia

Darevskia chlorogaster (Boulenger, 1908)

Lacerta chlorogaster Boulenger, 1908: 934

Type locality: Bandar-e Anzali, Gilan Province, Iran

Lectotype: BMNH 1946.9.2.29

Synonym:

Lacerta mostoufii Baloutch, 1976 Type locality: Deh Salm (31°12'N, 59°19'E), about 200 km N Malek Mohammad Mountains Holotype: MMTT 1582

Distribution: southern Azerbaijan and northwestern Iran: eastern Hyrcanian Forest

Darevskia clarkorum (Darevsky & Vedmederja, 1977)

Lacerta clarkorum Darevsky & Vedmederja, 1977: 50

Type locality: Cankurtaran Pass between Hopa and Borçka, Artvin province, northeastern Turkey

Holotype: CAS 105610

Synonym:

Lacerta dryada Darevsky & Tuniyev, 1997 Type locality: Adzharistan, Khelvachaury district, gorge of Charnali, W Republic of Georgia. Holotype: CNR 1103

Distribution: northeastern Turkey: northern Pontic Mountains; southwestern Georgia: western slopes Mtirala Mountains

Darevskia daghestanica (Darevsky, 1967)

Lacerta saxicola daghestanica Darevsky, 1967: 55

Type locality: Levaschi (=Lavaschi) Village (42°25' N 47°19' E), Levashinsky District, Dagestan, Russia

Lectotype: GNM 263

Synonym:

Lacerta saxicola gracilis Méhely, 1909 Type locality: Dagestan, Russia Syntypes: MC 26

Lacerta muralis caucasica Boulenger, 1913 Type locality: Pyatigorsk, Stavropol Krai, Russia Syntype: 1904.11.26.12

Distribution: northern Caucasus Range: southern Russia, northern Azerbaijan

Darevskia dahli (Darevsky, 1957)

Lacerta saxicola dahli Darevsky, 1957: 32

Type locality: near Shagali, northern Armenia

Holotype: ZIA 892

Distribution: northern Armenia, southern Georgia

Darevskia defilippii (Camerano, 1877)

Lacerta saxicola defilippii Camerano, 1877: 90

Type localities: Lar Valley and Damavand Mountain, Tehran, northern Iran

Syntypes: MRSN R2734, R2713

Synonym:

Lacerta muralis persica Bedriaga, 1886 Type locality: ? Type: ?

Distribution: northern Iran: western Alborz Mountains

Darevskia derjugini (Nikolsky, 1898)

Darevskia derjugini derjugini (Nikolsky, 1898)

Lacerta derjugini Nikolsky, 1898: 284

Type locality: Artvin, Turkey

Syntypes: ?? 9101-02

Distribution: Georgia; northeastern Turkey

Darevskia derjugini abchasica Bischoff, 1982

Lacerta derjugini abchasica Bischoff, 1982: 44 Type locality: Adzharia, Batumi, Georgia

Holotype: MM ?

Distribution: Georgia: vicinity of Sukhumi

Darevskia derjugini barani (Bischoff, 1982)

Lacerta derjugini barani Bischoff, 1982: 32

Type locality: Botanical Garden of Batumi, Georgia

Holotype: MM ?

Distribution: northeastern Turkey: coastal mountains of the Black Sea

Darevskia derjugini boehmei (Bischoff, 1982)

Lacerta derjugini boehmei Bischoff, 1982: 46

Type locality: Mt. Achun, Sochi, Russia

Holotype: MTKD 14626

Distribution: southwestern Caucasus Range: Russia and Georgia

Darevskia derjugini orlowae (Bischoff, 1984)

Lacerta derjugini orlowae Bischoff, 1984: 104

Type locality: near Pasanauri, Georgia

Holotype: ZFMK 38047

Distribution: southern Great Caucasus: Georgia and northwestern Azerbaijan.

Darevskia derjugini silvatica (Bartenjew & Rjesbikowa, 1931)

Lacerta derjugini silvatica Bartenjew & Rjesnikowa, 1931: 9

Type locality: Glade stock well on the river Belaya, northwest Caucasus, Russia

Holotype: ?

Distribution: Russia: northwestern Caucasus

Darevskia kamii Ahmadzadeh, Flecks, Carretero, Mozaffari, Böhme, Harris, Freitas & Rödder, 2013

Darevskia kamii Ahmadzadeh, Flecks, Carretero, Mozaffari, Böhme, Harris, Freitas & Rödder, 2013: 11

Type locality: Naharkhoran Forest (36° 46'33.61''N, 54°27'48.01''E), Gorgan, Golestan Province, Iran

Holotype: ZFMK 94118

Distribution: Iran: Golestan Province: western Hyrcanian forest

Darevskia kopetdaghica Ahmadzadeh, Flecks, Carretero, Mozaffari, Böhme, Harris, Freitas & Rödder, 2013

Darevskia kopetdaghica Ahmadzadeh, Flecks, Carretero, Mozaffari, Böhme, Harris, Freitas & Rödder, 2013: 12

Type locality: Sarani Protected Area (37°44'4.96''N, 58°5'25.09''E), Kopet Dagh, northern Khorasan Province, Iran

Holotype: ZFMK 94124

Distribution: Kopet Dagh Range: northern Iran, southern Turkmenistan

Darevskia lindholmi (Szczerbak, 1962)

Lacerta saxicola lindholmi Szczerbak, 1962: 1378

Type locality: Yalta (44°29'N, 34°9'E), Autonomous Republic of Crimea, Ukraine

Lectotype: NMNH SR 2057/14097

Distribution: Ukraine: Crimea mountains

Darevskia mixta (Méhely, 1909)

Lacerta mixta Méhely, 1909: 581

Type locality: Abastumani, Republic of Georgia

Holotype: SMF 12087

Distribution: northeastern Turkey; southwestern Georgia

Darevskia parvula (Lantz & Cyrén, 1913)

Lacerta saxicola parvula Lantz & Cyrén, 1913: ?

Type locality: Artvin, northeastern Turkey

Lectotype: GNM 2862

Distribution: northeastern Turkey: Çoruh River Valley

Darevskia pontica (Lantz & Cyrén, 1919)

Lacerta praticola pontica Lantz & Cyrén, 1919: 30

Type locality: Gagry, on the Black Sea coast, Republic of Georgia

Lectotype: ZISP 22853

Distribution: western Caucasus Range: northwestern Georgia, southwestern Russia

Darevskia portschinskii (Kessler, 1878)

Darevskia portschinskii portschinskii (Kessler, 1878)

Lacerta portschinskii Kessler, 1878: 160

Type locality: Tbilisi, Republic of Georgia

Holotype: MNKNU 27226

Distribution: southeastern Georgia, Armenia, southern Azerbaijan

Darevskia portschinskii nigrita (Bakradse, 1976)

Lacerta portschinskii nigrita Bakradse, 1976: 54

Type locality: Dmanisi, Marneuli district, Georgia

Holotype: ZISP 18604

Distribution: Georgia: Mashavera Valley; Armenia: Debeta Valley

Darevskia praticola (Eversmann, 1834)

Darevskia praticola praticola (Eversmann, 1834)

Lacerta praticola Eversmann, 1834: 345

Type locality: Kislovodsk, southern Russia

Lectotype: ZISP 22848

Synonyms:

Lacerta vivipara stenolepis Nikolsky, 1911 Type locality: unknown Holotype: ZISP 7203

Lacerta plicata Bartenjew & Reznikova, 1931 Type locality: firebreak in streams in the small mountain river on the sunny glade (43° 53' N, 40° 31' E), on the road from the Veriyut (=Nikitino Village) on the Malaya Laba River to the Malyi Pambak Mount, Caucasian Biosphere Reserve, Mostovskoy District, Russia Holotype: ZISP 15204

Distribution: Azerbaijan, Georgia, Russia

Darevskia praticola hungarica (Sobolewsky, 1930)

Lacerta praticola hungarica Sobolewsky, 1930: 6

Type locality: Mehadia (44°54' N, 22°22' E), Caraş-Severin County, Romania

Lectotype: ZISP 9814

Distribution: northeastern Serbia, southern Romania, northern Greece, Bulgaria, European Turkey

Darevskia praticola hyrcanica Tuniyev, Doronin, Kidov & Tuniyev, 2011

Darevskia praticola hyrcanica Tuniyev, Doronin, Kidov & Tuniyev, 2011: 304

Type locality: Talysh Ridge, natural boundary of Gada-Zyga-Khi (38°28' N, 48°35' E), Astara District, Azerbaijan

Holotype: SNP 1473

Distribution: southern Azerbaijan, northeastern Iran

Darevskia praticola loriensis Tuniyev, Doronin, Tuniyev, Aghasyan, Kidov & Aghasyan, 2013

Darevskia praticola loriensis Tuniyev, Doronin, Tuniyev, Aghasyan, Kidov & Aghasyan, 2013: 229

Type locality: village Gyulagarak (40°56'35.0'' N, 44°28'19.1'' E), vicinity of Stepanavan, Lori Marz (Province), Armenia

Holotype: SNP 1568

Distribution: northern Armenia

Darevskia raddei (Boettger, 1892)

Darevskia raddei raddei (Boettger, 1892)

Lacerta saxicola raddei Boettger, 1892: 142

Type locality: Aras River Valley, SE Armenia

Lectotype: SMF 12054

Synonym:

Lacerta caucasica tenuis Nikolsky, 1910 Type locality: Khân Bolâghî, Azerbaijan

Syntype: MNKNU 27252

Distribution: Armenia, southern Azerbaijan, northwestern Iran

Darevskia raddei nairensis (Darevsky, 1967)

Lacerta saxicola nairensis Darevsky, 1967: 68

Type locality: Lchashen, beach of Lake Sevan,

Armenia Holotype: ZISP 17941

Distribution: western Armenia; southern Georgia; northeastern Turkey

Darevskia raddei vanensis (Eiselt, Schmidler & Daversky, 1993)

Lacerta raddei vanensis Eiselt, Schmidler & Daversky, 1993: 66

Type locality: Burgberg, Van Province, Turkey

Holotype: NHMW 32999

Distribution: Turkey: eastern Lake Van basin

Darevskia raddei chaldoranensis Rastegar-Pouyani, Karamiani, Orae, Khosrawani & Rastegar-Pouyani, 2011

Darevskia raddei chaldoranensis Rastegar-Pouyani, Karamiani, Orae, Khosrawani & Rastegar-Pouyani, 2011: 217

Type locality: Chaldoran area (39° 03' N, 44° 22' E) in the northern regions of west Azerbaijan Province, Iran.

Holotype: RUZM- LL70.1

Distribution: Iran: western Azerbaijan Province: Chaldoran area

Darevskia rostombekowi (Darevsky, 1957)

Lacerta saxicola rostombekowi Darevsky, 1957: 35

Type locality: near Idzhevan, northern Armenia

Holotype: ZIA 900

Distribution: northern Armenia; western Azerbaijan

Darevskia rudis (Bedriaga, 1886)

Darevskia rudis rudis (Bedriaga, 1886)

Lacerta depressa rudis Bedriaga, 1886: 275

Type locality: Trabzon, Turkey

Holotype: MZUT 2737:1

Distribution: northeastern Turkey, near the Black Sea

Darevskia rudis bischoffi (Böhme & Bubak, 1977)

Lacerta rudis bischoffi Böhme & Bubak, 1977

Type locality: Artvin village, 6 km W Arhavi, eastern Turkey

Holotype: SZE 183/977

Distribution: Georgia: southern Black Sea coast; Turkey: northeastern coast

Darevskia rudis bolcardaghica Arribas, Ilgaz, Kumlutaş, Durmuş, Avcı & Üzümlü, 2013

Darevskia rudis bolcardaghica Arribas, Ilgaz, Kumlutaş, Durmuş, Avcı & Üzümlü, 2013: 422

Type locality: Karagöl, Ulukışla, Niğde, Central Anatolia, Turkey

Holotype: ZDEU 144/2009

Distribution: Turkey: south-central Anatolia

Darevskia rudis chechenica (Eiselt & Darevsky, 1991)

Lacerta rudis chechenica Eiselt & Darevsky, 1991: 17

Type locality: canyon of the Argun River, northern gorge of the central Caucasus, Chechen Republic

Holotype: ZISP 17882
Distribution: Russia, Georgia, Azerbaijan

Darevskia rudis macromaculata (Darevsky, 1967)

Lacerta rudis macromaculata Darevsky, 1967: 104
Type locality: vicinity of Akhalkalaki, Georgia
Holotype: ZISP 17940
Distribution: Kura Valley: Georgia, Turkey, Armenia

Darevskia rudis mirabilis Arribas, Ilgaz, Kumlutaş, Durmuş, Avci & Üzümlü, 2013

Darevskia rudis mirabilis Arribas, Ilgaz, Kumlutaş, Durmuş, Avci & Üzümlü, 2013: 421
Type locality: Ovit Pass, Rize (40°37' N, 40°49' E), Trabzon, northeastern Anatolia, Turkey
Holotype: ZDEU 145/2002
Distribution: Turkey: northeastern Anatolia

Darevskia rudis obscura (Lantz & Cyrén, 1936)

Lacerta saxicola obscura Lantz & Cyrén, 1936: 177
Type locality: valley of the Borjomka River, near Borjom, Georgia
Lectotype: ZISP 17171.1
Distribution: southern Georgia; northeastern Turkey

Darevskia rudis svanetica (Darevsky & Eiselt, 1980)

Lacerta rudis svanetica Darevsky & Eiselt, 1980: 32
Type locality: Mestia in Svanetia (43°03'N, 42°45'E), Georgia
Holotype: ZISP 17875: 1.
Distribution: Great Caucasus Range: Georgia: Inguri Valley; Russia: Mount Atsetuk

Darevskia sapphirina (Schmidtler, Eiselt & Darevsky, 1994)

Lacerta sapphirina Schmidtler, Eiselt & Darevsky, 1994: 55
Type locality: 30 km NW Erçis, Van Province, Turkey
Holotype: ZSM 170/1993
Distribution: Turkey: north of Lake Van

Darevskia saxicola (Eversmann, 1834)

Lacerta saxicola Eversmann, 1834: 349
Type locality: southern slope of the Borgustan Ridge (42°44' N, 43°57' E), Kislovodsk, Stavropol Territory, Russia
Neotype: ZISP 25731
Distribution: Russia: northern slopes of the Caucasus; Stavropol plateau

Darevskia schaeckeli Ahmadzadeh, Flecks, Carretero, Mozaffari, Böhme, Harris, Freitas & Rödder, 2013

Darevskia schaeckeli Ahmadzadeh, Flecks, Carretero, Mozaffari, Böhme, Harris, Freitas & Rödder, 2013: 12
Type locality: Firouzkooh (52° 449 48.580' E, 35°449 54.560' N), Tehran Province, Iran
Holotype: ZFMK 94200
Distribution: northern Iran: Alborz Mountains

Darevskia steineri (Eiselt, 1995)

Lacerta steineri Eiselt, 1995: 63
Type locality: Gole-Loweh, near Minou-dasht (37°20'53"N, 55°40' 28"E), SE Gonbad-e-Gavous, northeastern Iran

Holotype: NHMW 33715
Distribution: Iran: Golestan Province

Darevskia unisexualis (Darevsky, 1966)

Lacerta unisexualis Darevsky, 1966: 148

Type locality: Achta, in the canyon of the Razdan River, Armenia

Holotype: ZISP 17929

Distribution: Armenia, northeastern Turkey, southern Georgia

Darevskia uzzelli (Darevsky & Danielyan, 1977)

Lacerta uzzelli Darevsky & Danielyan, 1977: 55

Type locality: Kars, Turkey

Holotype: CAS 105689

Distribution: Turkey: Erzurum & Kars Provinces

Darevskia valentini (Boettger, 1892)

Darevskia valentini valentini (Boettger, 1892)

Lacerta muralis valentini Boettger, 1892: 145

Type locality: Bazarkent, northeastern Armenia

Holotype: SMF 12064

Synonym:

Lacerta saxicola terentjevi Darevsky, 1957 Type locality: Verin Ghukasyan, Gukasyan district, Aremnia Holotype: ZISP 16677

Distribution: Azerbaijan; Armenia; northeastern Turkey: Kars Plateau; southern Georgia

Darevskia valentini lantzicyreni (Darevsky & Eiselt, 1967)

Lacerta saxicola lantzicyreni Darevsky & Eiselt, 1967: 107

Type locality: Mount Erciyes, Turkey

Lectotype: NHMG 2459

Distribution: Turkey: eastern & central mountains ranges

Darevskia valentini spitzenbergerae (Eiselt, Darevsky & Schmidtler, 1992)

Lacerta valentini spitzenbergerae Eiselt, Darevsky & Schmidtler, 1992: 13

Type locality: Mergan Yayla, Cilo-Sat Mountains, Turkey

Holotype: NHMW 32224

Distribution: southeastern Turkey: Hakkari. Province: Cilo-Sat Mountains.

Dinarolacerta Arnold, Arribas & Carranza, 2007 Arnold, E.N., O. Arribas & S. Carranza. 2007.



Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata:

Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* 1430: 1-86

Type species: *Lacerta mosorensis*

Kolombatovic, 1886

Dinarolacerta mosorensis Benny Trapp

Dinarolacerta montenegrina Ljubisavljevic, Arribas, Dukic & Carranza, 2007

[Dinarolacerta montenegrina Ljubisavljevic, Arribas, Dukic & Carranza, 2007: 11](#)

Type locality: Ebeza Mt., Prokletije Mountain Massif (42° 37'N, 19°33'E), Montenegro.

Holotype: G 21594

Distribution: Montenegro: Prokletije Mountains

Dinarolacerta mosorensis (Kolombatovic , 1886)

[Lacerta mosorensis Kolombatovic, 1886: 26](#)

Type locality: Mount Mosor, Dalmatia, Croatia

Syntypes: NHMW 16090:1,2

Distribution: Croatia, Bosnia-Herzegovina, Montenegro, Albania

Eremias Wiegmann, 1834

Wiegmann, A.F.A. 1834. *Herpetologica Mexicana, seu descriptio amphibiorum*



Novae Hispaniae. Pars Prima, Saurorum Species, Berlin: C.G. Lüderitz. vi + 54 pp. + Plates I-X. Type species: *Lacerta variabilis* Pallas, 1814

Eremias suphani  Sergé Bogaerts

Eremias acutirostris (Boulenger, 1887)

[Scapteira acutirostris Boulenger, 1887: 114](#)

Type locality: near Nushki, northern Balochistan, Pakistan

Holotype: BMNH 1946.8.7.46

Synonym:

Scapteira aporosceles Alcock & Finn, 1897 Type locality: west of Robat Island, northern Balochistan, Pakistan Syntypes: BMNH 1917.3.6.40, .43; ZSI 14132, 14134, 14137, 14139–42, 14144–45, 14147–51, 14153

Distribution: Iran: Sistan basin; southern Afghanistan; Pakistan: Registan Desert

Eremias afghanistanica Böhme & Scerbak, 1991

[Eremias afghanistanica Böhme & Scerbak, 1991: 137](#)

Type locality: Dasht-e-Nawar, Ghazni Province, eastern Afghanistan

Holotype: ZFMK 8584

Distribution: eastern Afghanistan

Eremias andersoni Darvesky & Szczerbak, 1978

Eremias andersoni Darvesky & Szczerbak, 1978: 13

Type locality: Dashte-Kavir Desert, 45 km E of Lake Daryache Namak, Iran

Holotype: MMTT 1671

Distribution: central Iran: Dasht-e-Kavir Desert

Eremias argus Peters, 1869

Eremias argus argus Peters, 1869

Eremias argus Peters, 1869: 61

Type locality: Yantai Shi, Shandong, China

Lectotype: ZMB 4532

Distribution: northeastern China, eastern Mongolia, North Korea, South Korea

Eremias argus barbouri Schmidt, 1925

Eremias barbouri Schmidt, 1925: 2

Type locality: Mai Tai Chao, northern Shansi, China

Holotype: AMNH 24045

Distribution: Russia: south bank of the Lake Baikal; central Mongolia; China: Inner Mongolia

Eremias arguta (Pallas, 1773)

Eremias arguta arguta (Pallas, 1773)

Lacerta arguta Pallas, 1773: 718

Type locality: between Ural River and Emba River, northwestern Kazakhstan

Neotype: ZISP 13205

Synonyms:

Lacerta variabilis Pallas, 1814 Type locality: Siberia, Russia Holotype: ZMB 1102

Lacerta pustulata Eichwald, 1831 Type locality: Ural Mountains, Kazakhstan

Holotype: ?

Podarcis irritans Menetries, 1832 Type locality: Saliane, near Baku, Azerbaïdjan Holotype: MNHNP 0.546

Lacerta boristhenis Dvigubsky, 1832 Type locality: ? Holotype: ?

Eremias arguta potanini Bedriaga, 1912 Type locality: Salburdi, Sakha Republic, Russia

Holotype: ZISP 5165

Distribution: Kazakhstan, Mongolia, northwestern China

Eremias arguta darevskii Tsaruk, 1986

Eremias arguta darevskii Tsaruk, 1986: ?

Type locality: Issyk Kul Lake, Kazakhstan

Holotype: ?

Distribution: Kyrgyzstan: basin of the Issyk Kul

Eremias arguta deserti (Gmelin, 1789)

Lacerta deserti Gmelin, 1789: 1076

Type locality: Ural Mountains, Russia

Holotype: ZMB, not located

Distribution: southeast Romania: Black Sea coast; southwestern Russia: Dagestan; northern Azerbaijan; Kazakhstan

Eremias arguta transcaucasica Darevsky, 1953

Eremias arguta transcaucasica Darevsky, 1953: 118

Type locality: Metz Mazra vilage, Basargecharsky area, Armenia

Holotype: ZMMU 2811

Distribution: Armenia, Azerbaijan, Georgia, Turkey: area of Lake Van

Eremias arguta uzbekistanica Tschernow, 1934

[Eremias arguta uzbekistanica Tschernow, 1934: ?](#)

Type locality: ?

Holotype: ?

Distribution: western Uzbekistan; southern Kazakhstan; Kyrgyzstan: Osch region, Talas Valley, Chuy Valley

Eremias aria Anderson & Leviton, 1967

[Eremias aria Anderson & Leviton, 1967: 1](#)

Type locality: 5-10 mi ENE Nimla on old Kabul-Jalalabad rd, and ca. 10 SW of Balabagh (30°19'21"N, 70°10'15"E), Afghanistan

Holotype: CAS 96204

Distribution: eastern Afghanistan: Kabul River valley

Eremias brenchleyi Günther, 1872

[Eremias brenchleyi Günther, 1872: 419](#)

Type locality: unknown

Yarkand and Chotan BMNH 1946.8.7.63

Distribution: northeastern China; southeastern Russia

Eremias buechneri Bedriaga, 1906

[Eremias buechneri Bedriaga, 1906: 184](#)

Type locality: Yarkand to Chotan, Xinjiang Uyghur Autonomous Region, China

Holotype: ZISP 7081

Distribution: western China: southern edge of Tarim basin desert from Yarkant and Quarquan Rivers, Shinkiang District, and doubtfully Quinghai.

Eremias cholistanica Baig & Masroor, 2006

[Eremias cholistanica Baig & Masroor, 2006: 167](#)

Type locality: Baghdad-i-Jadeed Campus (29°22'43"N, 71°46'10"E), Islamia University Bahwalpur, Pujab province, Pakistan

Holotype: PMNH 1685

Distribution: Pakistan: Cholistan Desert

Eremias dzungarica Orlova, Poyarkov, Chirirova, Nazarov, 2017

[Eremias dzungarica Orlova, Poyarkov, Chirirova, Nazarov, 2017: 12](#)

Type locality: 7 km west from Uyench Sum (= Uench-somon) (46° 05' N; 91° 56' E), Khovd Aimaq, Khoovor, Mongolia

Holotype: ZMMU R-12845

Distribution: Mongolian Altai Mountains ; China: Xinjiang Uyghur Autonomous Region north from Tian Shan Mountains; Kazakhstan: East Kazakhstan Province

Eremias fasciata Blandford, 1874

[Eremias fasciata Blandford, 1874: 32](#)

Type locality: Kerman, Iran

Lectotype: ZMB 9329

Distribution: eastern Iran, southern Afghanistan, western Pakistan

Eremias grammica (Lichtenstein, 1823)

[Lacerta grammica Lichtenstein, 1823: 100](#)

Type locality: Karakum Desert, Turkmenistan

Lectotype: ZMB 1095

Synonym:

Scapteira zarudnyi Lantz, 1928 Type locality: Tsharakhs, Zir-Kuh, Korasan, Iran

Syn types: ZISP 9322, 9323

Distribution: Turkmenistan; northeastern Iran; Uzbekistan; southern Kazakhstan; western China:

Ili River valley, Sinkiang province; southwestern Tajikistan; northern Afghanistan

Eremias intermedia (Strauch, 1876)

[Podarces intermedia Strauch, 1876: 28](#)

Type locality: Karakum Desert, Turkmenistan

Lectotype: ZISP 3664

Synonyms:

Eremias transcaspica Nikolsky, 1905 Type locality: Baherden, Turkmenistan

Holotype: ZISP 14429

Eremias intermedia oxyrrhina Bedriaga, 1912 Type locality: ? Holotype: ?

Distribution: southern Kazakhstan, Tajikistan, Turkmenistan, Uzbekistan, northeastern Iran.

Eremias isfahanica Rastegar-Pouyani, Hosseinian, Rafiee, KamiI, Rajabizadeh & Wink, 2016

[Eremias isfahanica Rastegar-Pouyani, Hosseinian, Rafiee, KamiI, Rajabizadeh & Wink, 2016: ?](#)

Type locality: 54 km northwest of Isfahan city, near Hassanije village within the Ghomishloo National Park (32.84° N, 51.10° E), Iran

Holotype. SUHC 3012

Distribution: Iran: Isfahan Province

Eremias kavirensis Mozaffari & Parham, 2007

[Eremias kavirensis Mozaffari & Parham, 2007: 569](#)

Type locality: Maranjab sand dunes (34°17'51''N, 51°50'57''E), Isfahan Province, Iran

Holotype: CAS 238636

Distribution: Iran: Isfahan Province: Kavir Desert.

Eremias kokshaaliensis Eremchenko & Panfilov, 1999

[Eremias kokshaaliensis Eremchenko & Panfilov, 1999: ?](#)

Type locality: Settlement Sary-Jaz, Terekti Ravine, Kyrgyzstan,

Holotype: NAS KR R 000580

Distribution: eastern Kyrgyzstan: Sary-Jaz Basin; China: Xinjiang province: Kokshaal Range

Eremias kopetdaghica Szczerbak, 1972

[Eremias strauchi kopetdaghica Szczerbak, 1972: 83](#)

Type locality: vicinity of Ai-Dere-Tuzli-Tepe, Kara-Kalinskii, Turkmenistan

Holotype: ZIK 4

Distribution: northeastern Iran, southern Turkmenistan

Eremias lalezharica Moravec, 1994

[Eremias lalezharica Moravec, 1994: 61](#)

Type locality: Lalezhar (29°31'N, 56°51'E), N foot of Mt. Lalezhar, Kerman Province, Iran

Holotype: NMP6V 34555/3

Distribution: southeastern Iran: Mt. Lalezhar and Darb-e Behesht

Eremias lineolata (Nikolsky, 1897)

[Scapteira lineolata Nikolsky, 1897: 330](#)

Type locality: between Feizabad and Nusi, eastern Iran

Lectotype: ZISP 8801

Synonym:

Scapteira grumgrzimai Bedriaga, 1906: 240 Type locality: ? Holotype: ?

Distribution: Turkmenistan, Uzbekistan, northeastern Iran, northern Afghanistan, southwestern Tajikistan, southern Kazakhstan

Eremias montana Rastegar-Pouyani & Rastegar-Pouyani, 2001

[Eremias montana Rastegar-Pouyani & Rastegar-Pouyani, 2001: 108](#)

Type locality: upland regions of the Zagros mountains, 60 km NE of city of Kermanshah (47°5' E, 34°52'N), Kermanshah Province, western Iran

Holotype: not located

Synonym:

Eremias novo Mozaffari & Parham, 2007 Type locality: upland area surrounded by the Zagros Mountains, about 21 km southwest of the city of Hamedan, Hamedan Province, western Iran
Syntypes: RUZM 96T-104T

Distribution: Iran: Kermanshah, Kurdistan, Hamedan, Qazvin Provinces

Eremias multiocellata Günther, 1872

Eremias multiocellata multiocellata Günther, 1872

[Eremias multiocellata Günther, 1872: 419](#)

Type locality: Gobi Desert, Mongolia

Holotype: BMNH 1946.8.7.69

Synonyms:

Eremias yarkandensis saturata Blanford, 1875 Type locality: Kuenlun Valleys, S. of Yarkand, Xinjiang Uygur Autonomous Region, China Syntypes: BMNH 1946.8.7.60-62

Podarces planiceps Strauch, 1876 Type locality: Gansu Province, China Holotype: ?

Eremias multiocellata reticulata Bedriaga, 1912 Type locality: eastern Junggar Basin, Xinjiang Uygur Autonomous Region, China Holotype: ?

Distribution: southern Mongolia, northern China

Eremias multiocellata bannikowi Szczerbak, 1973

[Eremias multiocellata bannikowi Szczerbak, 1973: 86](#)

Type locality: ?

Holotype: ?

Distribution: western Mongolia; Russia: southern Tuvinia, Tyva Republic

Eremias multiocellata kozłowi Bedriaga, 1907

[Eremias multiocellata kozłowi Bedriaga, 1907: 183](#)

Type localities: Osis Ssachsheu, Donhuang Co., and Nanshan, Guansu Prov., China

Syntypes: ?

Distribution: central China

Eremias multiocellata tsaganbogdensis Munkhabayar & Borkin, 2010

[Eremias multiocellata tsaganbogdensis Munkhabayar & Borkin, 2010: 123](#)

Type locality: Tsagaan Bogd Uul Mountain, Bayanhongor Aymak, southern Mongolia

Holotype: ZISP.19956.1

Distribution: Mongolia: Bayanhongor Aymak: Tsagaan Bogd Uul Mountain.

Eremias nigrocellata Nikolsky, 1896

[Eremias nigrocellata Nikolsky, 1896: 371](#)

Type locality: Sistan, Iran

Lectotype: ZISP 8800

Distribution: northeastern Iran, southern Turkmenistan, southern Uzbekistan, southwestern Tajikistan, northern Afghanistan

Eremias nikolskii Nikolsky, 1905

[Eremias nikolskii Nikolsky, 1905: 478](#)

Type locality: Altay Mountain Range, Kyrgyzstan

Holotype: ZISP ?

Distribution: eastern Uzbekistan; Kyrgyzstan: Chatkal Range, Fergan Kyrka Range, Alai Mts; southeastern Kazakhstan; northern Tajikistan: Turkestan Range

Eremias papenfussi Mozaffari, Ahmadzadeh & Parham, 2011

[Eremias papenfussi Mozaffari, Ahmadzadeh & Parham, 2011: ?](#)

Type locality: Sooleghan Mountains (35°47' 44.9"N, 51°14'20.2"E), Tehran Province in the Alborz Mountain Range, Iran

Holotype: ZFMK 91701

Distribution: Iran: Alborz Mountain Range

Eremias persica Blanford, 1875

[Eremias persica Blanford, 1874: 31](#)

Type locality: near Isfahan, Iran

Lectotype: BMNH 1946.8.7.32

Synonym:

Eremias nigrolateralis Rastegar-Pouyani & Nilson, 1998 Type locality: 150 km NE of Shiraz (53°9'E, 30°52'N), Fars Prov., south-central Iran Holotype: NHMG 5147

Distribution: southern Azerbaijan, Iran, southern Turkmenistan, Afghanistan, southwestern Pakistan.

Eremias pleskei Nikolsky, 1905

[Eremias pleskei Nikolsky, 1905: 481](#)

Type locality: Nakhichevan, Azerbaijan

Lectotype: ZISP 6724

Distribution: northeastern Turkey, Armenia, Azerbaijan, northwestern Iran

Eremias przewalskii (Strauch, 1876)

Eremias przewalskii przewalskii (Strauch, 1876)

[Podarces przewalskii Strauch, 1876: 43](#)

Type locality: Alashan Desert, Inner Mongolia, China

Holotype: BMNH 1946.8.7.30

Synonyms:

Podarces kessleri Strauch, 1876 Type localities: Alashan Desert, Inner Mongolia and Gansu Prov., China Syntypes: ?

Podarces brachydactyla Strauch, 1876 Type locality: Alashan Desert, Inner Mongolia, China Holotype: ?

Distribution: northern China: Sinkiang, Qinghai, Nei Mongolia provinces; southwestern Mongolia

Eremias przewalskii tuvensis Szczerbak, 1970

Eremias przewalskii tuvensis Szczerbak, 1970: 35

Type locality: between Tes River and Lake Tere-Khol, Borig Del Sands, Erzin district, Tuva Republic, Russia

Holotype: ZIK ?

Distribution: western Mongolia; Russia: southern Tuva Republic

Eremias quadrifrons (Strauch, 1876)

Podarces quadrifrons Strauch, 1876: 34

Type locality: Alashan Desert, Inner Mongolia, China.

Holotype: ?

Distribution: China: Nei Mongol province: Alashan Desert

Eremias regeli Nikolsky, 1905

Eremias regeli Nikolsky, 1905: 479

Type locality: Shirabad, Uzbekistan

Holotype: ZISP 6115

Distribution: eastern Turkmenistan, southwestern Tajikistan, northern Afghanistan

Eremias scripta (Strauch, 1867)

Eremias scripta scripta (Strauch, 1867)

Podarces scripta Strauch, 1867: 424

Type locality: Aralo-Caspian desert, Turkmenistan

Lectotype: ZISP 3669

Synonym:

Scapteira bilkewitschi Nikolsky, 1905 Type locality: Turkmenistan Holotype: MNKNU 26919

Distribution: southern Kazakhstan, Uzbekistan, Turkmenistan, southern Afghanistan, southwestern Pakistan

Eremias scripta lasdini (Zarewskij, 1918)

Scapteira lasdini Zarewskij, 1918: 84

Type locality: Turmenistan

Holotype: ZISP 12107

Distribution: southeastern Uzbekistan, southwestern Tajikistan

Eremias scripta pherganensis Szczerbak & Nashetko, 1973

Eremias scripta pherganensis Szczerbak & Washetko, 1973: 85

Type locality: Fergana Valley, Uzbekistan

Holotype: ?

Distribution: Uzbekistan: Fergana Valley

Eremias strauchi Kessler, 1878

Eremias strauchi Kessler, 1878: 166

Type locality: Echmiadzin, Armenia

Lectotype: MLSU 166

Distribution: Turkey: eastern Anatolia; Armenia; Azerbaijan; northwestern Iran

Eremias stummeri Wettstein, 1940

Eremias velox stummeri Wettstein, 1940: 83

Type locality: region of Lake Issyk Kul, vicinity of the town Przhevalsk, Kyrgyzstan
Holotype: NHMW 15664
Distribution: Kyrgyzstan: Issyk-Kul area; Kazakhstan: Naryn steppe; northwestern China.

Eremias suphani Basoglu & Hellmich, 1968

Eremias velox suphani Basoglu & Hellmich, 1968: 4
Type locality: Lake Van, Turkey
Holotype: ZDEU 31/1957
Distribution: eastern Turkey, northwestern Iran

Eremias szczerbaki Eremchenko, Panfilov & Zarinenko, 1992

Eremias multiocellata szczerbaki Eremchenko, Panfilov & Zarinenko, 1992: ?
Type locality: Arpa River Valley, Kyrgyzstan
Holotype: NAS KR 002056
Distribution: Kyrgyzstan: Arpa Valley

Eremias velox (Pallas, 1771)

Eremias velox velox (Pallas, 1771)

Lacerta velox Pallas, 1771: 406

Type locality: Inder Mountains, lower Ural River, northwestern Kazakhstan

Neotype: ZISP 16233

Synonyms:

Lacerta cruenta Pallas, 1771: 406 Type locality: ? Holotype: ?

Lacerta coccinea Merrem, 1820 Type locality: ? Holotype: ?

Lacerta tigrina Kuhl, 1820 Type locality: ? Holotype: ?

Lacerta leucosticta Lichtenstein, 1823 Type locality: Kazakhstan Syntypes: ZMB 1105, 1107

Lacerta gracilis Eichwald, 1831 Type locality: ? Holotype: ?

Lacerta vittata Eversmann, 1834 Type locality: Russia Holotype: ?

Eremias coeruleo-ocellata Duméril & Bibron, 1839 Type locality: Bakou, Azerbaijan

Holotype: MNHNP 0.2769

Eremias velox borkini Eremchenko & Panfilov, 1999

Eremias velox borkini Eremchenko & Panfilov, 1999: ?

Type locality: Issyk Kul, Kyrgyzstan

Holotype: ?

Distribution: Kyrgyzstan: Tien Shan mountains, Issyk Kul area

Eremias velox caucasia Lantz, 1928

Eremias velox caucasia Lantz, 1928: ?

Type locality: Azerbaijan

Syntypes: FMNH 83977-78

Distribution: Russia: Daghestan: upper Volga & Kalmuk Steppe; Turkmenistan, Azerbaijan

Eremias velox roborowskii Bedriaga, 1906

Eremias velox roborowskii Bedriaga, 1906: 181

Type localities: Ssatschsheu (=Dunhuang Co.). Gansu Prov. and Lukchun, Xinjian Uygur
Autonomous Region, China

Syntypes: ?

Distribution: western China

Eremias vermiculata Blanford, 1875

[Eremias vermiculata Blanford, 1875: 194](#)

Type locality: southern Xinjian Uygur Autonomous Region, China

Syntypes: ZSI 5308, 5340

Synonym:

Poderces pylzowi Strauch, 1876 Type locality: Alashan Desert, Inner Mongolia, China

Holotype: ?

Distribution: eastern Kazakhstan: Zaysan Lake area; northwestern China: Xinjiang, Ningxia, Gansu, and Nei Mongol provinces; and southern Mongolia

Eremias yarkandensis Blanford, 1875

[Eremias yarkandensis Blanford, 1875: 194](#)

Type locality: town Yarkend, Kashgaria, Xinjiang-Uyghur Autonomous region, China

Syntypes: ZSI 6306, 6340; MCZ R-7167

Distribution: Western China, Kyrgystan

Gallotia Oudart, 1839

Oudart, P-L., P.B. Webb & S. Berthelot. 1839. *Histoire naturelle des Iles Canaries*.



Vol. 2 (2): pl. 1, Paris :Béthune, éditeur, 1836-1850 Type species: *Lacerta galloti* Oudart, 1839

Gallotia galloti eisentrauti A & S Troidl

Gallotia atlantica (Peters & Doria, 1882)

Gallotia atlantica atlantica (Peters & Doria, 1882)

[Lacerta atlantica Peters & Doria, 1882: 433](#)

Type localities: Arrecife, Isla de Lanzarote, and Santa Cruz, Isla de Tenenife, Canary Islands, Spain

Syntypes: ZMB 10289, MSNG 10116

Synonyms:

Gallotia atlantica delibesi Castroviejo, Mateo & Collado, 1985 Type locality: Aringa, Isla de Gran Canaria, Canary Islands Holotype: EBD 18153

Gallotia atlantica ibagnezi Castroviejo, Mateo & Collado, 1985 Type locality: Isla de Alegranza, Las Palmas de Gran Canaria Prov., Canary Islands Holotype: EBD 16887

Gallotia atlantica laurae Castroviejo, Mateo & Collado, 1985 Type locality: Cueva de los Verdes, northeastern Isla de Lanzarote, Canary Islands Holotype: EBD 17054

Distribution: Spain: Canary Islands: Lanzarote, Graciosa, Montaña Clara, Alegranza and Roque del Este Islands

Gallotia atlantica mahoratae Bischoff, 1985

[Gallotia atlantica mahoratae Bischoff, 1985: 48](#)

Type locality: Paraja, Isla de Fuerteventura, Canary Islands
Holotype: ZFMK 35034
Distribution: Spain: Canary Islands: Isla de Fuerteventura, Isla de Lobos

Gallotia auaritae Mateo, Garcia-Marquez, López-Jurado & Barahona, 2001
[Gallotia simonyi auaritae Mateo, Garcia-Marquez, López-Jurado & Barahona, 2001: 54](#)
Type locality: Roque de Mazo, Isla de la Palma, Canary Islands
Holotype: DBULPGC 50
Distribution: Spain: Canary Islands: Isla de la Palma (probably extinct)

Gallotia bravoana Hutterer, 1985
[Gallotia simonyi bravoana Hutterer, 1985: 379](#)
Type locality: Barranca de Chiguarime (28°01'10"N, 17°10'10"W), Isla de La Gomera, Canary Islands
Holotype: ZFMK 42392
Distribution: Spain: Canary Islands: Isla de La Gomera: Parque Rural de Valle Gran Rey

Gallotia caesaris (Lehrs, 1914)
Gallotia caesaris caesaris (Lehrs, 1914)
[Lacerta caesaris Lehrs, 1914: 681](#)
Type locality: Las Lapas, Isla de El Hierro, Canary Islands
Syntypes: BMNH 1969.446-455
Distribution: Spain: Canary Islands: Isla de El Hierro, Roque Grande de Salmor

Gallotia caesaris gomerae (Boettger & Müller, 1914)
[Lacerta galloti gomerae Boettger & Müller, 1914: 73](#)
Type locality: Isla de La Gomera, Canary Islands
Holotype: SMF 6041.2a
Distribution: Spain: Canary Islands: Isla de La Gomera

Gallotia galloti (Oudart, 1839)
Gallotia galloti galloti (Oudart, 1839)
[Lacerta galloti Oudart, 1839: 4](#)
Type locality: Güfmar, Isla de Tenerife, Canary Islands
Lectotype: MNHNP 0.2744
Distribution: Spain: Canary Islands: central & southern Isla de Tenerife

Gallotia galloti eisentrauti Bischoff, 1982
[Gallotia galloti eisentrauti Bischoff, 1982: 371](#)
Type locality: Bajamar, Anaga Peninsula, Isla de Tenerife, Canary Islands
Holotype: ZFMK 34800
Distribution: Spain: Canary Islands: northern Isla de Tenerife

Gallotia galloti insulanagae Martin, 1985
[Gallotia galloti insulanagae Martin, 1985: 524](#)
Type locality: Roque de Fuera de Anaga, Isla de Tenerife, Canary Islands
Holotype: DZLU 194
Distribution: Spain: Canary Islands: Isla de Tenerife: Roque Fuera de Anaga

Gallotia galloti palmae (Boettger & Müller, 1914)
[Lacerta galloti palmae Boettger & Müller, 1914: 71](#)

Type locality: Isla de La Palma, Canary Islands
Holotype: ZSM 241/1913
Distribution: Spain: Canary Islands: Isla de La Palma

Gallotia intermedia Barrbadillo, Lacoba, Pérez-Mellado, Sancho & López-Jurado, 1999

Gallotia intermedia Barrbadillo, Lacoba, Pérez-Mellado, Sancho & López-Jurado, 1999: 65

Type locality: Risco de La Jaqueta in the Acanilado de Los Gigantes (28°15'N, 16°50'W), Isla de Tenerife, Canary Islands

Holotype: DZLU 2450

Distribution: Spain: Canary Islands: western Isla de Tenerife: Acanilado de Los Gigantes

Gallotia simonyi (Steindachner, 1889)

Gallotia simonyi simonyi (Steindachner, 1889)

Lacerta simonyi Steindachner, 1889: 260

Type locality: Roques del Zalmor, near Isla de Hierro, Canary Islands

Syntypes: NHMW 16254-56

Distribution: Spain: Canary Islands: Isla de Hierro: Roques del Zalmor

Gallotia simonyi machadoi López-Jurado, 1989

Gallotia simonyi machadoi López-Jurado, 1989: 266

Type locality: La Fuga de Gorreta, Isla de Hierro, Canary Islands

Holotype: MISN ?

Distribution: Spain: Canary Islands: Isla de Hierro: Risco de Tibataje, in La Fuga de Gorreta

Gallotia stehlini (Schenkel, 1901)

Lacerta galloti stehlini Schenkel, 1901: 187

Type locality: Las Palmas, Isla de Gran Canaria, Canary Islands

Lectotype: NHMB 4021

Distribution: Spain: Canary Islands: Isla de Gran Canaria

Gastropholis Fischer, 1886

Fischer, J. G. 1886. Herpetologische Notizen. *Abhandlungen des*

Naturwissenschaftlichen Vereins in Hamburg 9: 1-19 Type species: *Gastropholis vittata* Fischer, 1886



Gastropholis prasina Tristan

Gastropholis echinata (Cope, 1862)

Lacerta echinata Cope, 1862: 189

Type locality: West Africa

Holotype: USNM 5995

Synonym:

Lacerta langi Schmidt, 1919 Type locality: Medje, Ituri Forest, DR Congo Holotype: AMNH R10525

Distribution: Liberia, Ivory Coast, Ghana, Cameroon, Equatorial-Guinea, Gabon, northeastern Democratic Republic of Congo

Gastropholis prasina Werner, 1904

Gastropholis prasina Werner, 1904: 462

Type locality: Tanga, Tanzania

Holotype: ZMB 18087

Synonym:

Bedriagaia moreaui Loveridge, 1936 Type locality: Amani, Usambara Mts. Tanzania

Holotype: MCZR-39799

Distribution: northeastern Tanzania, southeastern Kenya

Gastropholis tropidopholis (Boulenger, 1916)

Bedriagaia tropidopholis Boulenger 1916: 113

Type locality: Modje, Ituri, DR Congo

Holotype: RMCA.1646

Distribution: Democratic Republic of Congo

Gastropholis vittata Fischer, 1886

Gastropholis vittata Fischer, 1886: 51

Type locality: coastal northern Tanzania

Syntypes: BMNH 1946.8.4.91-92

Synonym:

Gastropholis lutzei Tornier, 1900 Type locality: Tanga, Tanzania Holotype: ZMB 1581

Distribution: southeastern Kenya; eastern Tanzania; northeastern Mozambique

Heliobolus Fitzinger, 1843 Fitzinger, L. 1843. *Systema Reptilium*, Fasciculus Primus, Amblyglossae.



Vienna: Braumüller & Seidel. 106 pp
Type species: *Lacerta lugubris* Smith, 1838

Heliobolus lugubris  Froot

Heliobolus lugubris (Smith, 1838)

Lacerta lugubris Smith, 1838: 93

Type locality: district immediately beyond the northern frontier of the Cape Colony, South Africa
Syntypes: BMNH 1946.8.6.37-45
Distribution: Angola, Mozambique, Zimbabwe, Botswana, South Africa, Namibia

Heliobolus neumanni (Tornier, 1905)

Eremias neumanni Tornier, 1905: 376

Type locality: Brussa Valley, north of Lake Stephanie, Ethiopia

Holotype: ZMB 18356 (lost)

Distribution: southern Ethiopia: north bank of Chew Bahir (Lake Stephanie), and west and northeast of Lake Chamo

Heliobolus nitidus (Günther, 1872)

Heliobolus nitidus nitidus (Günther, 1872)

Eremias nitida Günther, 1872: 381

Type locality: western Africa

Holotype: BMNH, not located

Distribution: Guinea, Mali, Cote d'Ivoire, Ghana, Togo, Benin, Niger, Nigeria

Heliobolus nitidus garambensis (Schmidt, 1919)

Eremias nitida garambensis Schmidt, 1919: 511

Type locality: Garamba, DR Congo

Holotype: AMNH R10686

Distribution: Central African Republic, Cameroon, DR Congo

Heliobolus nitidus quadrinasalis (Chabanaud, 1918)

Eremias quadrinasalis Chabanaud, 1918: 108

Type locality: Fort Arschambo, Chad

Holotype: MNHNP 1904.161

Distribution: Chad

Heliobolus spekii (Günther, 1872)

Heliobolus spekii spekii (Günther, 1872)

Eremias spekii Günther, 1872: 381

Type locality: 5° 7' S, between the coast and Unjamwesi, Tanzania

Syntypes: BMNH 1946.8.6.8-9

Synonym:

Eremias rugiceps Peters, 1878 Type locality: Taita, Kenya Syntypes: ZMB 9287

Distribution: southeastern Kenya; Tanzania

Heliobolus spekii scortecii (Ariillo, Balletto & Spanò, 1965)

Eremias spekii scortecii Ariillo, Balletto & Spano, 1965: ?

Type locality: Migiurtinia, Somalia

Holotype: MSNG ?

Distribution: eastern Somalia

Heliobolus spekii sextaeniata (Stejneger, 1894)

Eremias sextaeniata Stejneger, 1894: 718

Type locality: Tana River, Kenya.

Holotype: USNM 20080

Distribution: South Sudan; southern Ethiopia; western Somalia; northern Uganda; northern Kenya

Hellenolacerta Arnold, Arribas & Carranza, 2007 Arnold, E.N., O. Arribas & S. Carranza. 2007.



Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* 1430: 1-86
Type species: *Lacerta graeca* Bedriaga, 1886

Hellenolacerta graeca  Benny Trapp

Hellenolacerta graeca (Bedriaga, 1886)

[Lacerta graeca Bedriaga, 1886: 290](#)

Type locality: Taygetos Mountains, near Kalamata, Greece

Holotype: BMNH 1940.9.3.46

Distribution: Greece: Peloponnese

Holaspis Gray, 1863 Gray, J.E. 1864. Description of two new genera of lizards (*Holaspis* and



Poriodogaster, A. Smith, MS). *Annals and Magazine of Natural History* (3) 13: 101-104. Type species: *Holaspis guentheri* Gray, 1863

Holaspis guentheri  J-F Trape

Holaspis guentheri Gray, 1863

[Holaspis guentheri Gray, 1863: 152](#)

Type locality: unknown

Holotype: BMNH 1946.8.7.31

Distribution: Sierra Leone, Guinea, Liberia, Cote d'Ivoire, Ghana, Togo, Benin, Nigeria, Cameroon, Central African Republic, DR Congo, Uganda, Equatorial Guinea, Gabon and Angola

Holaspis laevis Werner, 1896

Holaspis guentheri laevis Werner, 1896: 191

Type locality: Usambara Mts., Tanzania

Holotype: ?

Distribution: Kenya, Tanzania, Malawi, Mozambique, eastern DR Congo

Iberolacerta Arribas, 1997

Arribas, O. J. 1997. Morfologia, filogenia y bibliografía de las lagartijas de alta montana de los Pirineos [Microforma].

Tesis doctoral - Universitat Autònoma de Barcelona. *Publicaciones de la Universitat Autònoma de Barcelona*, 08193 Bellaterra (Barcelona), 353 pp. Type species: *Lacerta muralis monticola* Boulenger, 1905



Iberolacerta galani unknown

Iberolacerta aranica (Arribas, 1993)

Lacerta bonnali aranica Arribas, 1993: 135

Type locality: Coll de Barradós, Viella, Vall d'Aràn. Lleida, Spain

Holotype: OA 92081007

Distribution: France and Spain: Central Pyrenees: Maubèrme Massif

Iberolacerta aurelioi (Arribas, 1994)

Lacerta aurelioi Arribas, 1994: 334

Type locality: Estany de Sotillo, Macizo de la Pica d'Estats, Lérida, Spain

Holotype: CA 91070110

Distribution: France, Spain and Andorra: eastern Pyrenees: massifs of Montroig, Pica d'Estats, Coma Pedrosa and Sorteny

Iberolacerta bonnali (Lantz, 1927)

Lacerta monticola bonnali Lantz, 1927: 58

Type locality: Lac Bleu de Bigorre, Pyrenees, France

Holotype: BMNH 1947.1.1.19

Distribution: France and Spain: central Pyrenees: from El Portalé to La Bonaigua.

Iberolacerta cyreni (Müller & Hellmich, 1937)

Iberolacerta cyreni cyreni (Müller & Hellmich, 1937)

Lacerta monticola cyreni Müller & Hellmich, 1937: 67

Type locality: between Puerto de Cotos and Puerto de Navacerrada, La Granja de San Idefonso, Segovia, Spain.

Neotype: MNCN 39934

Distribution: Spain: Sierra del Guadarrama

Iberolacerta cyreni castilliana (Arribas, 1996)

Lacerta cyreni castilliana Arribas, 1996: 50

Type locality: Laguna Grande, Sierra de Gredos, central Spain

Holotype: CA 86071819

Distribution: Spain: Sierra de Bejar, Sierra de Gredos, Sierra La Serrota

Iberolacerta galani Arribas, Carranza & Odierna, 2006

Iberolacerta galani Arribas, Carranza & Odierna, 2006: 31

Type locality: Laguna de los Peces, Zamora, Sa Segundera, Spain

Holotype: IPE 4000

Distribution: Spain:

Iberolacerta horvathi (Méhely, 1904)

Lacerta horvathi Méhely, 1904: 363

Type locality: Jasenak, Croatia

Syntypes: NHMW 16025:1,2; BMNH 1946.9.1.24

Distribution: Croatia, Slovenia, northeastern Italy, eastern Austria

Iberolacerta martinezricai (Arribas, 1996)

Lacerta cyreni martinezricai Arribas, 1996: 54

Type locality: Peña de Francia, Salamanca, Spain

Holotype: IPE 3001

Distribution: Spain : Sierra de la Pena de la Francia

Iberolacerta monticola (Boulenger, 1905)

Iberolacerta monticola monticola (Boulenger, 1905)

Lacerta monticola Boulenger, 1905: 365

Type locality: Serra da Estrela, Portugal

Holotype: BMNH 1885.12.30.15

Distribution: Portugal: Serra da Estrela

Iberolacerta monticola astur Arribas, Galán, Remón & Naveira, 2014

Iberolacerta monticola astur Arribas, Galán, Remón & Naveira, 2014: 219

Type locality: Salientes, Palacios del Sil, León, Spain

Holotype: MNCN 44652

Distribution: Spain: Sierra de Gistreo

Iberolacerta monticola cantabrica (Mertens, 1929)

Lacerta monticola cantabrica Mertens, 1929: 284

Type locality: Rodiezmo, border of Leon and Asturias Provinces, northwestern Spain

Holotype: SMF 22098

Distribution: Spain: mountains of Cantabria and Galicia

Ichnotropis Peters, 1854 Peters, W.C.H. 1854. Diagnosen neuer Batrachier, welche zusammen mit der früher



(24. Juli und 17. August) gegebenen Übersicht der Schlangen und Eidechsen mitgeteilt werden. *Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlichen Preussische Akademie der Wissenschaften zu Berlin*. 1854: 614-628 Type species: *Ichnotropis macrolepidota* Peters, 1854

Ichnotropis capensis Johan Marais

Ichnotropis bivittata Bocage, 1866

Ichnotropis bivittata bivittata Bocage, 1866

[Ichnotropis bivittata Bocage, 1866: 43](#)

Type locality: Duque de Bragança, Angola

Syntype: ZMB 5827

Distribution: Angola; Gabon; DR Congo: southern Katanga; northern Zambia; northern Malawi; Tanzania

Ichnotropis bivittata pallida Laurent, 1964

[Ichnotropis bivittata pallida Laurent, 1964: 64](#)

Type locality: Boca da Humpata, Huila, Angola

Holotype: MBL 1854

Distribution: southwestern Angola: Serra da Bandeira

Ichnotropis capensis (Smith, 1838)

Ichnotropis capensis capensis (A. Smith, 1838)

[Algyra capensis Smith, 1838: 94](#)

Type locality: sandy deserts around Latakoo (Kuruman), Northern Cape province, South Africa

Holotype: lost

Synonyms:

Tropidosaura dumerelii Smith, 1849 Type locality: South Africa Holotype: not located

Ichnotropis microlepidota Peters, 1854 Type locality: Maputo, Mozambique Syntype: ZMB 6123

Ichnotropis capensis overlaeti Witte & Laurent, 1942 Type locality: Kapanga, Lualaba Province, DR Congo Holotype: RMCA 9691

Ichnotropis longipes Boulenger, 1902 Type locality: Mazoe, Mashonaland, Zimbabwe Syntypes: BMNH 1946.8.4.21-23

Distribution: Angola, Botswana, Mozambique, Namibia, South Africa, eastern DR Congo, Zambia

Ichnotropis capensis nigrescens Laurent, 1952

[Ichnotropis capensis nigrescens Laurent, 1952: 201](#)

Type locality: Bolobo, Mai-Ndombe Province, western DR Congo
Holotype: RMCA 14671
Distribution: central DR Congo

Ichnotropis chapini Schmidt, 1919

Ichnotropis chapini Schmidt, 1919: 508

Type locality: Aba, Uelle Region, DR Congo

Holotype: AMNH R10674

Holotype: FMNH 74285 northern DR Congo Last seen 1925

Ichnotropis grandiceps Broadley, 1967

Ichnotropis grandiceps Broadley, 1967: 1

Type locality: 25 miles west of Mohembo, Botswana, on the border of the Caprivi Strip

Holotype: USNM 163989

Distribution: Botswana, Namibia

Ichnotropis microlepidota Marx, 1956

Ichnotropis microlepidota Marx, 1956: 5

Type locality: foot of Mount Moco, Benguela Province, Angola

Holotype: FMNH 74285

Distribution: Angola / Benguela Province Last seen 1954

Ichnotropis tanganicana Boulenger, 1917

Ichnotropis tanganicana Boulenger, 1917: 278

Type locality: eastern shore of Lake Tanganyika, Tanzania.

Holotype: BMNH 1946.9.3.49

Distribution: western Tanzania Last seen 1896

Iranolacerta Arnold, Arribas & Carranza, 2007 Arnold, E.N., O. Arribas & S. Carranza. 2007.



Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. Zootaxa 1430: 1-86
Type species: *Lacerta brandtii* De Filippi, 1863

Iranolacerta brandtii brandtii Benny Trapp

Iranolacerta brandtii (De Filippi, 1863)

Iranolacerta brandtii brandtii (De Filippi, 1863)

Lacerta brandtii De Filippi, 1863: 387

northwestern Iran

Holotype: MRSN R2702

Distribution: northwestern Iran, southern Azerbaijan

Iranolacerta brandtii esfahanica (Nilson, Rastegar-Pouyani, Rastegar-Pouyani and Andr en. 2003)

[Lacerta brandtii esfahanica Nilson, Rastegar-Pouyani, Rastegar-Pouyani and Andr en. 2003: 15](#)

Type locality: main Zagros mountain Range, at Fereydun [= Feredun] Shahr, W of Daran (32°56'N, 50°05'E), Iran

Holotype: NHMG 5609

Distribution: Iran: southern and central Zagros Mountains

Iranolacerta zagrosica (Rastegar-Pouyani & Nilson, 1998)

[Lacerta zagrosica Rastegar-Pouyani & Nilson, 1998: 268](#)

Type locality: 3 km NW Fereyidin Shahr (32°58'N, 50°40'E), about 140 km NW Esfahan, Zagros Mountains, Esfahan Province, Iran

Holotype: NHMG 5149

Lacerta Linnaeus, 1758 Linnaeus, C. 1758. *Systema natur  per regna tria natur , secundum classes,*



ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio decima, reformata. Laurentii Salvii, Holmi . 10th Edition: 824 pp Type species: *Lacerta agilis* Linnaeus, 1758

Lacerta schreiberi Ronald Peck

Lacerta agilis Linnaeus, 1758

Lacerta agilis agilis Linnaeus, 1758

[Lacerta agilis Linnaeus, 1758: 203](#)

Type locality: southern Sweden

Syntypes: NRM 132-134

Synonyms:

Lacerta stirpium Dum ril & Bibron, 1839 Type locality: vicinity of Paris, France Syntypes: MNHNP0.2678-79

Lacerta fragilis Pamstruch & Swartz, 1808 Type locality: ? Holotype: Iconotype: pl. 27

Distribution: western Austria, Belgium, Denmark, Finland, France, western Germany, Liechtenstein, Luxembourg, Netherlands, Sweden, Switzerland, United Kingdom

Lacerta agilis argus (Laurenti, 1768)

[Seps argus Laurenti, 1768: 61](#)

Type locality: unknown

Holotype: ?

Distribution: eastern Austria, Croatia, Czech Republic, eastern Germany, Hungary, northeastern Italy, Poland, Romania, Serbia, Slovakia, Slovenia, Ukraine

Lacerta agilis boemica Suchow, 1929

[Lacerta boemica](#) Suchow, 1929: 117

Type locality: Vladikavkaz, Republic of North Ossetia–Alania, Russia

Holotype: ?

Distribution: Russia: northeastern Caucasus.

Lacerta agilis bosnica Schreiber, 1912

[Lacerta agilis bosnica](#) Schreiber, 1912: 483

Type locality: Bosnia

Holotype: ?

Distribution: Albania, Bosnia and Herzegovina, Bulgaria, Croatia, northern Greece, Kosovo, Macedonia, Montenegro

Lacerta agilis brevicaudata Peters, 1958

[Lacerta agilis brevicaudata](#) Peters, 1958: 127

Type locality: Stepanavan, Armenia

Holotype: ZISP 12900

Distribution: Armenia, Georgia, northeastern Turkey

Lacerta agilis chersonensis Andrzejowski, 1832

[Lacerta chersonensis](#) Andrzejowski, 1832: 327

Type locality: Carpathian Mountains, Romania

Holotype: ?

Synonym:

Lacerta agilis euxinica Fuhn & Vancea, 1964 Type locality: Danube delta, Black Sea coast, Romania Holotype: SMF 59716

Distribution: Belarus, Bulgaria, Estonia, Latvia, Lithuania, Moldova, Poland, Romania, western Russia, western Ukraine

Lacerta agilis exigua Eichwald, 1831

[Lacerta exigua](#) Eichwald, 1831: 188

Type locality: ?

Holotype: ?

Synonyms:

Lacerta agilis colchica Eichwald, 1841 Type locality: ? Holotype: ?

Lacerta agilis dilepis Lichtenstein, 1856 Type locality: Kazakhstan Syntype: ZMB 49520

Synonym:

Lacerta paradoxa Bedriaga, 1886 Type locality: Sochumi, Abkhazia, Russia Holotype: ?

Distribution: Azerbaijan, northwestern China, Kazakhstan, Kyrgyzstan, Mongolia, Russia, Ukraine

Lacerta agilis garzoni Palacios & Castroviejo, 1975

[Lacerta agilis garzoni](#) Palacios & Castroviejo, 1975: 6

Type locality: western outskirts of Puig de Basa, Tossas, Pirineos, provincia de Gerona, Spain.

Holotype: MNCN 44271

Distribution: Andorra, northern Spain

Lacerta agilis grusinica Peters, 1960

[Lacerta agilis grusinica](#) Peters, 1960: 280

Type locality: Suchum-Kale, Abkhazia

Syntypes: BMNH 1946.8.6.53-54
Distribution: western Georgia, northeastern Turkey

Lacerta agilis ioriensis Peters & Muskhelischwili, 1968
[Lacerta agilis ioriensis Peters & Muskhelischwili, 1968: 213](#)
Type locality: Tianeti, Georgia
Holotype: ZIT ?
Distribution: Georgia: upper Iori River in the Caucasus

Lacerta agilis mzymtensis Tuniyev & Tuniyev, 2008
[Lacerta agilis mzymtensis Tuniyev & Tuniyev, 2008: 64](#)
Type locality: mountain Aishkha-2, Adler Rayon of Sochi City, Krasnodar Kray, Russia
Holotype: ZISP 24648
Distribution: Russia: upper Mzymta River

Lacerta agilis tauridica Suchov, 1926
[Lacerta agilis tauridica Suchov, 1926: 83](#)
Type locality: Simferopol, Chumakarka settl., Ukraine, Crimea
Neotype: ZISP 12620
Distribution: Ukraine: southern Crimea

Lacerta bilineata Daudin, 1802

Lacerta bilineata bilineata Daudin, 1802
[Lacerta bilineata Daudin, 1802: 152](#)
Type locality: vicinity of Paris, France
Holotype: MNHNP, not located
Distribution: Andorra, Austria, Croatia, France, western Germany, northern Italy, Monaco, Slovenia, northern Spain, southern Switzerland, United Kingdom: Jersey, Guernsey

Lacerta bilineata chloronota Rafinesque-Schmaltz, 1810
[Lacerta chloronota Rafinesque-Schmaltz, 1810: 7](#)
Type locality: Sicily, Italy
Holotype: ?
Distribution: Italy: Calabria and Sicily

Lacerta bilineata chlorosecunda Taddei, 1950
[Lacerta bilineata chlorosecunda Taddei, 1950: 207](#)
Type locality: Puglia: Taranto, Soleto (Lecce) and Capo S. Maria di Leuca, Italy
Syntypes: MZUF 17316-23
Distribution: Italy: Apulia District

Lacerta bilineata fejervaryi Vasvary, 1926
[Lacerta viridis fejervaryi Vasvary, 1926: 37](#)
Type locality: Naples, Italy
Holotype: HNHM 2812
Distribution: Central Italy.

Lacerta media Lantz & Cyrén, 1920

Lacerta media media Lantz & Cyrén, 1920
[Lacerta viridis media Lantz & Cyrén, 1920: 35](#)
Type locality: Tiflis, Georgia

Lectotype: ZISP 12387

Distribution: Armenia, Azerbaijan, Georgia, northwestern Iran, southern Russia, eastern Turkey

Lacerta media ciliciensis Schmidtler, 1975

Lacerta trilineata media Schmidtler, 1975: 62

Type locality: 15 km N Kozan, Adana Province, Turkey

Holotype: ZSM 89/1972/1

Distribution: Turkey: Mersin, Adana, Kayseri provinces

Lacerta media isaurica Schmidtler, 1975

Lacerta trilineata isaurica Schmidtler, 1975: 59

Type locality: 6 m E Hadım, Konya Prov., Turkey

Holotype: NHMW 20328:3

Distribution: Turkey: central highlands of the Konya province

Lacerta media israelica Peters, 1964

Lacerta trilineata israelica Peters, 1964: 244

Type locality: Jerusalem

Holotype: TAU R990

Distribution: northern Israel, northern Jordan, southern Lebanon

Lacerta media wolterstorffi Mertens, 1922

Lacerta strigata wolterstorffi Mertens, 1922: 193

Type locality: Beirut, Lebanon

Holotype: SMF 13244

Distribution: southeastern Turkey: Hatay province; northwestern Syria; Lebanon

Lacerta pamphylica Schmidtler, 1975

Lacerta trilineata pamphylica Schmidtler, 1975: 57

Type locality: 15-25 km N. Alanya, Antalya Prov., Turkey

Holotype: ZSM 5/1971

Distribution: Turkey: central-southern coast

Lacerta schreiberi Bedriaga, 1878

Lacerta schreiberi Bedriaga, 1878: 299

Type locality: Spain

Syntypes: BMNH 1946.9.2.60-63

Distribution: North and Central Spain and Portugal

Lacerta strigata Eichwald, 1831

Lacerta strigata Eichwald, 1831: 189

Type locality: Kislovodsk, southern Russia

Holotype: unknown

Synonyms:

Lacerta viridis woosnami Boulenger, 1917 Type locality: south coast of the Caspian Sea, Iran
Syntypes: BMNH 1946.8.5.27-33

Lacerta quinquevittata Menetries, 1832 Type locality: Caucasus, Russia Syntypes: MNHNP
0.2692-93

Distribution: eastern Georgia, Armenia, Azerbaijan, northeastern Turkey, Caspian coast of northern Iran, southwestern Turkmenistan

Lacerta trilineata Bedriaga, 1886

Lacerta trilineata trilineata Bedriaga, 1886

[Lacerta viridis trilineata](#) Bedriaga, 1886: 99

Type locality: Greece

Holotype: BMNH 1946.9.2.59

Distribution: Macedonia; Greece: central and eastern mainland, Peloponnese, southern Ionian Islands; western Bulgaria

Lacerta trilineata cariensis Peters, 1964

[Lacerta trilineata cariensis](#) Peters, 1964: 241

Type locality: Bornova, İzmir Province, Turkey

Holotype: NHMG 4661

Distribution: Turkey: West Anatolia; offshore islands of the Mediterranean

Lacerta trilineata citrovittata Werner, 1935

[Lacerta viridis citrovittata](#) Werner, 1935: 155

Type locality: Tinos Island, the Cyclades, Greece

Holotype: ?

Distribution: Greece: Tinos, Syros, Andros, Naxos and Ios Islands

Lacerta trilineata diplochondrodes Wettstein, 1952

[Lacerta strigata diplochondrodes](#) Wettstein, 1952: 255

Type locality: Monolito, Rhodes Island, Greece

Syntypes: NHMW 14903:1,2

Distribution: Turkey: eastern Aegean coast; Greece: Rhodes Island

Lacerta trilineata dobrogica Fuhn & Mertens, 1959

[Lacerta trilineata dobrogica](#) Fuhn & Mertens, 1959: 34

Type locality: Malul Lake, Bugeae to Almalau, Adamklissi District, Dobruscha, Romania

Holotype: SMF 55246

Distribution: Black Sea coast of Bulgaria, Romania, European Turkey; notheastern Greece

Lacerta trilineata galatiensis Peters, 1964

[Lacerta trilineata galatiensis](#) Peters, 1964: 242

Type locality: Serai Koi, at the railway line Old Schehir after Ankara, Turkey

Holotype: NHMW 10900:9

Distribution: Turkey: middle northern Anatolia

Lacerta trilineata hansschweizeri Müller, 1935

[Lacerta major hansschweizeri](#) Müller, 1935: 236

Type locality: Milos Island, the Cyclades, Greece

Syntypes: ZSM 2010

Distribution: Greece: Cylades: Milos, Kimolos, Sifnos, Serifos Islands.

Lacerta trilineata major Boulenger, 1887

[Lacerta viridis major](#) Boulenger, 1887: 16

Type locality: Dalmatia, Croatia

Holotype: BMNH 1946.9.2.44

Distribution: Croatia, Montenegro, Bosnia-Herzegovina, Albania; western Greece, including Corfu and Paxos Islands

Lacerta trilineata polylepidota Wettstein, 1952

Lacerta strigata polylepidota Wettstein, 1952: 255

Type locality: Chania, northwestern Crete, Greece

Syntypes: NHMW 16261:1, 2; 16262: 1-11; 1490:1-25

Distribution: Greece: Crete

Lacerta viridis (Laurenti, 1768)

Lacerta viridis viridis (Laurenti, 1768)

Seps viridis Laurenti, 1768: 62

Type locality: Vienna, Austria

Holotype: no located

Synonyms:

Lacerta viridis brandenburgiensis Hecht, 1930 Type locality: Mark Brandenburg, Germany

Holotype: ZMB 32718

Lacerta viridis magnifica Sobolevsky, 1930 Type locality: Yalta, Crimea, Ukraine Syntypes:

ZMMU 2470-73

Lacerta viridis istriensis Werner, 1897 Type locality: Istria Peninsula Syntypes: NHMW 10835

Distribution: Albania, Austria, Bosnia and Herzegovina, Bulgaria, Croatia, Czech Republic, Germany, Greece, Hungary, Kosovo, Macedonia, Moldova, Montenegro, Romania, Serbia, Slovakia, Slovenia, Ukraine

Lacerta viridis guentherpetersi Rykena, Nettmann & Mayer, 2001

Lacerta viridis guentherpetersi Rykena, Nettmann & Mayer, 2001: 90

Type locality: Euboea, Dirfis Mountains, at the pass above Steni, at the old tavern, Greece

Holotype: NHMW 31693.2

Distribution: Greece: Island of Evia, Dirfis Mountains

Lacerta viridis infrapunctata Schmidtler, 1986

Lacerta viridis infrapunctata Schmidtler, 1986: 42

Type locality: coast E Esbiye, Prov. Giresun, Turkey

Holotype: NHMW, not located

Distribution: Turkey: eastern Black Sea coast

Lacerta viridis meridionalis Cyrén, 1933

Lacerta viridis meridionalis Cyrén, 1933: 231

Type locality: Büyükdere, Istanbul Province, Turkey

Syntypes: ? 4568-72

Distribution: eastern Romania, eastern Bulgaria, northwestern Turkey, northeastern Greece

Lacerta viridis paphlagonica Schmidtler, 1986

Lacerta viridis paphlagonica Schmidtler, 1986: 40

Type locality: plane forest at Kabala, southwest Sinop, Prov. Sinop, Turkey

Holotype: ZSM 193/1985

Distribution: Turkey : central Black Sea coast.

Latastia Bedriaga, 1884

Bedriaga, J. von. 1884. Die neue Lacertiden-Gattung *Latastia* und ihre Arten.



Annali del Museo Civico di Storia Naturale Giacomo Doria 20: 307-324

Type species: *Lacerta samharica*
Blanford, 1870

Latastia longicaudata longicaudata

 Tomas Mazuch

Latastia boscai Bedriaga, 1884

Latastia boscai boscai Bedriaga, 1884

[Latastia boscai Bedriaga, 1884: 16](#)

Type locality: from the land of the Adal (Rugdeira Sogheira) and from the land of the Bogos (Keren), Somalia

Syntype: MSNG 10118

Synonym:

Latastia wachei Werner, 1913 Type locality: 70 km NW Harrar, Ethiopia Syntypes: ZMH R01224-25

Distribution: Ethiopia, northwestern Somalia, Eritrea, Djibouti.

Latastia boscai arenicola Parker, 1942

[Latastia boscai arenicola Parker, 1942: 71](#)

Type locality: Dolo, Somalia

Syntypes: BMNH 1946.9.4.94

Distribution: central Somalia

Latastia boscai burii Boulenger, 1907

[Latastia burii Boulenger, 1907: 393](#)

Type locality: inland of Berbera, Somalia

Syntypes: BMNH 1946.9.2.94-95

Distribution: northeastern Somalia

Latastia caeruleopunctata Parker, 1935

[Latastia caeruleopunctata Parker, 1935: 527](#)

Type locality: Haud region (08°37'N, 45°09'E), Ethiopia-Somali border

Holotype: BMNH 1946.9.2.77

Synonym:

Latastia ventralis Werner, 1917 Type locality: Somalia Holotype: ZMH, not located

Distribution: north and central Somalia, eastern Ethiopia

Latastia cherchii Arillo, Balletto & Spano, 1967

[Latastia cherchii Arillo, Balletto & Spano, 1967: 132](#)

Type locality: Obbia, Mudug region of Somalia

Holotype: MZUF 768

Distribution: Somalia: Mudug region

Latastia doriai Bedriaga, 1884

Latastia doriai doriai Bedriaga, 1884

[Latastia doriai Bedriaga, 1884: 7](#)

Type locality: Rugdeira Sogheira (Adal), northern Somalia

Syntypes: BMNH 1946.9.2.85-87; MSNG 28250

Distribution: northern Somalia, northern Ethiopia

Latastia doriai martensi Bedriaga, 1884

[Latastia doriai martensi Bedriaga, 1884: 316](#)

Type locality: Keren, Bogos country, Eritrea

Syntypes: MSNG 28251-52, 36540; BMNH 1946.9.2.84

Distribution: Eritrea

Latastia doriai scortecii Arillo, Balletto & Spano, 1967

[Latastia doriai scortecii Arillo, Balletto & Spano, 1967: 140](#)

Type locality: Mandah, Somalia

Holotype: BMNH 1902.12.13.42

Distribution: northwestern Somalia, Djibouti, northern Ethiopia.

Latastia johnstonii Boulenger, 1907

[Latastia johnstonii Boulenger, 1907: 392](#)

Type locality: Masuku and Nyika Plateau, Malawi

Holotype: BMNH 1946.9.2.93

Synonyms:

Latastia kidwelli Boulenger, 1919 Type locality: Eldorado, Zimbabwe Holotype: MMK ?

Latastia bredoi De Witte, 1942 Type locality: Musosa, Tanganyika District, DR Congo

Holotype: IRSNB 1470

Distribution: western Tanzania, Malawi, eastern DR Congo, Zambia, Mozambique, Zimbabwe

Latastia longicaudata (Reuss, 1834)

Latastia longicaudata longicaudata (Reuss, 1834)

[Lacerta longicaudata Reuss, 1834: 29](#)

Type locality: Massawa, Eritrea

Lectotype: SMF 13601

Synonyms:

Lacerta samharica Blanford, 1870 Type locality: Margableh, near Assab, Eritrea Holotype: MSNG, not located

Lacerta sturti Blanford, 1870 Type locality: Komaylo Pass, near Undul Wells, Ethiopia

Holotype: MSNG, not located

Distribution: Cameroon, Djibouti, Eritrea, Ethiopia, Mali, Senegal, Somalia, Sudan, South Sudan, southeastern Egypt, Ethiopia, northern Somalia

Latastia longicaudata andersonii Boulenger, 1921

[Latastia longicaudata andersonii Boulenger, 1921: 30](#)

Type locality: Al-Khobar, Yemen

Syntypes: BMNH 1946.9.2.81-83

Distribution: Yemen

Latastia longicaudata lanzai Arillo, Balletto & Spano, 1967

[Latastia longicaudata lanzai Arillo, Balletto & Spano, 1967: 128](#)

Type locality: Dinsoor, southwestern Bay region, Somalia

Holotype: MZUF 1705

Distribution: southern Somalia, northern and southeastern Kenya, northeastern and central Tanzania.

Latastia longicaudata revoili (Vaillant, 1882)

[Eremias revoili](#) Vaillant, 1882: 20

Type locality: Darror Valley, Somalia

Holotype: MNHNP 0.5898

Distribution: southern Ethiopia, northeastern Somalia

Latastia ornata Monard, 1940

[Latastia ornata](#) Monard, 1940: 158

Type locality: Bafatá, central Guinea-Bissau

Holotype: MHNCF 608

Distribution: Guinea-Bissau Last seen 1938

Latastia petersiana Mertens, 1938

[Latastia petersiana](#) Mertens, 1938: 6

Type locality: Type locality: Brava, Somalia

Holotype: ZMB 8028

Distribution: southern Somalia.

Latastia siebenrocki (Tornier, 1905)

[Eremias siebenrocki](#) Tornier, 1905: 386

Type locality: unknown

Holotype: ZMB 17954

Distribution: Tanzania: near Tabora; Kenya: Eldama River Last seen 1905

Latastia taylori Parker, 1942

[Latastia taylori](#) Parker, 1942: 72

Type locality: Buran Valley, Somalia

Holotype: BMNH 1931.7.20.337

Distribution: northeastern Somalia

Meroles Gray, 1838 Gray, J.E. 1838. Catalogue of the Slender-tongued Saurians, with Descriptions of many



new Genera and Species. *Annals and Magazine of Natural History* 1: 274-283, 388-394. Type species: *Lacerta knoxii* Milne-Edwards, 1829

[Meroles anchietae](#)  Buckham

Meroles anchietae (Bocage, 1867)

Pachyrhynchus anchietae Bocage, 1867: 227

Type locality: Rio Croco, Mossamedes, Angola

Holotype: MBL, probably lost in fire

Distribution: Namibia: Namib Desert; southwestern Angola

Meroles ctenodactylus (Smith, 1838)

Lacerta ctenodactylus Smith, 1838: 93

Type locality: Great Namaqualand, Namibia

Holotype: lost

Synonym:

Lacerta capensis A. Smith, 1838 Type locality: sandy districts of Great Namaqualand,
Namibia Syntypes: BMNH, not located

Distribution: southwestern Namibia; South Africa: western North Cape Province

Meroles cuneirostris (Strauch, 1867)

Scapteira cuneirostris Strauch, 1867: 411

Type locality: Damaraland, northwestern Namibia

Holotype: not located

Distribution: Namibia: southern Namib Desert

Meroles knoxii (Milne-Edwards, 1829)

Meroles knoxii knoxii (Milne-Edwards, 1829)

Lacerta knoxii Milne-Edwards, 1829: 76

Type locality: Cape of Good Hope

Syntypes: BMNH 1946.8.7.70-73, 1946.8.7.41

Distribution: South Africa: Western Cape, Northern Cape Provinces

Meroles knoxii pequensis (Hewitt, 1935)

Scaptira knoxii pequensis Hewitt, 1935: 324

Type locality: Lüderitz Bay, southern coast of Namibia

Syntypes: PEM R 16102-04

Distribution: southwestern Namibia

Meroles micropholidotus (Mertens, 1938)

Scapteira micropholidotus Mertens, 1938: 12

Type locality: Lüderitz Bay, Namibia

Holotype: SMF 22245

Distribution: coastal Namibia: between Lüderitz Bay and Walvis Bay

Meroles reticulatus (Bocage, 1867)

Scapteira reticulata Bocage, 1867: 225

Type locality: Mossamedes, Angola

Holotype: MBL, probably lost in fire

Synonym:

Eremias serripes Peters, 1870 Type locality: Benguella, Angola Syntype: ZMB 6475

Distribution: coastal central Namibia to southwestern Angola

Meroles squamulosus (Peters, 1854)

Ichnotropis squamulosa Peters, 1854: 617

Type locality: Tete, Mozambique
Syntypes: ZMB 6614-16
Distribution: southern Angola, eastern Namibia, Botswana, Zambia, Malawi, southern Tanzania, Zimbabwe, Mozambique, Swaziland, northeastern South Africa

Meroles suborbitalis (Peters, 1869)

Eremias suborbitalis Peters, 1869: 658

Type locality: Damaraland,
Namibia Lectotype: ZMB 6150

Distribution: South Africa: western Western Cape, western North Cape Provinces;
southern Namibia

Mesalina Gray, 1838 Gray, J.E. 1838. Catalogue of the Slender-tongued Saurians, with Descriptions of many new Genera and Species. *Annals and Magazine of Natural History* 1: 274-283, 388-394. Type species: *Mesalina lichtensteini* Gray, 1838



Mesalina balfouri Sindaco

Mesalina adramitana (Boulenger, 1917)

Eremias adramitana Boulenger, 1917: 279

Type locality: Hadramut, Yemen

Syntypes: BMNH 1946.8.6.61-68

Distribution: Yemen, Oman, Qatar, United Arab Emirates, eastern Saudi Arabia

Mesalina arnoldi Sindaco, Simó-Riudalbas, Sacchi, Carranza, 2018

Mesalina arnoldi Sindaco, Simó-Riudalbas, Sacchi, Carranza, 2018: 530

Type locality: Yemen, Amran Governorate, plateau between Zakatin village (Hababah) to Kawkaban (Haraz Mt.) (about 15.51°N, 43.86°E)

Holotype: MCCI-R890

Distribution: highlands of southwestern Saudi Arabia and western Yemen.

Mesalina austroarabica Sindaco, Simó-Riudalbas, Sacchi, Carranza, 2018

Mesalina austroarabica Sindaco, Simó-Riudalbas, Sacchi, Carranza, 2018: 528

Type locality: Oman, Dhofar Governorate, Jebel Samhan at 17.1161°N, 54.7131°E

Holotype: MCCI-R1611

Distribution: southern Arabia: Oman: from the Jebel Samhan in Dhofar to the Yemen Mountains

Mesalina ayunensis Arnold, 1980

Mesalina ayunensis Arnold, 1980: 309

Type locality: Ayun, Dhofar, Oman

Holotype: BMNH 1977.1169

Distribution: southwestern Oman

Mesalina bahaeldini Segoli, Cohen & Werner, 2002

Mesalina bahaeldini bahaeldini Segoli, Cohen & Werner, 2002

Mesalina bahaeldini Segoli, Cohen & Werner, 2002: 169

Type locality: near St. Catherine's Monastery, Sinai, Egypt.

Holotype: HUI R10712
Distribution: Egypt: southern Sinai

Mesalina bahaeldini curatorum Werner & Ashkenazi, 2010

Mesalina bahaeldini curatorum Werner & Ashkenazi, 2010: 126

Type locality: Suez, Egypt
Holotype: ZMB 1122
Distribution: Egypt: northern Sinai

Mesalina balfouri (Blandford, 1881)

Eremias balfouri Blandford, 1881: 467 Type

locality: Socotra, Yemen Syntypes:

BMNH 1946.8.7.26-29

Distribution: Yemen: Socotra, Darsa, Samha, Abd al-Kuri Islands

Mesalina bernoullii (Schenkel, 1901)

Eremias bernoullii Schenkel, 1901: 187

Holotype. NMB 4396. Type locality: 'Palmyra' (Syria).

Distribution. Sinai, Jordan, Syria, Iraq, south-western Iran, Kuwait, north-eastern Saudi Arabia

Mesalina brevirostris Blandford, 1874

Mesalina brevirostris brevirostris Blandford, 1874

Mesalina brevirostris Blandford, 1874: 32

Type localities: Kalabagh, Punjab, Pakistan; Tunb Islands, Iran

Syntypes: BMNH 1946.8.6.34; ZSI 3474

Distribution: Iran, Iraq, Pakistan, Qatar, Saudi Arabia, Syria, United Arab Emirates

Mesalina brevirostris fieldi (Haas & Werner, 1969)

Eremias brevirostris fieldi Haas & Werner, 1969: 356 Type

locality: Mahor Birinji, southwestern Iran

Holotype: MCZ 56617

Distribution: southwestern Iran

Mesalina ercolinii (Lanza & Poggesi, 1975)

Eremias ercolinii Lanza & Poggesi, 1975: 305

Type locality: Bud Bud (about 4°12' N, 46° 28' E), 70 km S of El Bur, central Somalia.

Holotype: MZUF 5253

Distribution: central Somalia Last seen 1968

Mesalina guttulata (Lichtenstein, 1823)

Lacerta guttulata Lichtenstein, 1823: 101

Type locality: lower Egypt

Lectotype: ZMB 1119 Synonyms:

Scapteira punctulata Gray, 1838 Type locality: Egypt Syntypes: BMNH 1946.8.7.76-77

Eremias pardalis Duméril & Bibron, 1839 Type locality: Egypt Holotype: MNHNP 0.2786

Eremias pardaloides Boulenger, 1887 Type locality: Hcnjam (=Qeshm Island) Holotype: ZSI 3381

Distribution: Algeria, Egypt, Israel, Jordan, Libya, Morocco, Niger, Saudi Arabia, Senegal, Sudan, Syria, Tunisia, Yemen, Iran: Persian Gulf

Mesalina kuri Joger & Mayer, 2002

Mesalina kuri Joger & Mayer, 2002: 502

Type locality: Abd al-Kuri Island (12°11'N, 53°14'E), Yemen

Holotype: HLMD RA-2796-HT

Distribution: Yemen: Abd al-Kuri Island

Mesalina martini (Boulenger, 1897)

Eremias martini Boulenger, 1897: 468

Type locality: Obok, Djibouti

Syntypes: MNHNP 1893.139-40, 1893.86-87

Distribution: Djibouti, southeastern Egypt, Eritrea, northern Ethiopia, northern Somalia, eastern Sudan, southwestern Yemen

Mesalina microlepis (Angel, 1936)

Eremias brevisrostris microlepis Angel, 1936: 112

Type locality: Haouarine, SE Homs, Syria

Holotype: MNHNP 1935.285

Distribution. Lebanon, Syria, northern Jordan and southern Turkey

Mesalina olivieri (Audouin, 1829)

Mesalina olivieri olivieri (Audouin, 1829)

Lacerta olivieri Audouin, 1829: 122

Type locality: Egypt.

Syntypes: BMNH 1946.8.6.56-57

Synonyms:

Eremias guichenoti Doumergue, 1901 Type locality: Oran, Algeria Syntypes: not located

Eremias guttulata susana Boulenger, 1918 Type locality: Susa, Tunisia Holotype: BMNH 1946.8.7.74

Eremias guttulata latasti Boulenger, 1918 Type locality: Ausala, Algiers prov., Algeria

Holotype: BMNH 1886.4.13.9

Distribution: Algeria, Egypt, Libya, Mauritania, Morocco, Senegal, Tunisia, Western Sahara

Mesalina olivieri schmidtii (Haas, 1951)

Eremias olivieri schmidtii Haas, 1951: 276

Type locality: Wadi Nefk, Israel

Holotype: HNJ 1765

Distribution: Egypt: Sinai Peninsula; Palestine, Israel, Jordan.

Mesalina pasteuri (Bons, 1960)

Eremias pasteuri Bons, 1960: 69

Type locality: Amguit, Ahaggar, Algeria

Holotype: MNHN 1932.141

Distribution: southern Morocco, Western Sahara, Algeria: Sahara; Mali, Niger, northwestern Egypt

Mesalina rubropunctata (Lichtenstein, 1823)

Lacerta rubropunctata Lichtenstein, 1823: 100

Type localities: Egypt, northern Sudan

Syntypes: ZMB 1115-16, 1127 (Egypt); 1113-14 (Sudan)

Synonyms:

Scapteira nebulosa Gray, 1838 Type locality: North Africa Holotype: BMNH 1946.8.7.37

Mesalina lichtensteini Gray, 1838 Type locality: Egypt Syntypes: BMNH 1946.8.7.52-55

Distribution: Western Sahara, southern Morocco, southern Algeria, Libya, Egypt, northern Sudan, Mali, Niger

Mesalina saudiarabica Moravec, Smid, Schmitz, Shobrak, Wilms, 2017

Mesalina saudiarabica Moravec, Smid, Schmitz, Shobrak, Wilms, 2017: 10

Type locality: Mahazat as-Sayd (22.237°N, 41.843°E, Makkah Province, Saudi Arabia)

Holotype: ZFMK 91912

Distribution: central-western Saudi Arabia on the central plateau

Mesalina simoni (Boettger, 1881)

Podarces simoni Boettger, 1881: 571

Type locality: near Casablanca, Morocco
Holotype: SMF 6087
Distribution: Morocco: north of High Atlas Range

Mesalina watsonana (Stoliczka, 1872)

Eremias watsonana Stoliczka, 1872: 86

Type locality: between Karachi and Sukkur, Sind, Pakistan
Syntypes: BMNH 1946.8.7.75; ZSI 4929, 5050, 5223-5225; NHMW 23474:1-3
Distribution: Iran, Turkmenistan, Afghanistan, Pakistan, northwestern India: Thar Desert

Nucras Gray, 1838 Gray, J.E. 1838. Catalogue of the Slender-tongued Saurians, with Descriptions of many new Genera and Species. *Annals and Magazine of Natural History* 1: 274-283, 388-394. Type species: *Lacerta lalandii* Milne-Edwards, 1829



Nucras ornata Tyrone Ping

Nucras aurantiaca Bauer, Childers, Broeckhoven, Mouton, 2019
Nucras aurantiaca Bauer, Childers, Broeckhoven, Mouton, 2019: 154
Type locality: Republic of South Africa, Western Cape Province, Lamberts Bay, Bosduifklip Restaurant (32°05'04.36"S, 18°21'26.9 E).
Holotype: NMB R11626
Distribution: South Africa: Western Cape Province: Fynbos Biome

Nucras boulengeri Neumann, 1900

Nucras boulengeri boulengeri Neumann, 1900

Nucras boulengeri Neumann, 1900: 56

Type locality: Luba's village, Busoga, Uganda

Syntypes: ?

Synonym:

Nucras emini Boulenger, 1907 Type locality: southern shore of Lake Victoria, Tanzania

Holotype: BMNH 1946.8.6.16

Distribution: western Kenya, northern Tanzania, southern Uganda

Nucras boulengeri kilosae Loveridge, 1922

Nucras kilosae Loveridge, 1922: 314

Type locality: Tuidiga, south of Kilosa, Tanzania

Syntypes: BMNH 1946.8.6.58-59

Distribution: southern Tanzania, northern Zambia

Nucras caesicaudata Broadley, 1972

Nucras caesicaudata Broadley, 1972: ?

Type locality: Sazale Pan, Gona-re-Zhou, southeastern Zimbabwe

Holotype: NMZB 12380

Distribution: southern Mozambique, southeastern Zimbabwe, northeastern South Africa

Nucras holubi (Steindachner, 1882)

Eremias holubi Steindachner, 1882: 83

Type locality: Crocodile River, Mpumalanga Province, South Africa

Lectotype: NHMW 19056

Distribution: Namibia, Botswana, Zimbabwe, western Mozambique, Swaziland, eastern South Africa.

Nucras intertexta (Smith, 1838)

Lacerta intertexta Smith, 1838: 93

Type locality: country near Latakoo, Cape Colony, South Africa

Holotype: BMNH 1946.8.6.1.

Distribution: southern Mozambique; South Africa: northern Kwa-Zulu Natal; Angola, Botswana, northern Namibia, Zimbabwe

Nucras lalandii (Milne-Edwards, 1829)

Lacerta lalandii Milne-Edwards, 1829: 84

Type locality: Cape of Good Hope, South Africa

Syntypes: MNHNP 0.259, 0.264, 0.2745

Distribution: South Africa: southern Western Cape, Eastern Cape, KwaZulu-Natal Province; Swaziland.

Nucras livida (Smith, 1838)

Lacerta livida Smith, 1838: 92

Type locality: northern parts of Cape Colony, South Africa

Syntypes: BMNH 1946.8.6.81-83

Distribution: South Africa: Eastern Cape Province

Nucras ornata (Gray, 1864)

Teira ornata Gray, 1864: 380

Type locality: near the Zambezi River, probably Zimbabwe

Holotype: BMNH 1864.1.9.30

Distribution: South Africa: KwaZulu-Natal, Mpumalanga, Limpopo Provinces; Zimbabwe, Mozambique, southeastern Tanzania, Zambia, Malawi, Swaziland

Nucras scalaris Laurent, 1964

Nucras scalaris Laurent, 1964: 58

Type locality: Alto Chicapa, Lunda, Angola

Holotype: MD 5401

Distribution: northeastern Angola: Lunda Sul Province

Nucras taeniolata (Smith, 1838)

Lacerta taeniolata Smith, 1838: 93

Type locality: Grahamstown, Albany District, Eastern Cape Province, South Africa

Lectotype: BMNH 1946.8.4.25

Distribution: South Africa: southeastern Eastern Cape Province.

Nucras tessellata (Smith, 1838)

Lacerta tessellata Smith, 1838: 92

Type locality: South Africa

Holotype: BMNH, not located

Synonym:

Lacerta elegans Smith, 1838 Type locality: Little Namaqualand and country toward the Orange River Syntypes: BMNH 1946.8.6.10-11

Distribution: southern Namibia, Botswana and southwestern South Africa

Omanosaura Lutz, Bischoff & Mayer, 1986.

Lutz, D., W. Bischoff & W. Mayer. 1986.



Chemosystematische Untersuchungen zur Stellung von *Lacerta jayakari* Boulenger, 1887, sowie der Gattungen *Gallotia* Boulenger und *Psammodromus* Fitzinger (Sauria; Lacertidae). *Zeitschrift für Zoologische Systematik und Evolutionsforschung* 24: 144-157 Type species: *Lacerta jayakari* Boulenger, 1887

Omanosaura cyanura  Sindaco

Omanosaura cyanura (Arnold, 1972)

Lacerta cyanura Arnold, 1972: 116

Type locality: Wadi Shawkah (25°06' N, 50°03' E), Trucial States, United Arab Emirates

Holotype: BMNH 1971.1291

Distribution: Hajar Mountains, northern Oman and United Arab Emirates

Omanosaura jayakari (Boulenger, 1887)

Lacerta jayakari Boulenger, 1887: 40

Type locality: near Muscat, Oman

Syntypes: BMNH 1887.11.11.17-20

Distribution: Hajar Mountains, northern Oman and United Arab Emirates

Ophisops Ménétries, 1832

Ménétries, E. 1832. Catalogue raisonné des objets de zoologie recueillis dans



un voyage au caucase et jusqu'aux frontières actuelles de la Perse, par ordre de S.M. L'Empereur. *St.-Petersbourg, de l'Impimerie de l'Academie Impériale des Sciences*, 4to (270 x 212mm). pp. (6), 271, xxxii, iv

Ophisops elegans  Benny Trapp

Ophisops beddomei (Jerdon, 1870)

Pseudophiops beddomei Jerdon, 1870: 72

Type locality: Bramagherry Hills, Wynaad, India

Holotype: BMNH 1946.9.4.5

Synonym:

Pseudophiops monticola Beddome, 1870 Type locality: Bramagherry Hills, Wynaad,
India Syntypes: BMNH 1946.9.4.6-15

Distribution: India: Tamil Nadu, Karnataka, Maharashtra, Goa, Gujarat States

Ophisops elbaensis Schmidt & Marx, 1957

Ophisops elbaensis Schmidt & Marx, 1957: 20

Type locality: Wadi Kansisrob, Gebel Elba, Egypt

Holotype: FMNH 73601

Distribution: Egypt: South Eastern and Central Eastern Deserts; southwestern Saudi Arabia,
Yemen

Ophisops elegans Ménétries, 1832

Ophisops elegans elegans Ménétries, 1832

Ophisops elegans Ménétries, 1832: 63

Type locality: Baku, Azerbaijan

Holotype: ZISP ?

Synonym:

Gymnops meizolepis Stoliczka, 1872 Type locality: SW of Kalabagh, Mianwali Dist.,
Punjab Prov., Pakistan Syntype: ZSI 5207

Distribution: Azerbaijan, Georgia, Iran, northern Iraq, Jordan, Pakistan, southwestern Russia,
eastern Turkey

Ophisops elegans basoglui Baran & Budak, 1978

Ophisops elegans basoglui Baran & Budak, 1978: 186

Type locality: Alanya, southern Anatolia, Turkey

Holotype: SZE 167/1975.1

Distribution: southern Turkey

Ophisops elegans blanfordi Schmidt, 1939

Ophisops elegans blanfordi Schmidt, 1939: 64

Type locality: Halfaya, 20 mi east of Amara, Iraq

Holotype: FMNH 19721

Distribution: southern Iraq, western Iran, Jordan

Ophisops elegans centralanatoliae Bodenheimer, 1944

Ophisops elegans centralanatoliae Bodenheimer, 1944: 33

Type locality: Ankara, Turkey

Holotype: ?

Distribution: Turkey: central, eastern and southern provinces

Ophisops elegans ehrenbergii (Wiegmann, 1835)

Amystes ehrenbergii Wiegmann, 1835: 5

Type locality: Syria

Syntypes: ZMB 1050

Distribution: southern Turkey, western Syria, Lebanon, Israel, western Jordan, Egypt:
northeastern Sinai

Ophisops elegans macrodactylus Berthold, 1840

Ophisops macrodactylus Berthold, 1840: 58

Type locality: Istanbul, Turkey

Holotype: ZFMK 21219

Distribution: southeastern Bulgaria, northeastern Greece, northwestern Turkey

Ophisops elegans persicus Boulenger, 1918

Ophisops elegans persicus Boulenger, 1918: 160

Type locality: Kirjawa, east of Lake Urmi, West Azerbaijan Province, Iran

Syntypes: BMNH 1946.8.4.43-45

Distribution: Iran: West Azerbaijan, Isfahan, Fars, Kerman Provinces

Ophisops elegans schlueteri Boettger, 1880

Ophisops elegans schlueteri Boettger, 1880: 176

Type locality: Cyprus

Lectotype: SMF 13841

Distribution: Cyprus

Ophisops jerdonii Blyth, 1853

Ophisops jerdonii Blyth, 1853: 653

Type locality: Mhow, Indore, Madhya Pradesh, India

Holotype: ZSI 2196

Synonyms:

Pseudophiops theobaldi Jerdon, 1870 Type locality: alpine Punjab, Pakistan Holotype: BMNH 1870.11.29.12

Ophiops bivittata Jerdon, 1870 Type locality: Punjab, Pakistan Holotype: ?

Calosaura chaperi Sauvage, 1884 Type locality: Bellary, Madras, India Holotype: BMNH 1946.8.4.52

Distribution: southeastern Afghanistan, India, Nepal, Pakistan

Ophisops kutchensis Agarwal, Khandekar, Ramakrishnan, Vyas & Giri, 2018

Ophisops kutchensis Agarwal, Khandekar, Ramakrishnan, Vyas & Giri, 2018: 21

Type locality: near Vithon (23.3430556°N 69.354444°E), Kutch district, Gujarat, India

Holotype: NCBS AU760

Distribution: India: Gujarat State: Kutch

Ophisops leschenaultii (Milne-Edwards, 1829)

Ophisops leschenaultii leschenaultii (Milne-Edwards, 1829)

Lacerta leschenaultii Milne-Edwards, 1829: 80

Type locality: Coromandel coast, India

Holotype: MNHNP 0.2753

Synonym:

Cabrita brunnea Gray, 1838 Type locality: unknown Holotype: BMNH, not located

Distribution: southern and eastern India

Ophisops leschenaultii lankae Deraniyagala, 1953

Ophisops leschenaultii lankae Deraniyagala, 1953: ?

Type locality: ?

Holotype: ?

Distribution: eastern Sri Lanka

Ophisops microlepis Blanford, 1870

Ophisops microlepis Blanford, 1870: 351

Type locality: Korba in Biláspúr, the eastern part of Chhatisgarh division, Madhya Pradesh State

Syntypes: BMNH 1946.8.6.12; ZSI 2236

Distribution: India: Gujarat, Bihar, Madhya Pradesh, Rajasthan States

Ophisops minor (Deraniyagala, 1971)

Ophisops minor minor (Deraniyagala, 1971)

Cabrita jerdoni minor Deraniyagala, 1971: 104

Type locality: Bulupitiya-Nilgala area in Uva Province, Sri Lanka
Holotype: NMSL 1971.1.1
Distribution: eastern Sri Lanka: Uva province

Ophisops minor nictans Arnold, 1989
[Ophisops nictans Arnold, 1989: 245](#)
Type locality: Cuddapah, Madras, India
Syntypes: BMNH 1946.9.3.89-92
Distribution: India:

Ophisops occidentalis Boulenger, 1887
[Ophisops occidentalis Boulenger, 1887: 75](#)
Type locality: Matmata, Tunisia
Holotype: BMNH 1885.4.13.3
Distribution: northeaster Morocco, northern Algeria, Tunisia, northern Libya, northwestern Egypt

Ophisops pushkarensis Agarwal, Khandekar, Ramakrishnan, Vyas & Giri, 2018
[Ophisops pushkarensis Agarwal, Khandekar, Ramakrishnan, Vyas & Giri, 2018: 15](#)
Type locality: near Pushkar (26.49318°N 74.56021°E), Rajasthan, India
Holotype: BNHS 2031
Distribution: India: Rajasthan State: near Pushkar and Ajmer

Parvilacerta Arnold, Arribas & Carranza, 2007

Arnold, E.N., O. Arribas & S. Carranza. 2007.



Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* 1430: 1-86
Type species: *Lacerta parva* Boulenger, 1887

Parvilacerta fraasii Wolfgang Bischoff

Parvilacerta fraasii (Lehrs, 1910)
[Lacerta fraasii Lehrs, 1910: 228](#)
Type locality: Cedars at Bscherreh, Lebanon
Holotype: SMNS 1446
Distribution: Lebanon: Mount Lebanon Range; Syria: Anti-Lebanon Mountains

Parvilacerta parva (Boulenger, 1887)
[Lacerta parva Boulenger, 1887: 22](#)
Type locality: Kayseri, Central Anatolia, Turkey
Holotype: BMNH 1946.8.6.47
Distribution: Turkey: central and eastern Anatolia; southern Armenia

Pedioplanis Fitzinger, 1843

Fitzinger, L. 1843. *Systema Reptilium*, Fasciculus Primus, Amblyglossae.

Vienna: Braumüller & Seidel. 106 pp

Type species: *Eremias burchelli*

Duméril & Bibron, 1839



Pedioplanis burchelli J C Els

Pedioplanis benguellensis (Bocage, 1867)

Eremias benguellensis Bocage, 1867: 229

Type locality: Benguela, Angola

Syntype: ZMB 7762

Distribution: southwestern Angola, northwestern Namibia

Pedioplanis breviceps (Sternfeld, 1911)

Eremias breviceps Sternfeld, 1911: 404

Type locality: Walvis Bay, Namibia

Holotype: ZMB 23445

Distribution: northwestern Namibia

Pedioplanis burchelli (Duméril & Bibron, 1839)

Eremias burchelli Duméril & Bibron, 1839: 303

Type locality: Karoo flats, South Africa

Holotype: BMNH 1946.8.7.82

Synonym:

Eremias burchelli quinquevittata Hewitt, 1926 Type locality: Matroosberg, Ceres Dist., Western Cape Prov., South Africa Holotype: SAM 14342

Distribution: South Africa: southern and eastern; Lesotho

Pedioplanis gaerdesi (Mertens, 1954)

Eremias undata gaerdesi Mertens, 1954: 176

Type locality: Tsisab Canyon, Brandberg, Damaraland, Namibia

Holotype: SMF 45954

Distribution: northwestern Namibia

Pedioplanis haackei Conradie, Measey, Branch & Tolley, 2012

Pedioplanis haackei Conradie, Measey, Branch & Tolley, 2012: 11

Type locality: along the road to Tambo (15°52'33.8''S, 12°12'21.0''E), Namibe Province, Angola

Holotype: PEM R18465

Distribution: southwestern Angola: Namibe Province

Pedioplanis huntleyi Conradie, Measey, Branch & Tolley, 2012

Pedioplanis huntleyi Conradie, Measey, Branch & Tolley, 2012: 15

Type locality: road to Oncocua, 7 km from Iona (16°51'29.9'' S, 12°36'45.9'' E), Namibe Province, Angola

Holotype: PEM R18479

Distribution: Angola: Namibe, Cunene Provinces

Pedioplanis husabensis Berger-Dell'mour & Mayer, 1989

Pedioplanis husabensis Berger-Dell'mour & Mayer, 1989: 89

Type locality: northern side of Husab Mountain, Namib Park (22° 40' S, 15° 08' E), Swakopmund Dist., Namibia

Holotype: NMNW 4424

Distribution: Namibia: central Namib Desert

Pedioplanis inornata (Roux, 1907)

Eremias inornata Roux, 1907: 427

Type locality: Orange River, Little Namaqualand, Northern Cape Prov., South Africa

Lectotype: ZMA 11049

Distribution: southern Namibia; South Africa: Northern Cape Province

Pedioplanis laticeps (Smith, 1838)

Eremias laticeps Smith, 1839: pl 46

Type locality: mouth of the Orange River, Northern Cape Prov., South Africa

Holotype: BMNH 1946.8.7.56

Distribution: South Africa: Northern Cape, Western Cape, Eastern Cape Provinces; southern Namibia.

Pedioplanis lineocellata (Duméril & Bibron, 1839)

Pedioplanis lineocellata lineocellata (Duméril & Bibron, 1839)

Eremias lineocellata Duméril & Bibron, 1839: 314

Type locality: South Africa

Syntypes: MNHNP 0.2788-89, 1999.8297

Synonym:

Eremias aspera Boulenger, 1917 Type locality: Mochudi, Botswana Syntypes: BMNH, not located

Distribution: eastern Namibia, Botswana, Lesotho, South Africa: North West, Free State Provinces.

Pedioplanis lineocellata inocellata (Mertens, 1955)

Eremias lineocellata inocellata Mertens, 1955: 66

Type locality: between Bay of Lüderitz and Aus, Great Namaqualand, Namibia

Holotype: SMF 13893

Distribution: coastal southern Namibia

Pedioplanis lineocellata pulchella (Gray, 1845)

Eremias pulchella Gray, 1845: 42

Type locality: South Africa

Holotype: BMNH 1946.8.7.39

Synonyms:

Eremias annulifera Smith, 1845 Type locality: interior of South Africa Holotype: BMNH 1946.8.7.24

Eremias formosa Smith, 1845 Type locality: vicinity of the Orange River, South Africa
Holotype: BMNH 1946.8.7.38

Eremias pulchra Smith, 1845 Type locality: dry flats in the interior of South Africa
Holotype: BMNH 1946.8.7.36

Distribution: southwestern Namibia; South Africa: Northern Cape, Western Cape, northwestern Eastern Cape Provinces

Pedioplanis namaquensis (Duméril & Bibron, 1839)

Eremias namaquensis Duméril & Bibron, 1839: 307

Type locality: Namaqualand, probably Namibia

Syntypes: BMNH 1946.8.7.83-95

Synonym:

Eremias namaquensis quadrangularis Hewitt, 1926 Type locality: Koffiefontein, Free State, South Africa. Holotype: SAM 16128

Distribution: southern Angola, Namibia, Botswana, South Africa: North West, Northern Cape, Western Cape, western Free State, western Eastern Cape Provinces

Pedioplanis rubens (Mertens, 1954)

Eremias undata rubens Mertens, 1954: 177

Type locality: Okatjikona Farm, Waterberges, Namibia

Holotype: SMF 45943

Distribution: Namibia: Waterberg Plateau

Pedioplanis undata (Smith, 1838)

Lacerta undata Smith, 1838: 93

Type locality: Windhoek, Namibia

Neotype: NHMW 318886

Distribution: northern and central Namibia, southern Angola

Philochortus Matschie, 1893 Matschie, P. 1893. Über einige von Herrn Oscar Neumann bei Aden



gesammelte und beobachtete Säugethiere, Reptilien und Amphibien.

Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin (1893): 24-31. Type species: *Philochortus neumannii* Matschie, 1893

Philochortus intermedius Tomas Mazuch

Philochortus hardeggeri (Steindachner, 1891)

Latastia hardeggeri Steindachner, 1891: 371

Type locality: Harar on the way from Heusa, Ethiopia

Syntypes: NMW 16810:1,2

Synonyms:

Eremias heterolepis Boettger, 1893 Type locality: Lafarug, Somalia Holotype: SMF 59441

Latastia degeni Boulenger, 1903 Type locality: Mandah, Somalia Holotype: BMNH
1946.9.3.41

Distribution: Somalia, Djibouti, eastern Ethiopia

Philochortus intermedius Boulenger, 1917

Philochortus intermedius Boulenger, 1917: 152

Type locality: Northern Somaliland at and near Berbera

Syntypes: BMNH 1946.9.3.26-35

Distribution: Djibouti, Ethiopia, Somalia

Philochortus neumanni Matschie, 1893

Philochortus neumanni Matschie, 1893: 30

Type locality: Scadi, near Lahij, Yemen

Holotype: ZMB 11148

Distribution: southwestern Saudi Arabia, Yemen

Philochortus phillipsi (Boulenger, 1898)

Latastia phillipsi Boulenger, 1898: 131

Type locality: Berbera, northern Somalia

Holotype: BMNH 1898.5.21.16

Distribution: Djibouti, northern Ethiopia, northern Somalia

Philochortus rudolfensis Parker, 1932

Philochortus intermedius rudolfensis Parker, 1932: 226

Type locality: mouth of Kaliokwel River, Lake Rudolf, Kenya

Holotype: BMNH 1946.9.3.44

Distribution: southern Ethiopia, northern Kenya

Philochortus spinalis (Peters, 1874)

Lacerta spinalis Peters, 1874: 369

Type locality: supposedly from Bogos, Eritrea

Holotype: ZMB 7633

Distribution: Djibouti, Eritrea, northern Ethiopia, northern Somalia

Philochortus zolii Scortecci, 1934

Philochortus zolii Scortecci, 1934: 305

Type locality: El Barkat, Ghat, Libya

Holotype: MCSN, destroyed in WWII

Synonym:

Philochortus lhotei Angel, 1936 Type locality: vicinity of Abezon, Hoggar, Mali

Holotype: MNHNP 1936.86

Distribution: northern Egypt, northern Libya, Mali, Niger

Phoenicolacerta Arnold, Arribas & Carranza, 2007



Arnold, E.N., O. Arribas & S. Carranza. 2007.

Systematics of the Palaearctic and Oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* 1430: 1-86
Type species: *Lacerta laevis* Gray, 1838

Phoenicolacerta laevis A&S Troidi

Phoenicolacerta cyanisparsa (Schmidtler & Bischoff, 1999)

Lacerta cyanisparsa Schmidtler & Bischoff, 1999: 132

Type locality: Al Bara, south of Ariha, Syria.

Holotype: ZSM 12/1999

Distribution: northwestern Syria, southeastern Turkey

Phoenicolacerta kulzeri (Müller & Wettstein, 1932)

Phoenicolacerta kulzeri kulzeri (Müller & Wettstein, 1932)

Lacerta kulzeri Müller & Wettstein, 1932: 219

Type locality: cedar forest near Bcharré, Lebanon

Holotype: ZSM 10/1931

Distribution: Lebanon: Mount Lebanon Range; Syria: Anti-Lebanon Mountains, and Jabal Al Arab (Druz)

Phoenicolacerta kulzeri khazaliensis Modrý, Necas, Rifai, Bischoff, Hamidan & Amr, 2013

Phoenicolacerta kulzeri khazaliensis Modrý, Necas, Rifai, Bischoff, Hamidan & Amr, 2013: 309

Type locality: Wadi Ramm, Djebel Khazali, Hashemite Kingdom of Jordan

Holotype: ZFMK 73470

Distribution: southwestern Jordan: Djebel Khazali

Phoenicolacerta kulzeri petraea (Bischoff & Müller, 1999)

Lacerta kulzeri petraea Bischoff & Müller, 1999: 246

Type locality: Petra, Jordan

Holotype: ZFMK 65019

Distribution: southwestern Jordan: highlands Esh Shar

Phoenicolacerta laevis (Gray, 1838)

Lacerta laevis Gray, 1838: 279

Type locality: unknown

Holotype: BMNH 1946.9.3.2

Synonym:

Lacerta judaica Camerano, 1877 Type locality: vicinity of Jerusalem Syntypes: MSNTO ?

Distribution: southern Turkey, western Syria, Lebanon, northern Israel, northwestern Jordan

Phoenicolacerta troodica (Werner, 1936)

Lacerta laevis troodica Werner, 1936: 655

Type locality: Platraes, Cyprus

Syntypes: NHMW 16015:1-5

Distribution: Cyprus

Podarcis Wagler, 1830. Wagler, J. 1830. Natürliches System der Amphibien, mit vorangehender



Classification der Säugthiere und Vögel.
Ein Beitrag zur vergleichenden Zoologie.
Munich, Stuttgart and Tübingen: J.G.
Cotta. vi + 354 pp. + one plate. Type
species: *Seps muralis* Laurenti 1768

Podarcis pityusensis formenterae
Manfred Rogner

Podarcis bocagei (Seoane, 1885)

Lacerta muralis bocagei Seoane, 1885: 18

Type locality: La Coruña, northeastern Spain

Lectotype: BMNH 1946.9.1.23

Distribution: northern Spain, northern Portugal

Podarcis carbonelli Perez Mellado, 1981

Podarcis carbonelli carbonelli Perez Mellado, 1981

Podarcis bocagei carbonelli Perez Mellado, 1981: 262

Type locality: Laguna de San Marcos, en el Término Municipal de La Alberca
Provincia de Salamanca, Spain

Holotype: VPM 07010680

Distribution: western and central Portugal; southwestern Spain

Podarcis carbonelli berlengensis Vicente, 1985

Podarcis bocagei berlengensis Vicente, 1985: 371

Type locality: Alto do Cemitério, Ile de Berlenga, Portugal

Holotype : MNHNP 1982.995

Distribution: Portugal: Berlenga Islands and Peniche Peninsula

Podarcis cretensis (Wettstein, 1952)

Lacerta erhardii cretensis Wettstein, 1952: 252

Type locality: Kastelli Kissamoui, northwestern Crete, Greece

Holotype: NHMW 8272

Synonyms:

Podarcis erhardii elaphonisii Wettstein, 1952 Type locality: Elafonisi Island, SW of Crete,
Greece Holotype: NHMW 8298

Podarcis erhardii leukaorii Wettstein, 1952 Type locality: Samaria, Crete, Greece Holotype: NHMW 8291

Podarcis erhardii punctigularis Wettstein, 1952 Type locality: Klippe, Prasoni Island, southwestern Crete, Greece Holotype: NHMW 8296

Podarcis erhardii rechingeri Wettstein, 1952 Type locality: Dragonada Island, Dionysades group, NE of Crete, Greece Holotype: NHMW 8307

Podarcis erhardii schiebeleri Wettstein, 1952 Type locality: Dhia Island, near Heraklion, Crete, Greece Holotype: NHMW 8304

Distribution: Greece: western Crete, and satellite islets: Elaphonisos, Artemis, Gaidouronisi, Mikronisi, Kavalloi, Elasa, Paximada, Dragonada, Prassonisi, Avgo, Dia, Karga, Ag. Nikolaos, Agio Theodoroi

Podarcis erhardii (Bedriaga, 1882)

Podarcis erhardii erhardii (Bedriaga, 1876)

[*Lacerta muralis erhardii* Bedriaga, 1876: 266](#)

Type locality: Seriphos Island, Cyclades, Greece

Holotype: not located

Distribution: Greece: Cyclades: Sifnos and Seriphos Islands

Podarcis erhardii amorgensis (Werner, 1933)

[*Lacerta erhardii amorgensis* Werner, 1933: 110](#)

Type locality: Amorgos Island, SE Cyclade Islands, Greece

Lectotype: NHMW 8261

Distribution: Greece: Cyclades Islands: Amorgos, Anáfi, Pano Koufonissi, Kato Koufonissi, Glaronisi, Keros, Antikeros, Anhydros, Gramvoussa Islands

Podarcis erhardii biinsulicola (Wettstein, 1937)

[*Lacerta erhardii biinsulicola* Wettstein, 1937: 84](#)

Type locality: Makria Island near Anaphi, Aegean Islands, Greece

Lectotype: NHMW 8294

Distribution: Greece: Cyclades: Makra and Mikro Fteno Islands

Podarcis erhardii buchholzi (Wettstein, 1956)

[*Lacerta erhardii buchholzi* Wettstein, 1956: 134](#)

Type locality: Ktenia Island, east of Naxos, Greece

Holotype: NHMW 15230

Distribution: Greece: Cyclades: Ktenia rocks (33 km east of Naxos).

Podarcis erhardii kinarensis (Wettstein, 1937)

[*Lacerta erhardii kinarensis* Wettstein, 1937: 87](#)

Type locality: Kinaros Island, east of Amorgos, SE Cyclade Islands, Greece

Holotype: NHMW 8268

Distribution: Greece: Cyclades: Kinaros Island.

Podarcis erhardii levithensis (Wettstein, 1937)

[*Lacerta erhardii levithensis* Wettstein, 1937: 88](#)

Type locality: Levitha Island, east of Amorgos, Greece

Holotype: NHMW 8300

Distribution: Greece: Cyclades: Levitha Island

Podarcis erhardii livadiaca (Werner, 1902)

Lacerta erhardii livadiaca Werner, 1902: 383

Type locality: central Greece

Holotype: ?

Distribution: central Greece & the Peloponnese, and Hydra Island

Podarcis erhardii makariaisi (Wettstein, 1956)

Lacerta erhardii makariaisi Wettstein, 1956: 135

Type locality: Haga Nikolaos Island, northernmost of the Makariais Islands, east of Naxos, Greece

Holotype: NMW15232

Distribution: Greece: Cyclades: Hagia Nikolaos Island

Podarcis erhardii megalophthenae (Wettstein, 1937)

Lacerta erhardii megalophthenae Wettstein, 1937: 84

Type locality: Megalo Phthena Island, the eastern and larger of the two Phthini Islands, Aegean Sea, Greece

Lectotype: NHMW 8289

Distribution: Greece: Cyclades: Megalo Fteno Island

Podarcis erhardii mykonensis (Werner, 1933)

Lacerta erhardii mykonensis Werner, 1933: 118

Type locality: Mykonos Island, Greece

Lectotype: NHMW 8276

Distribution: Greece: Cyclades: Andros, Tinos, Mykonos, Syros, Dilos, Rinia, Mikro, Megalo Rhemathiaris, and Cherronisi islands

Podarcis erhardii naxensis (Werner, 1899)

Lacerta muralis naxensis Werner, 1899: 835

Type locality: Naxos Island, Greece

Holotype: ?

Distribution: Greece: Cyclades: Naxos, Donousa, Koufonisi, Ios, Sxoinousa Islands; Islets of Paros, Heraklea, Sikinos, Kardiotissa, Pholegandros

Podarcis erhardii ophidusae (Wettstein, 1937)

Lacerta erhardii ophidusae Wettstein, 1937: 87

Type locality: Ofidusa Island, west of Astropalia, Aegean Sea, Greece

Lectotype: NHMW 8266

Synonym:

Lacerta erhardii werneriana Wettstein, 1952 Type locality: Gaidaronisi, near Hierapetra, southeastern Crete, Greece Syntypes: NHMW 8252:18-21

Distribution: Greece: Cyclades: Ofidoussa Island

Podarcis erhardii pachiae (Wettstein, 1937)

Lacerta erhardii pachiae Wettstein, 1937: 84

Type locality: Pachia Island, south of Anaphi, Cyclades, Greece

Lectotype: NHMW 8271

Distribution: Greece: Cyclades: Pachia Island

Podarcis erhardii phytiusae (Wettstein, 1937)

Lacerta erhardii phytiusae Wettstein, 1937: 88

Type locality: Phytiusa Island near the southern tip of Schinusa Island, south of Naxos, Greece
Lectotype: NHMW 8271
Distribution: Greece: Cyclades: Phytiusa Island

Podarcis erhardii riveti (Chabanaud, 1919)

Lacerta muralis riveti Chabanaud, 1919: 23

Type locality: Strkovo, a village east of Lake Prespa, Greece

Syntypes: MNHNP 1918.110-112

Distribution: southern Bulgaria; southern Serbia; Macedonia; Albania; Greece: Epirus, Macedonia, Thrace

Podarcis erhardii ruthveni (Werner, 1930)

Lacerta erhardii ruthveni Werner, 1930: 13

Type locality: Kyra Panagia Island, Bay of Hagios Petros, Northern Sporades, Greece.

Holotype: UMMZ 65516

Distribution: Greece: Northern Sporades

Podarcis erhardii subobscura (Wettstein, 1937)

Lacerta erhardii subobscura Wettstein, 1937: 86

Type locality: southwestern, largest island of the Tria Nisia, south of Syrina, Aegean Sea, Greece

Lectotype: NHMW 8302

Distribution: Greece: Cyclades: Tria Nisia islands

Podarcis erhardii syrinae (Wettstein, 1937)

Lacerta erhardii syrinae Wettstein, 1937: 85

Type locality: Syrina Island, near Astropalia, Cyclades, Greece

Lectotype: NHMW 8283

Distribution: Greece: Cyclades: Astypalaia, Dio Adelfia, Syrna Islands

Podarcis erhardii thermiensis (Werner, 1935)

Lacerta erhardii thermiensis Werner, 1935: 96

Type locality: Kythnos Island, western Cyclades Islands, Greece

Lectotype: NHMW 8281

Distribution: Greece: Cyclades: Kythnos Island

Podarcis erhardii thessalica (Cyrén, 1938)

Lacerta erhardii thessalica Cyrén, 1938: 78

Type locality: Sparmos, Mount Olympus, Thessaly Province, Greece

Holotype: NZMS 1690

Distribution: Greece: Thessaly Province & Paros Island

Podarcis erhardii zafranae (Wettstein, 1937)

Lacerta erhardii zafranae Wettstein, 1937: 85

Type locality: Megali Zafrana Island, south of Astropalia, Aegean Sea, Greece

Lectotype: NHMW 8287

Distribution: Greece: Cyclades: Megali Zofrano Island.

Podarcis filfolensis (Bedriaga, 1876)

Podarcis filfolensis filfolensis (Bedriaga, 1876)

Lacerta filfolensis Bedriaga, 1876: 18

Type locality: Filfla, a small island south of Malta

Syntypes: BMNH 1946.9.1.68-69

Distribution: Malta: Filfolia island

Podarcis filfolensis generalensis (Gulia, 1914)

Lacerta generalensis Gulia, 1914: 322

Type locality: Fungus or Generals Rock, a steep little islet, just few yards distant from the western coast of Gozo, and just at the mouth of the inlet called Cala Dwejra.

Holotype: ?

Distribution: Malta: General's Rock (Fungus Island)

Podarcis filfolensis kieselbachi (Fejervary, 1924)

Lacerta muralis kieselbachi Fejervary, 1924: 9

Type locality: St. Paul's Island, St. Paul's Bay, NE Malta

Syntypes: HNM 2794, 2805, 2803

Distribution: Malta: Selmunett (Saint Paul's Island).

Podarcis filfolensis laurentiimulleri (Fejervary, 1924)

Lacerta muralis laurentiimulleri Fejervary, 1924:11

Type locality: Linosa Island, Italy

Holotype: ZCBS/M 1147

Distribution: Italy: Linosa, Lampioni Islands

Podarcis filfolensis maltensis Mertens, 1921

Podarcis filfolensis maltensis Mertens, 1921: 236

Type locality: Malta

Holotype: SMF 12489

Distribution: Malta: Malta, Gozo, Comino Islands

Podarcis gaigeae (Werner, 1930)

Podarcis gaigeae gaigeae (Werner, 1930)

Lacerta taurica gaigeae Werner, 1930: 9

Type locality: Scyros Island, northern Sporades, Greece

Holotype: UMMZ 65544

Distribution: Greece: northern Sporades: Skyros archipelago and Piperi Island

Podarcis gaigeae weigandi (Gruber & Schultze-Westrum, 1971)

Lacerta erhardii weigandi Gruber & Schultze-Westrum, 1971: 120

Type locality: Piperi Island, northern Sporades, Greece

Holotype: ZFMK 10914

Distribution: Greece: northern Sporades: Piperi Island

Podarcis guadarramae (Boscá, 1916)

Podarcis guadarramae guadarramae (Boscá, 1916)

Lacerta muralis guadarramae Boscá, 1916: 330

Type locality: village of Lozoya (40.951°N, 3.792°W), Sierra de Guadarrama, province of Madrid, Spain

Neotype: MNHNP 2012.0262

Distribution: central Spain

Podarcis gadarramae lusitanicus Geniez, Sá-Sousa, Guillaume, Cluchier, Crochet, 2014
[Podarcis gadarramae lusitanicus Geniez, Sá-Sousa, Guillaume, Cluchier, Crochet, 2014: 14](#)
Type locality: 1 km past Âncora towards Póvoa de Varzim (41.794°N, 8.864°W) south of Caminha, district of Viana do Castelo, Portugal
Holotype: MNHNP 2012.0263
Distribution: northern Portugal; northwestern Spain

Podarcis hispanicus (Steindachner, 1870)

[Lacerta oxycephala hispanica Steindachner, 1870: 336](#)
Type locality: Mount Agudo, near Murcia, southeastern, Spain
Lectotype: NHMW 16088:1
Synonym:
Lacerta muralis hesperica Schreiber, 1912 Type locality: Zaragoza, Spain Syntypes: NHMW 16020, 16021:1-4
Distribution: southeastern Spain

Podarcis lewendis Lymberakis, Poulakakis, Kaliontzopoulou, Valakos & Mylonas, 2008

[Podarcis lewendis Lymberakis, Poulakakis, Kaliontzopoulou, Valakos & Mylonas, 2008: 315](#)
Type locality: Islet Pori, 7.4 km N of the island of Antikythira, between Crete and Peloponnesos and islet Lagouvardos (=Poreti) 3.4 km W–SW of Pori.
Holotype: NHMC 80.3.51.830
Distribution: Greece: Islets Pori and Lagouvardos

Podarcis lilfordi (Günther, 1874)

Podarcis lilfordi lilfordi (Günther, 1874)
[Zootoca lilfordi Günther 1874: 158](#)
Type locality: Isla de l'Aire, southeast coast of Minorca, Spain
Syntypes: BMNH 1946.9.1.64-67
Distribution: Spain: Balearic Islands: Menorca: Illa de l'Aire

Podarcis lilfordi addayae (Eisentraut, 1928)

[Lacerta lilfordi addayae Eisentraut, 1928: 122](#)
Type locality: larger of the two Addaya Islands on the east coast of Menorca, Spain
Lectotype: ZMB 36089
Distribution: Spain: Balearic Islands: Menorca: Illa Gran d'Addaia, Illa Petita d'Addaia, Illa de Ses Mones, Illot d'en Carbó Gran and Illot d'en Carbó Petit.

Podarcis lilfordi balearica (Bedriaga, 1879)

[Lacerta muralis balearica Bedriaga, 1879: 221](#)
Type locality: Isla del Rey, Balearic Islands, Spain
Holotype BMNH 1946.9.2.7
Distribution: Spain: Balearic Islands: Menorca: Illa del Rei.

Podarcis lilfordi brauni (Müller, 1927)

[Lacerta lilfordi brauni Müller, 1927: 261](#)
Type locality: Isla del Colon, Menorca, Spain
Holotype: ZSM 1539
Synonym:
Lacerta lilfordi hospitalis Eisentraut, 1928 Type locality: Isla del Hospital, in the port of Mahon, Menorca, Spain Lectotype: ZMB 36060

Distribution: Spain: Balearic Islands: Menorca: Illa d'en Colom

Podarcis lilfordi carbonerae Perez, Mellado & Salvador, 1988

[Podarcis lilfordi carbonerae Perez Mellado & Salvador, 1988: 173](#)

Type locality: Isla Carbonera, Spain

Holotype: ZFMK 11935

Distribution: Spain: Balearic Islands: Menorca: Isla Carbonera

Podarcis lilfordi codrellensis Perez Mellado & Salvador, 1988

[Podarcis lilfordi codrellensis Perez Mellado & Salvador, 1988: 174](#)

Type locality: Escull de Codrell Island, in the southern coast of Menorca, Balearic Islands, Spain

Holotype: MNCN 14207

Distribution: Spain: Balearic Islands: Menorca: Escull de Binicrodrell Gros and Escull de Binicrodrell Petit

Podarcis lilfordi colomi (Salvador, 1979)

[Lacerta lilfordi colomi Salvador, 1980: 491](#)

Type locality: Illot d'Es Colomé, north of Mallorca, Spain

Holotype: ?

Distribution: Spain: Balearic Islands: Mallorca: Illa es Colomer

Podarcis lilfordi conejerae (Müller, 1927)

[Lacerta lilfordi conejerae Müller, 1927: 264](#)

Type locality: Isla Conejera, north from Cabrera, Spain

Neotype: ZFMK 11898

Distribution: Spain: Balearic Islands: Cabrera Archipelago: S'Illa dels Conills and Illa Na Redona

Podarcis lilfordi espongicola (Salvador, 1979)

[Lacerta lilfordi espongicola Salvador, 1979: 181](#)

Type locality: Isla L'Esponge, situated north of Cabrera between Illa des Conis and Na Plana, Islas Baleares, Spain

Holotype: MNCN 14388

Distribution: Spain: Balearic Islands: Cabrera Archipelago: Escull l'Esponja

Podarcis lilfordi estelicola (Salvador, 1979)

[Lacerta lilfordi estelicola Salvador, 1979: 180](#)

Type locality: Estel de Fora, situated south of Cabrera, Islas Baleares, Spain

Holotype: MNCN 13577

Distribution: Spain: Balearic Islands: Cabrera Archipelago: S'Estell des Colls and S'Estell de Fora.

Podarcis lilfordi fahrae (Müller, 1927)

[Lacerta lilfordi fahrae Müller, 1927: 265](#)

Type locality: Isla Horadada, north of Plana, Dabrera, Spain

Holotype: ZSM 1546

Distribution: Spain: Balearic Islands: Cabrera Archipelago: Illot Na Foradada.

Podarcis lilfordi fenni (Eisentraut, 1928)

[Lacerta lilfordi fenni Eisentraut, 1928: 122](#)

Type locality: Isla del Porros, off Cabo Caballería, north coast of Menorca Spain
Lectotype: ZMB 36062
Distribution: Spain: Balearic Islands: Menorca: Illa d'es Porros

Podarcis lilfordi gigliolii (Bedriaga, 1879)

Lacerta muralis gigliolii Bedriaga, 1879: 223

Type locality: Isla Dragonera, Mallorca, Spain
Syntypes: BMNH 1946.9.1.63, 1946.9.1.74
Distribution: Spain: Balearic Islands: Mallorca: Illa Sa Dragonera

Podarcis lilfordi hartmanni (Wettstein, 1937)

Lacerta lilfordi hartmanni Wettstein, 1937: 296

Type locality: Isla Malgrats in the Bay of St. Ponza, SW coast of Mallorca, Spain
Syntypes: NHMW 15684:1-3
Distribution: Spain: Balearic Islands: Mallorca: Illa Malgrats and Illot d'es Conills

Podarcis lilfordi imperialensis (Salvador, 1979)

Lacerta lilfordi imperialensis Salvador, 1979: 179

Type locality: Isla Imperial, the extreme southeast of Cabrera, Islas Baleares, Spain
Holotype: MNCN 14387
Distribution: Spain: Balearic Islands: Cabrera Archipelago: Illot l'Imperial

Podarcis lilfordi jordansi (Müller, 1927)

Lacerta lilfordi jordansi Müller, 1927: 263

Type locality: Isla La Guardia, south of Salinas, Mallorca, Spain
Holotype: ZSM 1541
Distribution: Spain: Balearic Islands: Mallorca: Illa Na Guardis, Illa Na Moltona, Illot Na Pelada and Illot d'en Curt.

Podarcis lilfordi kuligae (Müller, 1927)

Lacerta lilfordi kuligae Müller, 1927: 263

Type locality: Isla Cabrera, south of Mallorca, Spain
Holotype: ZSM 1543
Synonym:

Lacerta lilfordi mulleri Eisentraut, 1928 Type locality: Isla Cabrera, with the exception of the lighthouse peninsula Holotype: ZMB 36042

Distribution: Spain: Balearic Islands: Cabrera Archipelago: Cabrera, S'Illa de ses Rates, Illa d'es Fonoll and Illot d'es Fonoll.

Podarcis lilfordi nigerrima (Salvador, 1979)

Lacerta lilfordi nigerrima Salvador, 1979: 178

Type locality: Isla Sas Bledas, on the east coast of Cabrera, Balearic Islands, Spain
Holotype: MNCN 14386
Distribution: Spain: Balearic Islands: Cabrera Archipelago: S'Illa de ses Bledes.

Podarcis lilfordi planae (Müller, 1927)

Lacerta lilfordi planae Müller, 1927: 265

Type locality: Isla Plana, north of Conejera, Cabrera, Spain
Holotype: ZSM 1549
Distribution: Spain: Balearic Islands: Cabrera Archipelago: Illot Na Plana.

Podarcis lilfordi pobrae (Salvador, 1979)

Lacerta lilfordi pobrae Salvador, 1979: 182

Type locality: Isla Na Pobra, situated to the north of Illa Des Conis, between Na Foradada and Na Plana, Islas Baleares, Spain.

Holotype: MNCN 14391

Distribution: Spain: Balearic Islands: Cabrera Archipelago: Illot Na Pobra.

Podarcis lilfordi porrosicola Perez Mellado & Salvador, 1988

Podarcis lilfordi porrosicola Perez Mellado & Salvador, 1988: 174

Type locality: Illa des Porros, in the Bay of Fornells, Menorca, Balearic Islands, Spain.

Holotype: MNCN 14194

Distribution: Spain: Balearic Islands: Menorca: Illot d'es Porros.

Podarcis lilfordi rodriguezii (Müller, 1927)

Lacerta lilfordi rodriguezii Müller, 1927: 261

Type locality: Isla des Ratas in the Bay of Mahon, Menorca

Holotype: ZSM 1540

Distribution: Spain: Balearic Islands: Menorca: Isla de Ratas; probably extinct

Podarcis lilfordi sargantanae (Eisentraut, 1928)

Lacerta lilfordi sargantanae Eisentraut, 1928: 465

Type locality: Isla Sargantana and Isla Robello in the Bay of Fornells, north coast of Menorca, Spain

Lectotype: ZMB 52784

Distribution: Spain: Balearic Islands: Menorca: Illa de ses Bledes, Illa Sargantana, Illa d'es Ravells and Illot d'en Tosqueta

Podarcis lilfordi toronis (Hartmann, 1953)

Lacerta lilfordi toronis Hartmann, 1953: 96

Type locality: Lighthouse island, Isla Toro near Mallorca, Spain

Holotype: ZMF 51891

Distribution: Spain: Balearic Islands: Mallorca: Illa Toro.

Podarcis lilfordi xapaticola (Salvador, 1979)

Lacerta lilfordi xapaticolus Salvador, 1979: 179

Type locality: Isla Xapat Gros, south of Cabrera, Islas Baleares, Spain

Holotype: MNCN 13578

Distribution: Spain: Balearic Islands: Cabrera Archipelago: El Carabassot de S'Estell d'en Terra, S'Estell d'en Terra and S'Estell de s'Esclata Sang

Podarcis liolepis (Boulenger, 1905)

Podarcis liolepis liolepis (Boulenger, 1905)

Lacerta muralis liolepis Boulenger, 1905: 363

Type locality: Valencia, Spain

Lectotype: BMHN 1886.12.29.3

Distribution: northeastern Spain, southern France.

Podarcis liolepis atratus (Boscá, 1916)

Lacerta hispanica atrata Boscá, 1916: 330

Type locality: Les Ferreres, Columbretes Island, Spain

Holotype: ?

Synonyms:

Lacerta atrata columbretensis Eisentraut, 1930 Type locality: Columbrete Grande, Spain

Holotype: ZMB 36076

Lacerta atrata mancolibrensis Eisentraut, 1930 Type locality: Isla Mancolibre, near

Columbretes, Spain Holotype: ZMB 36074

Distribution: Spain: Columbretes Archipelago

Podarcis liolepis cebennensis Geniez & Guillaume, 1986

Podarcis hispanica cebennensis Geniez & Guillaume, 1986:81

Type locality: Cap de Coste, between Mandagout and the Col de la Lusette, Massif de l'Aigoual, southern France

Holotype: MNHNP 1986.966

Distribution: southern France, northeastern Spain.

Podarcis liolepis sebastiani (Klemmer, 1964)

Lacerta muralis sebastiani Klemmer, 1964:493

Type locality: Monte Urgull, in San Sebastián, northwestern Spain

Holotype: SMF 59995

Distribution: Spain: Isla Santa Clara and Monte Urgull, San Sebastian.

Podarcis melisellensis (Braun, 1877)

Podarcis melisellensis melisellensis (Braun, 1877)

Lacerta melisellensis Braun, 1877: 50

Type locality: Melisello, near Lissa, Croatia, in the Adriatic Sea

Holotype: not located

Distribution: Croatia: Vis Archipelago

Podarcis melisellensis aeoli (Radovanovic, 1959)

Lacerta melisellensis aeoli Radovanovic, 1959: 412

Type locality: Mali Opuh Island, southeastern Kornati Archipelago, Croatia

Holotype: SMF 57701

Distribution: Croatia: Kornati Archipelago

Podarcis melisellensis bokicae (Radovanovic, 1956)

Lacerta melisellensis bokicae Radovanovic, 1956: 31

Type locality: Vrtlac Island, near Zirje, Croatia

Holotype: NHMW 15517

Distribution: Croatia: Vrtlac Islands

Podarcis melisellensis caprina (Radovanovic, 1970)

Lacerta melisellensis caprina Radovanovic, 1970: 66

Type locality: Kaprije Island, near Šibenik, Croatia

Holotype: ?

Distribution: Croatia: Šibenik Archipelago

Podarcis melisellensis curzolenis (Taddei, 1950)

Lacerta melisellensis curzolenis Taddei, 1950: 161

Type locality: Isola di San Massimo, Korčula Island, Croatia

Syntypes: MZUF 17478-88

Distribution: Croatia: Korčula Island

Podarcis melisellensis digenea (Wettstein, 1926)

Lacerta fiumana digenea Wettstein, 1926: 279

Type locality: Sveti Andrija Island, near Vis Island, Croatia

Syntypes: NHMW 16033:1-4

Distribution: Croatia: Sveti Andrija Island

Podarcis melisellensis fiumana (Werner, 1891)

Lacerta muralis fiumana Werner, 1891: 753

Type locality: Fiume near Cherso, Veglia, Brazza, Lesina, Croatia

Syntypes: NHMW 692, 696

Distribution: Italy: Trieste; Croatian coast, Bosnia-Herzegovina, Montenegro, northwestern Albania

Podarcis melisellensis galvagnii (Werner, 1908)

Lacerta melisellensis galvagnii Werner, 1908: ?

Type locality: Kamik Island, near Svetak, Croatia

Holotype: ?

Distribution: Croatia: Kamik Island

Podarcis melisellensis gigantea (Radovanovic, 1956)

Lacerta melisellensis gigantea Radovanovic, 1956: 21

Type locality: St. Andrea, near Dubrovnik, Croatia

Holotype: NHMW 15515

Distribution: Croatia: Sveti Andrija

Podarcis melisellensis gigas (Wettstein, 1926)

Lacerta fiumana gigas Wettstein, 1926: 286

Type locality: Scoglio Mali, Parsanj near Lissa

Syntypes: NHMW 16042:1,2; NMW 16043:1-6; NMW 16044:1-13

Distribution: Croatia: Lissa Archipelago

Podarcis melisellensis jidulae (Radovanovic, 1959)

Lacerta melisellensis jidulae Radovanovic, 1959: 402

Type locality: Jidula Island, Croatia

Holotype: SMF 57712

Distribution: Croatia; Srednji Archipelago

Podarcis melisellensis kammereri (Wettstein, 1926)

Lacerta fiumana kammereri Wettstein, 1926: 285

Type locality: Scoglio Mali, Barjak near Lissa

Syntypes: NMW 16050:1-2; NMW 16051:1-6

Distribution: Croatia: Lissa Archipelago

Podarcis melisellensis kornatica (Radovanovic, 1959)

Lacerta melisellensis kornatica Radovanovic, 1959: 405

Type locality: Purara Island, Kornati Archipelago

Holotype: SMF 57718

Distribution: Croatia: Kornati Archipelago

Podarcis melisellensis lissana (Werner, 1891)

Lacerta muralis lissana Werner, 1891: 752

Type locality: Vis Island, Croatia

Syntypes: NHMW 16049.1-2

Distribution: Croatia: Vis Island

Podarcis melisellensis lupa (Radovanovic, 1970)

Lacerta melisellensis lupa Radovanovic, 1970: 62

Type locality: Kurjak Island, Croatia

Holotype: ?

Distribution: Croatia: Kornati Archipelago

Podarcis melisellensis mikavicae (Radovanovic, 1959)

Lacerta melisellensis mikavicae Radovanovic, 1959: 414

Type locality: Mikavica Island, north of Zirje Island, Croatia

Holotype: SMF 57722

Distribution: Croatia: Šibenik Archipelago

Podarcis melisellensis plutonis (Radovanovic, 1959)

Lacerta melisellensis plutonis Radovanovic, 1959: 415

Type locality: Jerolim Island, near Hvar, Croatia.

Holotype: ?

Distribution: Croatia: Paklinski Islands

Podarcis melisellensis pomoensis (Wettstein, 1926)

Lacerta fiumana pomoensis Wettstein, 1926: 288

Type locality: Pomo Cliff, near Vis, Croatia

Syntypes: NHMW 16052:1,2; NHMW 11276:1-5

Distribution: Croatia: Vis Island

Podarcis melisellensis thetidis (Radovanovic, 1959)

Lacerta melisellensis thetidis Radovanovic, 1959: 414

Type locality: Veliki Opuh Island, Croatia

Holotype: SMF 57715

Distribution: Croatia: southeastern Kornati Archipelago

Podarcis melisellensis traguriana (Radovanovic, 1959)

Lacerta melisellensis traguriana Radovanovic, 1959: ?

Type locality: near Split, Dalmatian Coast, Croatia

Holotype: ?

Distribution: Croatia: central coast

Podarcis milensis (Bedriaga, 1882)

Podarcis milensis milensis (Bedriaga, 1882)

Lacerta muralis fusca Bedriaga, 1882: 98

Type locality: Milos, the Cyclades, Greece

Syntypes: BMNH 1946.9.1.59-60

Distribution: Greece: Milos Archipelago: Milos, Kimolos, Polyaiagos and Antimilos Islands.

Podarcis milensis adolfjordansi (Buchholz, 1962)

Lacerta muralis adolfjordansi Buchholz, 1962: 218

Type locality: Ananes Island, Cyclades, Greece

Holotype: ZFMK 1854

Distribution: Greece: Ananes Archipelago

Podarcis milensis gerakuniae (Müller, 1938)

Lacerta erhardi gerakuniae Muller, 1938: 39

Type locality: Gerakúnia Island, northwesst of Erimomilos, Cyclades, Greece

Holotype: ZSM 2409

Distribution: Greece: Falkonera and Velopoula Islands

Podarcis muralis (Laurenti, 1768)

Podarcis muralis muralis (Laurenti, 1768)

Seps muralis Laurenti, 1768: 61

Type locality: unknown

Holotype: not located

Synonym:

Podarcis muralis rubriventris Bonaparte, 1836 Type locality: Maritime Alps, Piedmont, Italy

Holotype: unknown

Distribution: Austria, Slovakia, Czech Republic, Poland, Hungary, Romania, southwestern Ukraine, northern Croatia, Slovenia, Italy: Alps

Podarcis muralis acrolampra (Costa, 1839)

Lacerta acrolampra Costa, 1839: 75

Type locality: Aspromontes, Calabria, southern Italy

Holotype: ?

Synonym:

Lacerta muralis appenninica Taddei, 1949 Type locality: Montalto, Aspromontes, Calabria, Italy Syntypes: MZUF 9592-96

Distribution: Italy: Apennine Mountains

Podarcis muralis albanica (Bolkay, 1919)

Lacerta muralis albanica Bolkay, 1919: 12

Type locality: Fieri, southern Albania

Syntypes: NHMW 16096:1-7

Distribution: southern Croatia, Kosovo, Bosnia and Herzegovina, Albania, Serbia, Montenegro, Greece: western & Peloponneses; Bulgaria, Macedonia

Podarcis muralis breviceps (Boulenger, 1905)

Lacerta muralis breviceps Boulenger, 1905: 1

Type locality: Italy

Holotype: BMNH 1946.9.1.61

Distribution: southern Italy

Podarcis muralis brongniardii (Daudin, 1802)

Lacerta brongniardii Daudin, 1802: 221

Type locality: Fontainebleau, France

Syntypes: MNHNP 0.2719-21

Synonyms:

Lacerta merremia Risso, 1826 Type locality: Nice, France Holotype: ?

Lacerta muralis occidentalis Knoepffler & Sochurek, 1956 Type locality: headwaters of the Têt River, Pyrénées-Orientales, France Holotype: NHMW 15205

Distribution: Spain: central, eastern Spain, northwestern; Andorra, France, Italy: western Liguria; Switzerland, Belgium, Luxembourg, Netherlands, western Germany

Podarcis muralis brueggemanni (Bedriaga, 1879)

Lacerta muralis brueggemanni Bedriaga, 1879: 304

Type locality: Arsenal-Mauern in Spezia, Liguria, Italy

Holotype: BMNH 1946.9.1.73

Distribution: northwestern Italy: southern Liguria, Toscana, Umbria, Lazio Provinces

Podarcis muralis colosii (Taddei, 1949)

Lacerta muralis colosii Taddei, 1949: 30

Type locality: Elba Island, Italy

Syntypes: MZUF 9788-93

Synonyms:

Lacerta muralis baldasseronii Taddei, 1949 Type locality: Isola Palmailoa, near Elba, Italy

Syntypes: MZUF 9777-81

Lacerta muralis marcuccii Lanza, 1956 Type locality: Argentarola Islet, Tuscany, Italy

Holotype: MZUF 489

Distribution: Italy: Tuscan Archipelago

Podarcis muralis insulanica Bedriaga, 1881

Lacerta muralis insulanica Bedriaga, 1881: 101

Type locality: Pianosa, near Elba, Italy

Holotype: ?

Distribution: Italy: Tuscan Archipelago: Isola Pianosa

Podarcis muralis maculiventris (Werner, 1891)

Lacerta muralis maculiventris Werner, 1891: 752

Type locality: Fiume (=Rijeka), Croatia

Syntype: NHMW 16098

Synonym:

Lacerta muralis borromeica Mertens, 1932 Type locality: Isola Madre, Lago Maggiore, Italy

Holotype: SMF 22130

Distribution: northern Italy, southern Switzerland, Slovenia: Istria, northwestern Croatia, southern Germany

Podarcis muralis muellerlorenzi (Taddei, 1949).

Lacerta muralis muellerlorenzi Taddei, 1949: 3

Type locality: Isolotto La Scuola, east side of Pianosa Island, Tuscan Archipelago, Italy

Syntypes: MZUF 9712-19

Distribution: Italy: Tuscan Archipelago

Podarcis muralis nigriventris Bonaparte, 1838

Podarcis muralis nigriventris Bonaparte, 1838: 41

Type locality: Rome, Italy

Syntypes: MNHNP 0.2727, 0.2736, 0.719-20

Synonyms:

Lacerta muralis vinciguerrai Mertens, 1932 Type locality: Isola di Gorgona, Tuscan

Archipelago, Italy Holotype: MSNG 31865

Lacerta muralis beccarii Lanza, 1958 Type locality: Port'Ercole Islet, Tuscany, Italy
Holotype: MZUF 1237

Podarcis muralis parkeri (Mertens, 1926)

Lacerta muralis parkeri Mertens, 1926: 322

Type locality: Isola di S. Stefano, Italy

Holotype: BMNH 1917.5.11. 129

Distribution: Italy: Latina Prov, Tyrrhenian Sea

Podarcis muralis paulinii (Taddei, 1953).

Lacerta muralis paulini Taddei, 1953: 86

Type locality: Monte Argentario, Tuscany, Italy

Syntypes: MZUF 9720-26

Distribution: Italy: Grosseto Province: Monte Argentario

Podarcis muralis sammichelii Lanza, 1976

Podarcis muralis sammichelii Lanza, 1976: 186

Type locality: Isolotto della Lacca, Corsica, France

Holotype: MZUF 19375

Distribution: France: Corsica: Archipelago of Isole Sanguinarie: Isole Porro, Isola Locca

Podarcis muralis tinettoi (Taddei, 1949)

Lacerta bedriagae tinettoi Taddei, 1949: 201

Type locality: Isola del Tinetto, Province of La Spezia, Italy

Syntypes: MZUF 16609-13

Distribution: Italy: Tinetto Island, near La Spezia

Podarcis peloponnesiacus (Bibron & Bory, 1833)

Podarcis peloponnesiacus peloponnesiacus (Bibron & Bory, 1833)

Lacerta peloponnesiaca Bibron & Bory, 1833: 66

Type locality: Mistra, southern Peloponnese, Greece

Holotype: MNHNP 0.2706

Distribution: Greece: western & southern Peloponnese; south-western section
of Athens

Podarcis peloponnesiacus lais (Buchholz, 1960)

Lacerta peloponnesiaca lais Buchholz, 1960: 99

Type locality: Kalavrita, northern Peloponnese, Greece

Holotype: ZFMK 12115

Synonym:

Lacerta peloponnesiaca phryne Buchholz, 1960 Type locality: Velia Mt., southern Kalavrita,
above the forest Holotype: ZFMK 12263

Distribution: Greece: northern Peloponnese

Podarcis peloponnesiacus thais (Buchholz, 1960)

Lacerta peloponnesiaca thais Buchholz, 1960: 102

Type locality: Aesculap sanctuary, Argolis, Greece

Holotype: ZFMK 12261

Distribution: Greece: northeastern Peloponnese and Psili Island

Podarcis pityusensis (Boscá, 1883)

Podarcis pityusensis pityusensis (Boscá, 1883)

Lacerta muralis pityusensis Boscá, 1883: 246

Type locality: Ibiza Island, Pityusic Islands, Spain

Syntypes: BMNH 1946.9.2.1-6

Synonyms:

Lacerta lilfordi miguelensis Eisentrut, 1928 Type locality: Isla del Bosque de San Miguel, in the north coast of Ibiza, in the Bay of San Miguel, Balearic Ids., Spain Lectotype: ZMB 36004

Distribution: Spain: Pityusic Islands: Punta Galera, Mesquida, Calders, Punta de Sa Ferradura, Bosc, Ses Illetes de Purroig, Sal Rossa, Negra de Llevant and Ibiza Islands.

Podarcis pityusensis affinis (Müller, 1927)

Lacerta lilfordi affinis Müller, 1927: 269

Type locality: Isla Malvin Pequeno, near Ibiza, Balearic Ids., Spain

Holotype: ZSM 1547

Distribution: Spain: Pityusic Islands: Isla Malvin Pequeno

Podarcis pityusensis ahorcadosi (Eisentraut, 1930)

Lacerta pityusensis ahorcadosi Eisentraut, 1930: 399

Type locality: Isla de los Ahorcados, Balearic Ids., Spain

Holotype: ZMB 35955

Distribution: Spain: Pityusic islands: Illa Penjats

Podarcis pityusensis algae (Wettstein, 1937)

Lacerta pityusensis algae Wettstein, 1937: 296

Type locality: Isla Alga, north of Formentera, Spain

Syntypes: NHMW 16226:1-8

Distribution: Spain: Pityusic Islands: Formentera: Illa Pouet

Podarcis pityusensis calaesaladae (Müller, 1928)

Lacerta lilfordi calaesaladae Müller, 1928: 387

Type locality: Isla de Cala Salada, west coast of Ibiza, Balearic Ids., Spain

Holotype: ZSM, uncataloged

Distribution: Spain: Pityusic Islands: Ibiza: Illot de Cala Salada

Podarcis pityusensis canaretensis Cirer, 1980

Podarcis pityusensis canaretensis Cirer, 1980: 122

Type locality: Illot d'es Canaret, Balearic Ids., Spain

Holotype: MZB 82-9463

Distribution: Spain: Pityusic Islands: Ibiza: Illot d'es Canaret

Podarcis pityusensis canensis (Eisentraut, 1928)

Lacerta lilfordi canensis Eisentraut, 1928: 466

Type locality: Isla Caná, east side of Ibiza, north of Isla Sta. Eulalia, Balearic Ids., Spain

Lectotype: ZMB 35984

Distribution: Spain: Pityusic Islands: Ibiza: Illa d'es Canar

Podarcis pityusensis caragolensis (Buchholz, 1954)

Lacerta pityusensis caragolensis Buchholz, 1954: 81

Type locality: Isla Caragole, Balearic Ids., Spain

Holotype: ZFMK 12497

Distribution: Spain: Pityusic Islands: Ibiza: Illot d'es Caragoler

Podarcis pityusensis carlkochi (Mertens & Müller, 1940)

Lacerta lilfordi carlkochi Mertens & Müller, 1940: 32

Type locality: Conillera Island, west of Ibiza, Balearic Islands, Spain

Holotype: ZSM 1542

Distribution: Spain: Pityusic Islands: Ibiza: Illa Sa Conillera

Podarcis pityusensis characae (Buchholz, 1954)

Lacerta pityusensis characae Buchholz, 1954: 86

Type locality: Isla Characa, off the north coast of Ibiza, in the Bay of Cala Characa, Balearic Islands, Spain

Holotype: ZFMK 12467

Distribution: Spain: Pityusic Islands: Ibiza: Isla Characa

Podarcis pityusensis espalmadoris (Müller, 1928)

Lacerta lilfordi espalmadoris Müller, 1928: 262

Type locality: Isla Espamador, south of Ibiza, Balearic Islands, Spain

Holotype: ZSM 1630

Distribution: Spain: Pityusic Islands: Es Freus: S`Espalmador

Podarcis pityusensis espardellensis (Eisentraut, 1928)

Lacerta lilfordi espardellensis Eisentraut, 1928: 123

Type locality: Isla Espardell, east of Isla Espamador, between Ibiza and Formentera, Balearic Islands, Spain

Lectotype: ZMB 35965

Distribution: Spain: Pityusic Islands: Es Freus: S`Espardell.

Podarcis pityusensis formenterae (Eisentraut, 1928)

Lacerta lilfordi formenterae Eisentraut, 1928: 123

Type locality: Isla Formentera, Balearic Islands, Spain

Lectotype: ZMB 35979

Distribution: Spain: Pityusic Islands: southern Isla Formentera

Podarcis pityusensis frailensis (Eisentraut, 1928)

Lacerta lilfordi frailensis Eisentraut, 1928: 467

Type locality: Espardell de Espartar, off the west coast of Ibiza, Balearic Islands, Spain

Lectotype: ZMB 36024

Distribution: Spain: Pityusic Islands: Ibiza: S`Espardell de S`Espartar (Fraile).

Podarcis pityusensis gastabiensis (Eisentraut, 1928)

Lacerta lilfordi gastabiensis Eisentraut, 1928: 123

Type locality: Isla Gastabi, between Ibiza and Formentera, southwest of Espamador, Balearic Islands, Spain

Lectotype: ZMB 35299

Distribution: Spain: Pityusic Islands: Es Freus: Illot de Gastavi

Podarcis pityusensis gorrae (Eisentraut, 1928)

Lacerta lilfordi gorrae Eisentraut, 1928: 122

Type locality: Isla Bleda Gorra, Balearic Islands, Spain

Lectotype: ZMB 36019

Distribution: Spain: Pityusic Islands: Bledas Islands: Na Gorra, Es Vaixell and Na Bosc.

Podarcis pityusensis grueni (Müller, 1928)

Lacerta lilfordi grueni Müller, 1928: 295

Type locality: Isla dos Trocadoo, near Espamador, Balearic Islands, Spain

Holotype: ZSM 1632

Distribution: Spain: Pityusic Islands: northern Formentera

Podarcis pityusensis hedwigkammerae (Müller, 1927)

Lacerta lilfordi hedwigkammerae Müller, 1927: 185

Type locality: Isla Margarita, NW Sant Ines, off Ibiza, Balearic Islands, Spain

Holotype: ZSM 1550

Distribution: Spain: Pityusic Islands: Ibiza: Ses Margalides

Podarcis pityusensis hortae (Buchholz, 1954)

Lacerta pityusensis hortae Buchholz, 1954: 86

Type locality: Isla de Hort, Balearic Ids., Spain

Holotype: ZFMK 12410

Distribution: Spain: Pityusic Islands: Ibiza: Illot de S'Hort

Podarcis pityusensis kameriana (Mertens, 1927)

Lacerta pityusensis kameriana Mertens, 1927: 302

Type locality: Isla Esparto, west coast of Ibiza, Pityusic Islands, Spain

Holotype: SMF 21315

Synonym:

Lacerta lilfordi zenonis Müller, 1928 Type locality: Escollo, off Esparta, on the west coast of Ibiza Holotype: ZSM, uncataloged (lost)

Distribution: Spain: Pityusic Islands: Ibiza: Illa S'Espartar and Escui de S'Espartar

Podarcis pityusensis maluquerorum Mertens, 1921

Podarcis pityusensis maluquerorum Mertens, 1921: 142

Type locality: Isla de las Bledas, near Ibiza, Pityusic Islands, Spain

Holotype: SMF 12554

Distribution: Spain: Pityusic Islands: Bledas Islands: Bleda Plana and Escull Vermell.

Podarcis pityusensis muradae (Eisentraut, 1928)

Lacerta lilfordi muradae Eisentraut, 1928: 122

Type locality: Isla Murada, on the west coast of Ibiza, north of Isla Margarita, Pityusic Islands, Spain

Holotype: ZMB 35997

Distribution: Spain: Pityusic Islands: Ibiza: Illa Murada

Podarcis pityusensis negrae (Eisentraut, 1928)

Lacerta lilfordi negrae Eisentraut, 1928: 468

Type locality: Isla Negra, west of Ahorcados, between Ibiza and Formetera, Balearic Ids., Spain

Lectotype: ZMB 35958

Distribution: Spain: Pityusic Islands: Es Freus: Illa Negra Norte

Podarcis pityusensis puercosensis (Buchholz, 1954)

Lacerta pityusensis puercosensis Buchholz, 1954: 77

Type locality: Isla Puercos, Balearic Ids., Spain

Holotype: ZFMK 12411

Distribution: Spain: Pityusic Islands: Es Freus: Illa Pou.

Podarcis pityusensis purroigensis (Buchholz, 1954)

Lacerta pityusensis purroigensis Buchholz, 1954: 85

Type locality: Isla Purroige, Balearic Ids., Spain

Holotype: ZFMK 12409

Distribution: Spain: Pityusic Islands: Isleta de Purroige

Podarcis pityusensis ratae (Eisentraut, 1928)

Lacerta lilfordi ratae Eisentraut, 1928: 466

Type locality: Isla Ratas, southwest of the Bay of Ibiza, Balearic Ids., Spain

Lectotype: ZMB 35951

Distribution: Spain: Pityusic Islands: Ibiza: Illa de ses Rates

Podarcis pityusensis redonae (Eisentraut, 1928)

Lacerta lilfordi redonae Eisentraut, 1928: 123

Type locality: Isla Redona, east of Sta. Eulalia, Ibiza, Balearic Ids., Spain

Lectotype: ZMB 35980

Synonym:

Lacerta lilfordi grossae Müller, 1929 Type locality: Ille La Grossa, east of Sta. Eulalia off Ibiza, Pityusic Islands, Balearic Islands, Spain Holotype: ZSM 1668

Distribution: Spain: Pityusic Islands / Ibiza / Illa Redona de Santa Eulària and Illa Grossa de Santa Eulària

Podarcis pityusensis sabinae (Buchholz, 1954)

Lacerta pityusensis sabinae Buchholz, 1954: 79

Type locality: Isla Sabina, north of the harbor of Formentera, Balearic Ids., Spain

Holotype: ZFMK 12607

Distribution: Spain: Pityusic Islands: Formentera Island: Isla Sabina

Podarcis pityusensis schreitmuelleri (Müller, 1927)

Lacerta lilfordi schreitmülleri Müller, 1927: 268

Type locality: Isla Malvin, near Ibiza, Balearic Islands, Spain

Holotype: ZSM 1544

Distribution: Spain: Pityusic Islands: Ibiza: Illa Malvi Pla and Illa Malvi Rodó

Podarcis pityusensis subformenterae (Buchholz, 1954)

Lacerta pityusensis subformenterae Buchholz, 1954: 78

Type locality: Conejo de Formentera, Formentera Island, Balearic Ids., Spain

Holotype: ZFMK 30229

Distribution: Spain: Pityusic Islands: Isla Formentera

Podarcis pityusensis tagomagensis (Müller, 1927)

Lacerta lilfordi tagomagensis Müller, 1927: 267

Type locality: Isla Tagomago, east of Ibiza, Balearic Islands, Spain

Holotype: ZSM 1545

Distribution: Spain: Pityusic Islands: Ibiza: Illa de Tagomago.

Podarcis pityusensis torretensis (Buchholz, 1954)

Lacerta pityusensis torretensis Buchholz, 1954: 76

Type locality: Isla de Torretas, Balearic Ids., Spain

Holotype: ZFMK 12408

Distribution: Spain: Pityusic Islands: Es Freus: Illa de Sa Torreta.

Podarcis pityusensis vedrae (Müller, 1927)

Lacerta lilfordi vedrae Müller, 1927: 190

Type locality: Isla Vedra, west of the southwestern tip of Ibiza, near Cabo Jueu, Balearic Islands, Spain

Holotype: ZSM 1551

Synonym:

Lacerta lilfordi vedranellensis Müller, 1928 Type locality: Isla Vedranell, on the west coast of Ibiza, Balearic Islands, Spain Holotype: ZSM 2149

Distribution: Spain: Pityusic Islands: Ibiza: Illa Es Vedrà and Illa Es Vedranell

Podarcis raffonei (Mertens, 1952)

Podarcis raffonei raffonei (Mertens, 1952)

Lacerta sicula raffonei Mertens, 1952: 313

Type locality: Strombolicchio Island, northeast of the island of Stromboli, Aeolian Islands, Italy

Holotype: SMF 43919

Distribution: Italy: Aeolian Islands: Strombolicchio islet near Stromboli

Podarcis raffonei alvearioi (Mertens, 1955)

Lacerta sicula alvearioi Mertens, 1955: 31

Type locality: Faraglione di Pollara, northwest coast of Salina, Liparian Islands, Italy

Holotype: SMF 48296

Synonym:

Podarcis siculus cucchiarai Di Palma, 1980 Type locality: Faraglione di La Canna, Isola di Filicudi, Italy Holotype: MZUP 33

Distribution: Italy: Aeolian Islands: La Canna near Filicudi, Scoglio Faraglione near Salina and Vulcanello Peninsula on Vulcano Island

Podarcis raffonei antoninoi (Mertens, 1955)

Lacerta wagleriana antoninoi Mertens, 1955: 32

Type locality: Isola Marettimo, Aegadian Islands, west of Sicily, Italy

Holotype: SMF 50519

Distribution: Italy: Aeolian Islands: Trapani Prov.: Marettimo Island

Podarcis siculus (Rafinesque-Schmaltz, 1810)

Podarcis siculus siculus (Rafinesque-Schmaltz, 1810)

Lacerta sicula Rafinesque-Schmaltz, 1810: 8

Type locality: Sicily, Italy

Holotype: not located

Synonyms:

Podarcis muralis albiventris Bonaparte, 1838 Type locality: Italy Holotype: MNHNP, not located

Lacerta muralis elegans Eimer, 1874 Type locality: Isola Capri Holotype: ?

Lacerta sicula mertensi Wettstein, 1931 Type locality: Paestum, Gulf of Salerno, Italy
Holotype: SMF 12797
Lacerta muralis joachinimurati Taddei, 1949 Type locality: Pizzo di Calabria, Italy Syntypes:
MZUF ?
Lacerta sicula ventotenensis Taddei, 1949 Type locality: Isola di Ventotene, Pontine Islands,
Tyrrhenian Sea Holotype: MZUF, not located
Lacerta sicula liscabiancae Mertens, 1952 Type locality: Lisca Bianca Island, east of Panaria,
Lipari Islands, Italy Holotype: SMF 43901
Lacerta sicula trischittai Mertens, 1952 Type locality: Bottaro Island, between Lisca Bianca and
Lisca Nera Islands, east of Panaria, Lipari Islands, Italy Holotype: SMF 43911
Distribution: southern Italy, Sicily (including Mal di Ventre, San Pietro, San Antioco,
Tavolara, Asinara and Maddalena), Aeolian Islands and Pontine islands

Podarcis siculus adriaticus (Werner, 1902)

Lacerta serpa adriatica Werner, 1902: 386

Type locality: Isole di Pelagosa Piccola, Adriatic Sea, Croatia

Holotype: ?

Distribution: Croatia: Palagruža Archipelago: Isole di Pelagosa Piccola

Podarcis siculus aemiliani Capolongo, 1984

Podarcis sicula aemiliani Capolongo, 1984: 197

Type locality: Scoglio Minore de Apani, near Brindisi

Holotype: DC 6816

Distribution: Italy: Scogli di Àpani: Scoglio Minore de Apani, near Brindisi

Podarcis siculus amparoeae Capolongo, 1979

Podarcis siculus amparoeae Capolongo, 1979: 8

Type locality: Isola di Dino, ca. 2 km south of Praia a Mare (Cosenza), Calabria, Italy

Holotype: MZUF 19840.

Distribution: Italy: Isola di Dino

Podarcis siculus astorgae (Mertens, 1937)

Lacerta sicula astorgae Mertens, 1937: 333

Type locality: Astorga Island, west coast of Istria, Croatia

Holotype: SMF 22314

Distribution: Croatia: Adriatic Sea: Istria: Sturago Island

Podarcis siculus bagnolensis (Mertens, 1937)

Lacerta sicula bagnolensis Mertens, 1937: 332

Type locality: Bagnola Island, west coast of Istria, Croatia

Holotype: SMF 22317

Distribution: Croatia: Adriatic Sea: Istria: Banjol

Podarcis siculus bolei (Breljih, 1961)

Lacerta sicula bolei Breljih, 1961: 78

Type locality: Vrsar, Istria, Tondo piccolo, Orsera beach, Croatia

Holotype: ML 731

Distribution: Croatia: Istria: Tovaez Island

Podarcis siculus calabra (Taddei, 1949)

Lacerta sicula calabra Taddei, 1949: 219

Type locality: Isola di Montecristo, Calabria, Italy
Syntypes: MZUF 1401-07, 14409-17
Distribution: Italy: Calabria : Isola di Montecristo

Podarcis siculus calabresiae (Taddei, 1949)

Lacerta sicula calabresiae Taddei, 1949: 22

Type locality: Isola di Montecristo, Calabria, Italy
Syntypes: SMF 40945-46
Distribution: Italy: Archipelago Toscano, Tirreno Sea : Isola di Montecristo

Podarcis siculus campana (Taddei, 1949)

Lacerta sicula campana Taddei, 1949: 220

Type locality: Campania and Basilicata, southern Italy
Syntypes: MZUF 14232, 14340-42
Distribution: Italy: Nisida, Capri, Procida, Vivara, Ischia Islands

Podarcis siculus campestris (De Betta, 1857)

Lacerta muralis campestris De Betta, 1857: 152

Type locality: Verona, southwestern Veneto, Italy
Syntypes: MSNV 22
Distribution: Italy: Corsica, northern Italy, northern Croatia

Podarcis siculus caporiacoi (Taddei, 1949)

Lacerta sicula catanensis Taddei, 1949: 216

Type locality: Isola Capraia, Italy
Syntypes: : MZUF 9847-54
Distribution: Italy: Isola Capraia

Podarcis siculus catanensis (Taddei, 1949)

Lacerta sicula catanensis Taddei, 1949: 216

Type locality: Catania, Sicily
Syntypes: MZUF 14418, 14542-53
Distribution: Italy: eastern Sicily

Podarcis siculus cattaroi (Taddei, 1950)

Lacerta sicula cattaroi Taddei, 1950: 154

Type locality: Dalmatia, Croatia
Syntypes: MZUF 17341-56
Distribution: Croatia: Dalmatian coast

Podarcis siculus cerbolensis (Taddei, 1949)

Lacerta sicula cerbolensis Taddei, 1949: 28

Type locality: Isla de Cerboli, Tuscan Archipelago, Italy
Lectotype: MZUF 9811
Distribution: Italy: Tuscan Archipelago: Isolotto di Cerboli

Podarcis siculus cettii (Cara, 1872)

Lacerta podarcis cettii Cara, 1872: 30

Type locality: Sardinia, Italy
Holotype: ?

Distribution: Italy: Sardinia & small islands of Asinara, San Pietro, Sant' Antioco, Tavolara;
France: southern Corsica

Podarcis siculus ciclopica (Taddei, 1949)

Lacerta siculus ciclopica Taddei, 1949: 224

Type locality: Isola di Trezza, Catania, Italy

Syntypes: MZUF 17692-701

Distribution: Italy: Isola Lachea (near Acitrezza), and Scoglio della Madonnina (Isole dei Ciclopi), (near Acitrezza)

Podarcis siculus coeruleus (Eimer, 1872)

Lacerta muralis coerulea Eimer, 1872: 11

Type locality: Isole Faraglione de Mezzo, near Capri Island, Italy

Syntypes: ZFMK 21332-21333

Distribution: Italy: Campania: Tyrrhenian Sea :Faraglione di Mezzo and Faraglione di Fuori o Esterno, near l'Isola di Capri.

Podarcis siculus dupinici (Radovanovic, 1956)

Lacerta sicula dupinici Radovanovic, 1956: 42

Type locality: Mali Dupinic, Croatia

Holotype: NHMW 15521

Distribution: Croatia: Dupinic Mali Island

Podarcis siculus fasciatus (Radovanovic, 1956)

Lacerta sicula fasciata Radovanovic, 1956: 49

Type locality: Duzac, Istria, near Zirje, Croatia

Holotype: NHMW 15525

Distribution: Croatia: Otocic Duzac Island

Podarcis siculus fiumanoideus (Brelj, 1963)

Lacerta sicula fiumanoidea Brelj, 1963: 110

Type locality: Kvarner Gulf, Visoki Island, near Cresu, Croatia

Holotype: ML 1447

Distribution: Croatia: Visoki Island

Podarcis siculus flavigulus (Mertens 1937)

Lacerta sicula flavigula Mertens, 1937: 336

Type locality: Isolotto San Giovanni Faro, west coast of Istria

Holotype: SMF 22315

Distribution: Italy: Lecce Province: Isolotto San Giovanni Faro

Podarcis siculus gallensis (Eimer, 1881)

Lacerta muralis gallensis Eimer, 1881: 395

Type locality: Isole Gallo Maggiore, Campania, Italy

Holotype: ?

Distribution: Italy: Campania: Gulf of Salerno: Isole Gallo Maggiore, Isole La Castelluccia

Podarcis siculus hadzii (Brelj, 1961)

Lacerta sicula hadzii Brelj, 1961: 90

Type locality: Porer cliffs, western Istria, Croatia

Holotype: ML, not located
Distribution: Croatia: western Istria: Cape Kamenjak : Porer islet

Podarcis sculus hieroglyphicus (Berthold, 1840)

Lacerta hieroglyphica Berthold, 1840: 54

Type locality: Istanbul, Turkey

Syntype: ZFMK 108205

Distribution: northwestern Turkey

Podarcis sculus insularus (Mertens, 1937)

Lacerta sicula insularum Mertens, 1937: 334

Type locality: La Longa Island, Istria, Croatia

Holotype: SMF 22316

Distribution: Croatia: Istria: Otočić Lunga

Podarcis sculus klemmeri (Lanza & Capolongo, 1972)

Lacerta sicula klemmeri Lanza & Capolongo, 1972: 21

Type locality: Licosa Islet, 50 km south of Salerno, Campania, Italy.

Holotype: MZUF 12923

Distribution: Italy: Isola Licosa

Podarcis sculus kolombatovici (Karaman, 1928)

Lacerta serpa kolombatovici Karaman, 1928: 138

Type locality: Velika Island, Croatia

Holotype: ?

Distribution: Croatia: Velika and Mala Kluda Islands

Podarcis sculus laganjensis (Radovanovic, 1956)

Lacerta sicula laganjensis Radovanovic, 1956: 46

Type locality: Laganj Mali, Croatia

Holotype: NHMW 15523

Distribution: Croatia: peninsula of Istria: Laganj Mali Island

Podarcis sculus lanzai (Mertens, 1967)

Lacerta sicula lanzai Mertens, 1967: 131

Type locality: Isola di Gavi, Pontine Islands, Italy

Holotype: SMF 58827

Distribution: Italy: Isola di Gavi, near Ponza

Podarcis sculus latastei (Bedriaga, 1879)

Lacerta muralis latastei Bedriaga, 1879: 209

Type locality: Isola Ponza, Pontine Archipelago, Italy

Holotype: ?

Distribution: Italy: Isola Ponza and Scoglio della Madonna

Podarcis sculus massinei (Mertens, 1961)

Lacerta sicula massinei Mertens, 1961: 34

Type locality: Isola Rotonda, Li Galli Archipelago, Gulf of Salerno, Italy

Holotype: SMF 56848

Distribution: Italy: Isola Rotonda

Podarcis siculus medemi (Mertens, 1942)

[Lacerta siculus medemi Mertens, 1942: 44](#)

Type locality: Isola Bella, near Taormina. Sicily

Holotype: SMF 34393

Distribution: Italy: Sicily: Isola Bella

Podarcis siculus mediofasciatus (Radovanovic, 1959)

[Lacerta sicula mediofasciata Radovanovic, 1959: 423](#)

Type locality: Dužac Island, west of Isto, Adriatic Sea, Croatia

Holotype: NHMW 15525

Distribution: Croatia: Duzac and Mala Sestrica Islands

Podarcis siculus monaconensis (Eimer, 1881)

[Lacerta muralis monacone Eimer, 1881: 392](#)

Type locality: Monacone Island, near Capri, Italy

Holotype: ?

Distribution: Italy: Campania: Faraglione Monacone, near Isola di Capri

Podarcis siculus muelleri (Taddei, 1949)

[Lacerta sicula muelleri Taddei, 1949: 23](#)

Type locality: La Scuola, Pianosa Island, Archipelago Toscano, Italy

Syntypes: MZUF 9712-19

Distribution: Italy: Archipelago Toscano: Isola Pianosa

Podarcis siculus nikolici (Breljih, 1961)

[Lacerta sicula nikolici Breljih, 1961: 89](#)

Type locality: Gusti Island, western Istria, Croatia

Holotype: ML, not located

Distribution: Croatia: Istria: Gusti Skolj

Podarcis siculus oristanensis (Taddei, 1949)

[Lacerta sicula oristanensis Taddei, 1949: 209](#)

Type locality: L'isola di Mal di Ventre, Oristano Prov., Sardinia, Italy

Syntypes: MZUF 4733

Distribution: Italy: Sardinia: Oristano Prov

Podarcis siculus palmarolae (Mertens, 1967)

[Lacerta sicula palmarolae Mertens, 1967: 134](#)

Type locality: Palmarola, Pontine Islands, Tyrrhenian Sea, Italy

Holotype: SMF 58837

Distribution: Italy: Isola Palmarola and Scoglio di Mezzagiomo.

Podarcis siculus pasquinii (Lanza, 1952)

[Lacerta sicula pasquinii Lanza, 1952: 511](#)

Type locality: Scoglio Cappello, S of Palmarola Island, Latium, Italy

Holotype: MZUF 4350

Distribution: Italy: Scoglio Capello.

Podarcis siculus patrizii (Lanza, 1952)

Lacerta sicula patrizii Lanza, 1952: 69

Type locality: Zannone Island, Latium, Italy

Syntypes: MZUF 8203-13

Distribution: Italy: Isola Zannone

Podarcis siculus paulae (Lanza, Adriani & Romiti, 1971)

Lacerta sicula paulae Lanza, Adriani & Romiti, 1971:5

Type locality: Santo Ianni Islet, Basilicata, Italy

Holotype: MZUF 7323

Distribution: Italy: Isola di Santo Ianni

Podarcis siculus pelagosae (Bedriaga, 1886)

Lacerta muralis pelagosae Bedriaga, 1886: 212

Type locality: Pelagosa Grande (=Vela Palagruža) Island, Adriatic Sea, Croatia

Holotype: ?

Synonym:

Lacerta serpa cazzae Schreiber, 1912 Type locality: Cazza, Dalmatia, Croatia Syntypes:
NHMW 16235:1-13

Distribution: Croatia: Palagruža Archipelago: Vela Palagruža

Podarcis siculus pirosoensis (Mertens, 1937)

Lacerta sicula pirosoënsis Mertens, 1937: 335

Type locality: Piroso Grande Island, west coast of Istria, Croatia

Holotype: SMF 22361

Distribution: Croatia: Veli Piruzi Island

Podarcis siculus pohlibensis (Radovanovic, 1970)

Lacerta sicula pohlibensis Radovanovic, 1970: 61

Type locality: Pohlib Island, Croatia

Holotype: ?

Distribution: Croatia: Otočić Pohlib, Planičić, Hrid Sip, Planik, Olib, Morovnik Islands

Podarcis siculus polenci (Breljih, 1961)

Lacerta sicula polenci Breljih, 1961: 88

Type locality: Calbula Cliffs, West Istria, Croatia

Holotype: ML, not located

Distribution: Croatia: Brijuni Islands: Cabula Island

Podarcis siculus premudanus (Radovanovic, 1959)

Lacerta sicula premudana Radovanovic, 1959: 418

Type locality: Permuda Island, Dalmatian Islands, Croatia

Holotype: SMF 57739

Distribution: Croatia: Dalmatian Islands: Permuda and Masarina Islands

Podarcis siculus premudensis (Radovanovic, 1959)

Lacerta sicula premudensis Radovanovic, 1959: 419

Type locality: Lutrosnjak, northwest of Permuda

Holotype: SMF 55742

Distribution: Croatia: Dalmatian Islands: Otočić Lutrošnjak

Podarcis siculus pretneri (Breljih, 1961)

Lacerta siculus pretneri Breljih, 1961: 90

Type localities: Gustinja and Pisulj Islets, West Istria, Croatia

Syntypes: ML, not located

Distribution: Croatia: West Istria

Podarcis siculus radovanovici (Breljih, 1961)

Lacerta sicula radovanovici Breljih, 1961: 88

Type locality: Orada, near Porec, Croatia

Holotype: ML, not located

Distribution: Croatia: West Istria

Podarcis siculus ragusae (Wettstein, 1931)

Lacerta sicula ragusae Wettstein, 1931: 282

Type locality: Ragusa (= Dubrovnik), Dalmatia, Croatia

Holotype: NHMW 16242

Distribution: Croatia: Dubrovnik

Podarcis siculus roberti (Taddei, 1949)

Lacerta sicula roberti Taddei, 1949: 27

Type locality: Isola Formica Grande di Grosset, Aegadian Islands, Sicily, Italy

Lectotype: MZUF 9979

Distribution: Italy: Sicily: Aegadian Islands

Podarcis siculus salfii (Lanza, 1954)

Lacerta sicula salfii Lanza, 1954: 136

Type locality: isolotto di Vivaro di Nerano, near southern coast of Salento Peninsula, southern Italy

Holotype: MZUF 8159

Distribution: Italy: Scoglio Vivaro di Nerano

Podarcis siculus samogradi (Radovanovic, 1956)

Lacerta sicula samogradi Radovanovic, 1956: 39

Type locality: Samograd, Korcula Island, Croatia

Holotype: NHMW15519

Distribution: Croatia: Korcula Island

Podarcis siculus sanctinicolai (Taddei, 1949)

Lacerta sicula sanctinicolai Taddei, 1949: 235

Type locality: Isola di San Nicola, Isole Tremiti Archipelago, Italy

Syntypes: MZUF 14385-89

Distribution: Italy: Isole Tremiti Archipelago

Podarcis siculus sanctistephani (Mertens, 1926)

Lacerta sicula sanctistephani Mertens, 1926: 320

Type locality: San Stefano Island near Ventotene, Tyrrhenian Sea.

Holotype: BMNH 1946.9.2.18

Distribution: Italy: Maddalena Archipelago: San Stefano Island

Podarcis siculus scopelicus (Taddei, 1950)

Lacerta sicula scopelica Taddei, 1950: 151

Type locality: Scoglio Sant' Andrea, near Vis Island, Croatia

Syntypes: MZUF 17468-72

Distribution: Croatia: Istria: Sveti Andrija Island

Podarcis siculus scortecii (Taddei, 1949)

Lacerta sicula scortecii Taddei, 1949: 24

Type locality: Isola di Giannutri, Tuscan Archipelago, Italy

Lectotype: MZUF 9960

Distribution: Italy: Tuscan Archipelago: Isola di Giannutri

Podarcis siculus strombolensis (Taddei, 1949)

Lacerta sicula strombolensis Taddei, 1949: 217

Type locality: Isola di Stromboli, Tyrrhenian Sea, Italy

Syntypes: MZUF 16795-804

Distribution: Italy: Aeolian Islands: Stromboli, Basiluzzo, Panaria Islands

Podarcis siculus tyrrhenicus (Mertens, 1932)

Lacerta sicula tyrrhenica Mertens, 1932: 244

Type locality: Isola di Giglio, Tuscan Archipelago, Tyrrhenian Sea, Italy

Holotype: SMF 22129

Distribution: Italy: Tuscan Archipelago: Isola di Giannutri, Isola del Giglio and Formiche di Grosseto

Podarcis siculus vesseljuchi (Radovanovic, 1959)

Lacerta sicula vesseljuchi Radovanovic, 1959: 421

Type locality: Veseljeh Island, Kornati Archipelago, Croatia

Holotype: SMF 57747

Distribution: Croatia: Kornati Archipelago: Otok Veseljeh

Podarcis siculus zaratinus (Taddei, 1950)

Lacerta sicula zaratina Taddei, 1950: 152

Type localities: Zara, Scardona, Capocesto, Spatato, Croatia

Syntypes: MZUF 14763-66

Distribution: Croatia: central Dalmatian coast

Podarcis siculus zeii (Breljih, 1961)

Lacerta sicula zeii Breljih, 1961: 89

Type locality: Kal cliffs (= Sc. la Calle), West Istria, Croatia

Holotype: ML, not located

Distribution: Croatia: West Istria: Kal Island

Podarcis tauricus (Pallas, 1814)

Podarcis tauricus tauricus (Pallas, 1814)

Lacerta taurica Pallas, 1814: 30

Type locality: Crimea, Ukraine

Holotype: lost

Synonym:

Lacerta taurica thasopulae Kattainger, 1942 Type locality: Thasopoulos Island, Greece

Holotype: ?

Distribution: southern Russia, southern Ukraine, southern Moldova, eastern and southern Romania (excluding the Danube Delta), Hungary, Bulgaria, Macedonia, eastern and southern Serbia, Albania, land Greece (incl. Ionian Islands), northwestern Turkey.

Podarcis tauricus ionicus (Lehrs, 1902)

Lacerta ionica Lehrs, 1902: 232

Type locality: Kefalonia Island, Ionian Islands, Greece

Syntypes: NHMW 15721:1-9.

Distribution: Albania, western Greece, Ionian Islands

Podarcis tiliguerta (Gmelin, 1789)

Podarcis tiliguerta tiliguerta (Gmelin, 1789)

Lacerta tiliguerta Gmelin, 1789: 1070

Type locality: Sardinia, Italy

Holotype: ?

Distribution: Italy: Sardinia; France: Corsica

Podarcis tiliguerta contii Lanza & Brizzi, 1977

Podarcis tiliguerta contii Lanza & Brizzi, 1977: 158

Type locality: Isola Piana di Cavallo, southern Corsica, France

Holotype: MZUF 18913

Distribution: France: Corsica: Isola Piana di Cavallo

Podarcis tiliguerta eiselti (Lanza, 1972)

Lacerta tiliguerta eiselti Lanza, 1972: 381

Type locality: Pietricaggiosa, Piana, Maestro Maria Islands, Corsica, France

Holotype: MZUF 13804

Distribution: France: Corsica: Isola di Maestro Maria, Isola Piana, Isola Pietricaggiosa

Podarcis tiliguerta granchii Lanza & Brizzi, 1974

Podarcis tiliguerta granchii Lanza & Brizzi, 1974: 157

Type locality: Isola Poraggia Grande, Corsica, France

Holotype: MZUF 16291

Distribution: France: Corsica: Isolotto Porraggia Grande, Isolotto Porraggia Piccola

Podarcis tiliguerta grandisonae (Lanza, 1972)

Lacerta tiliguerta grandisonae Lanza, 1972: 387

Type locality: Vacca Island, southeastern Corsica, France

Holotype: MZUF 13794

Distribution: France: Corsica: Isolotto della Vacca

Podarcis tiliguerta maresi (Lanza, 1972)

Lacerta tiliguerta maresi Lanza, 1972: 389

Type locality: Toro Piccolo and Toro Grande Islands, Corsica, France

Holotype: MZUF 13758

Distribution: France: Corsica: Cerbicale Archipelago: Isolotto del Toro Piccolo, Isolotto del Toro Grande

Podarcis tiliguerta pardii Lanza & Brizzi, 1974

Podarcis tiliguerta pardii Lanza & Brizzi, 1974: 157

Type locality: Giraglia Island, northern Corsica, France
Holotype: MZUF 15861
Distribution: France: Corsica: Isola Giraglia.

Podarcis tiliguerta ranzii (Lanza, 1966)
[Lacerta tiliguerta ranzii](#) Lanza, 1966: 516
Type locality: Molarotto Island, Sardinia, Italy
Holotype: MZUF 4342
Distribution: Italy: Sardinia: Isolotto Molarotto

Podarcis tiliguerta rudolphisimonii Brizzi & Lanza, 1975
[Podarcis tiliguerta rudolphisimonii](#) Brizzi & Lanza, 1975: 61
Type locality: Isolotto di Terra, northeastern Corsica, France
Holotype: MZUF 19200
Distribution: France: Corsica: Macinaggio Islands

Podarcis tiliguerta toro (Mertens, 1932)
[Lacerta muralis toro](#) Mertens, 1932: 246
Type locality: Toro Island, southwestern Sardinia, Italy
Holotype: MSNG 31864
Distribution: Italy: Sardinia: Isolotto di Toro.

Podarcis vaucheri (Boulenger, 1905)
[Lacerta muralis vaucheri](#) Boulenger, 1905: 365
Type locality: Tangier, Morocco
Syntypes: BMNH 1946.9.1.25-30
Distribution: southern Spain, central and northern Morocco, northern Algeria, northern Tunisia.

Podarcis virescens Geniez, Sá-Sousa, Guillaume, Cluchier & Crochet, 2014
[Podarcis virescens](#) Geniez, Sá-Sousa, Guillaume, Cluchier & Crochet, 2014: 20
Type locality: 1 km past Villanueva de los Escuderos towards Cuenca by the road CUV-7037 (40.0436°N / 2.2916°W, province of Cuenca, Spain)
Holotype: MNHNP 2012.0264
Distribution: southern Spain, southern Portugal

Podarcis waglerianus Gistel, 1868
Podarcis waglerianus waglerianus Gistel, 1868
[Podarcis wagleriana](#) Gistel, 1868: 149
Type locality: Sicily
Holotype: not located
Distribution: Italy: Sicily; Favignana, Levanzo, Isola Grande dello Stagnone and Maraone Islands

Podarcis waglerianus marettimensis (Klemmer, 1956)
[Lacerta wagleriana marettimensis](#) Klemmer, 1956: 377
Type locality: Marettimo Island, Egadian Islands, west of Sicily, Italy
Holotype: SMF 50519
Distribution: Italy: Marettimo Island

Poromera Boulenger, 1887 Boulenger, G. A. 1887. *Catalogue of the Lizards in the British Museum*. III.



Lacertidae, Gerrhosauridae, Scincidae, Anelytropsidae, Dibamidae, Chamaeleontidae. London: 575pp Type species: *Tachydromus fordii* Hallowell, 1857

Poromera fordii Dan Portik

Poromera fordii (Hallowell, 1857)

Tachydromus fordii Hallowell, 1857: 48

Type locality: West Africa, Gaboon country

Holotype: ANSP 9202

Distribution: Cameroon, southwestern Central African Republic, Gabon, Equatorial Guinea (including the island of Bioko), Congo Rep., southwestern DR Congo

Psammodromus Fitzinger, 1826 Fitzinger, L. 1826. *Neue Classification der Reptilien nach ihren natürlichen*



Verwandtschaften nebst einer Verwandtschafts-Tafel und einem Verzeichnisse der Reptilien-Sammlung des K. K. Zoologischen Museums zu Wien. J. G. Heubner, Wien, 66 pp. Type species: *Psammodromus hispanicus* Fitzinger, 1826

Psammodromus algirus algirus A Miralles

Psammodromus algirus (Linnaeus, 1758)

Psammodromus algirus algirus (Linnaeus, 1758)

Lacerta algira Linnaeus, 1758: 203

Type locality: Mauritania

Holotype: not located

Synonyms:

Pantodactylus nicefori Burt & Burt, 1931 Type locality: unknown Holotype: AMNH R1082
Psammodromus manuelae Busack, Salvador & Lawson, 2006 Type locality: Manzanares el Real (40°44' N, 3° 52' W), Madrid Province, Spain Holotype: MNCN 36792
Psammodromus jeanneae Busack, Salvador & Lawson, 2006 Type locality: 25.6 km NE of Facinas (36° 08' N, 5° 42' W) along CA-221, Cádiz Province, Spain Holotype: MNCN 11941
Distribution: southern France, Spain, Portugal, Andorra, Gibraltar northern Tunisia, northern Algeria, northwestern Morocco, Italy: Conigli Islet

Psammodromus algirus doriae (Bedriaga, 1886)

Tropidosaura algira doriae Bedriaga, 1886: 410

Type locality: Galita Archipelago, Tunisia

Holotype: ?

Distribution: Tunisia: Galita Archipelago

Psammodromus algirus nollii (Fischer, 1887)

Troidosaurus algira nollii Fischer, 1887: ?

Type locality: Touggourt, Algeria

Syntype: ZMH, lost

Distribution: northern Algeria, northeastern Morocco, Tunisia

Psammodromus algirus ketamensis Galan, 1931

Psammodromus algirus ketamensis Galan, 1931: 364

Type locality: Telata de Ketama, Morocco

Holotype: MNCN 7853

Distribution: Morocco: Rif Mountains

Psammodromus blanci (Lataste, 1880)

Zerzoumia blanci Lataste, 1880:297

Type locality: Algiers, Algeria

Holotype: BMNH 1946.9.4.22

Distribution: northern Algeria, northeastern Morocco, northern Tunisia

Psammodromus edwardsianus (Dugès, 1829)

Lacerta edwardsiana Dugès, 1829: 339

Type locality: lowland Languedoc region, France

Holotype: MNHNP, not located

Distribution: eastern Spain, southern France

Psammodromus hispanicus Fitzinger, 1826

Psammodromus hispanicus Fitzinger, 1826: 22

Type locality: Perales de Tajuña, Madrid, Spain

Neotype: MNCN 41745

Distribution: central Spain

Psammodromus microdactylus (Boettger, 1881)

Algira microdactyla Boettger, 1881: 571

Type locality: near Casablanca, Morocco

Holotype: BMNH 1946.8.5.44

Distribution: Morocco: Rif, Moyen and Haut Atlas Mountains

Psammodromus occidentalis Fitz, Gonzalez-Jimena, San-Jose, San Mauro & Zardoya, 2012

Psammodromus occidentalis Fitz, Gonzalez-Jimena, San-Jose, San Mauro & Zardoya, 2012: 47

Type locality: Colmenar del Arroyo (40° 27' 22.14''N, 4° 10' 28.36''W), Madrid, Spain
Holotype: MNCN 34516
Distribution: Portugal, western and central Spain

Pseuderemias Boettger, 1883 Boettger, O. 1883. Die Reptilien und Amphibien von Marocco. II.



Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft 13: 93-146 Type species: *Acanthodactylus mucronatus* Blandford, 1870

Pseuderemias brenneri Tomáš Mazuch

Pseuderemias brenneri (Peters, 1869)

Eremias brenneri Peters, 1869: 432

Type locality: Brava, Somalia

Holotype: ZMB 6443

Synonym:

Eremias edwardsii Mocquard, 1888 Type locality: Comalis country, Somalia Holotype: ?

Distribution: Somalia, Djibouti, eastern Ethiopia

Pseuderemias erythrostickta (Boulenger, 1891)

Eremias erythrostickta Boulenger, 1891: 10

Type locality: route from Obbia to Berbera, Somalia

Syntypes: BMNH 1946.8.6.48-50

Distribution: Somalia

Pseuderemias mucronata (Blandford, 1870)

Acanthodactylus mucronatus Blandford, 1870: 453

Type locality: Anseba, Eritrea

Holotype: BMNH 1946.8.6.29

Distribution: Djibouti, southeastern Egypt, Eritrea, eastern Ethiopia, north & central Somalia, eastern Sudan

Pseuderemias savagei (Gans, Laurent & Pandit, 1965)

Eremias savagei Gans, Laurent & Pandit, 1965: 39

Type locality: Kandala, Somalia

Holotype: MCZ R-71844

Distribution: northeastern Somalia Last seen 1961

Pseuderemias septemstriata (Parker, 1942)

Eremias septemstriata Parker, 1942: 62

Type locality: Migiurtinia, Halin district, Somalia

Holotype: BMNH 1946.8.6.13

Distribution: northeastern and central Somalia

Pseuderemias smithii (Boulenger, 1895)

Eremias smithii Boulenger, 1895: 534

Type locality: Milmil (08°18'N 43°53'E), Ethiopia

Holotype: BMNH 1946.8.6.35

Distribution: northwestern and southern Somalia, eastern Ethiopia, northern Kenya

Pseuderemias striatus (Peters, 1874)

Pseuderemias striatus striatus (Peters, 1874)

Eremias brenneri striatus Peters, 1874: 370

Type locality: Brava, Somalia

Syntypes: ZMB 8120, 39143

Distribution: central and southern Somalia, eastern Ethiopia, northeastern Kenya

Pseuderemias striatus gardoensis (Arilla, Balletto & Spanò, 1965)

Eremias striatus gardoensis Arilla, Balletto & Spanò, 1965: ?

Type locality: Migiurtinia, Somalia

Holotype: ?

Distribution: northern Somalia, Djibouti, eastern Ethiopia.

Scelarcis Fitzinger, 1843

Fitzinger, L. 1843. *Systema Reptilium*, Fasciculus Primus, Amblyglossae. Vienna:

Braumüller & Seidel. 106 pp Type species: *Lacerta perspicillata* Duméril & Bibron, 1839



Scelarcis perspicillata chabanaudi A Miralles

Scelarcis perspicillata (Duméril & Bibron, 1839)

Scelarcis perspicillata perspicillata (Duméril & Bibron, 1839)

Lacerta perspicillata Duméril & Bibron, 1839: 249

Type locality: Algiers, Algeria

Holotype: MNHNP 0.2749

Distribution: coastal Algeria, Morocco: eastern High Atlas Mts

Scelarcis perspicillata chabanaudi (Werner, 1931)

Lacerta perspicillata chabanaudi Werner, 1931: 286

Type locality: Chellah of Rabat, Fes, Morocco

Syntypes: NHMW 16225, 11153

Distribution: Morocco : High and Middle Atlas, Central Plateau, Fez, and Rabat.

Scelarcis perspicillata pellegrini (Werner, 1929)

[Lacerta perspicillata pellegrini](#) Werner, 1929: 5

Type locality: Sefrou, Morocco Syntypes:

NHMW 11154, MCZ R-27390

Distribution: Morocco : High and Middle Atlas.

Takydromus Daudin, 1802 Daudin, F. M. 1802. *Histoire Naturelle, générale et particulière des reptiles,*



ouvrage faisant suite, a l'histoire naturelle, générale et particulière composée par Leclerc de Buffon, et redigée par C. S. Sonnini, vol. 3. F. Dufart, Paris. Type Species: *Takydromus sexlineatus* Daudin, 1802

Takydromus smaragdinus  Lubos Melen

Takydromus albomaculosus Wang, Gong, Liu & Wang, 2017

[Takydromus albomaculosus](#) Wang, Gong, Liu & Wang, 2017: 447

Type locality: Tianjingshan Forestry Station (24.69°N, 113.03°E), Ruyuan County, Guangdong Province, China

Holotype: SYS r001624

Distribution: Nanling Mountains, Ruyuan County, Guangdong Province, southern China

Takydromus amurensis Peters, 1881

[Tachydromus amurensis](#) Peters, 1881: 71

Type locality: Kossakewitcha, in the Amur River Valley, Russia

Holotype: ZMB 9869

Synonyms:

Takydromus kwanguensis Doi, 1920 Type locality: Mt. Kwanak, near Seoul, South Korea

Holotype: ?

Takydromus aurorealis Doi, 1929 Type locality: Seoul, South Korea Holotype: ?

Distribution: Russia, northeast China, Korea, Japan: Tsushima Island

Takydromus dorsalis Stejneger, 1904

[Takydromus dorsalis](#) Stejneger, 1904: 294

Type locality: Ishigaki shima, Yaeyama Archipelago, Ryu Kyu Islands, Japan

Holotype: USNM 34162

Distribution: Japan: Ryu Kyu Islands: Yaeyama Islands

Takydromus formosanus Boulenger, 1894

[Takydromus formosanus](#) Boulenger, 1894: 462

Type localities: Taipei and near Taiwanfoo, Taiwan

Syntypes: BMNH 1946.8.4.50, 1946.9.3.93-97, 1946.8.4.29

Distribution: Taiwan: coastal regions

Takydromus hani Chou, Truong & Pauwels, 2001

[Takydromus hani](#) Chou, Truong & Pauwels, 2001: 499

Type locality: 12 Km S Rao-AnTown (18°20' 56"N, 105°14' 47"E), Ha Tinh Province, Hung Son District, Vietnam

Holotype: NMNS 3370
Distribution: Viet Nam: Ha Tinh, Quang Nam Provinces

Takydromus haughtonianus Jerdon, 1870

Takydromus haughtonianus Jerdon, 1870: 72
Type locality: Goalpara, Assam, India
Holotype: ZSI 2195
Distribution: India: western Assam Last seen 1870

Takydromus hsuehshanensis Lin & Cheng, 1981

Takydromus hsuehshanensis Lin & Cheng, 1981: 43
Type locality: Mt. Hsiao-hsueh, Taiwan
Holotype: THUPL002
Distribution: Taiwan: Mt. Hsiao-hsueh

Takydromus intermedius Stejneger, 1924

Takydromus intermedius Stejneger, 1924: 120
Type locality: Xin-Kai-Si, Mt. Emi, near Leshan Shi, Sichuan, China
Holotype: USNM 64437
Distribution: southwest China: Yunnan, Sichuan, Guizhou, Guangxi, Hunan Provinces

Takydromus khasiensis Boulenger, 1917

Takydromus khasiensis Boulenger, 1917: 221
Type locality: Khasi Hills, Garo-Khasi Range, Meghalaya State, India
Syntypes: BMNH 1946.8.4.77-82
Distribution: Garo-Khasi Range: northeast India and Gangetic Plains, West Bengal; northern Bangladesh; northwestern Burma

Takydromus kuehnei Van Denburgh, 1909

Takydromus kuehnei kuehnei Van Denburgh, 1909
Takydromus kuehnei Van Denburgh, 1909: 50
Type locality: Kuantzuling, Taiwan
Holotype: CAS 18002
Synonyms:
Tachydromus chinensis Vogt, 1914 Type locality: northern Guangdong Prov., China
Holotype: ZMB 24096
Platyplacopus kuehnei carinatus Gressitt, 1938 Type locality: Da'an, Hainan Island, Hainan Prov., China Holotype: MVZ 23519
Distribution: Taiwan, China: Hainan, Guangdong Provinces

Takydromus kuehnei vietnamensis Ziegler & Bischoff, 1999

Takydromus kuehnei vietnamensis Ziegler & Bischoff, 1999: 209
Type locality: Ky Anh - Ke Go, Ha Tinh Province, Vietnam
Holotype: ZFMK 70342
Distribution: northern Viet Nam: Ha Tinh, Nghe An, Thanh Hoa, Ninh Binh Provinces

Takydromus lueyanus Lue & Lin, 2008

Takydromus lueyanus Lue & Lin, 2008: 390
Type locality: on the bank of Luye Stream, the branch of Beinan River Luye2 (2° 53' 47.0" N, 121° 05' 19.0" E), Taidong County, Taiwan
Holotype: NMNS 4433

Distribution: southeastern Taiwan

Takydromus madaensis Bobrov, 2013

Takydromus madaensis Bobrov, 2013: 97

Type locality: Ma Da forest, Dong Nai Province, southern Vietnam

Holotype: ZMMU R-7772

Distribution: Vietnam: Dong Nai Province Last seen 1990

Takydromus sauteri Van Denburgh, 1909

Takydromus sauteri Van Denburgh, 1909: 50

Type locality: Hengchun, Taiwan

Holotype: CAS 18001

Distribution: south and central Taiwan, including the island Lanyu

Takydromus septentrionalis Günther, 1864

Takydromus septentrionalis Günther, 1864: 69

Type locality: Ningbo Shi, Zhejiang Province, China

Holotype: BMNH 1946.8.4.27

Distribution: China: Henan, Shaansi, Gansu, Zhejiang, Sichuan, Guizhou, Hubei, Anhui, Jiangsu, Jiangxi, Hunan, Fujian, Guangdong, Guangxi, Shandong Provinces

Takydromus sexlineatus Daudin, 1802

Takydromus sexlineatus sexlineatus Daudin, 1802

Takydromus sexlineatus Daudin, 1802: 256

Type locality: Java, Indonesia

Holotype: MNHNP, not located

Synonym:

Tachydromus sexlineatus aenofuscus Peters, 1863 Type locality: Bangkok, Thailand

Holotype: ZMB 4534

Distribution: southern Burma; Malaysia: West Malaysia, East Malaysia; Indonesia: Kalimantan, Natuna Islands, Sumatra, Bangka, Java

Takydromus sexlineatus ocellatus Cuvier, 1829

Tachydromus ocellatus Cuvier, 1829: pl 11

Type locality: East Indies

Holotype: MNHNP, not located

Synonyms:

Tachydromus typus Gray, 1838 Type locality: China Syntypes: BMNH
1946.8.4.64-67

Tachydromus kwangsiensis Ahl, 1930 Type locality: Kwangsi, China Holotype: ZMB
34033

Distribution: northeastern West Malaysia; Thailand; eastern Burma; Laos; Cambodia;
Vietnam; China: Guizhou, Fujian, Hong Kong, Guangxi, Guangdong, Hainan, Yunnan
Provinces

Takydromus sikkimensis Günther, 1888

Takydromus sikkimensis Günther, 1888: 166

Type locality: Dalep village, (27°14'N, 88°28'E), nearby Singtam, south Sikkim, India

Neotype: ZSI 25761

Distribution: India: Sikkim State: Teesta Valley

Takydromus smaragdinus Boulenger, 1887

Takydromus smaragdinus Boulenger, 1887: 147

Type locality: central Ryu-Kyu Islands, Japan

Syntypes: BMNH 1946.8.4.53-63

Distribution: Japan: northern and central Ryukyu Archipelago: Okinawa, Amami and Tokara group

Takydromus stejnegeri Van Denburgh, 1912

Takydromus stejnegeri Van Denburgh, 1912: 243

Type locality: Taipei, Taiwan

Holotype: CAS 18417

Distribution: Taiwan and Pescadores Islands

Takydromus sylvaticus (Pope, 1928)

Apeltonotus sylvaticus Pope, 1928: 372

Type locality: Chogan, northwestern Fujian Province, China

Holotype: AMNH 34975

Distribution: China: Fujian Province

Takydromus tachydromoides (Schlegel, 1838)

Takydromus tachydromoides tachydromoides (Schlegel, 1838)

Lacerta tachydromoides Schlegel, 1838: 101

Type locality: Nagasaki, Japan

Syntypes: ZMB 1046, MNHNP 0.2264

Synonyms:

Tachydromus japonicus Duméril & Bibron, 1839 Type locality: China

Holotype: MNHNP 0.3262

Tachydromus holsti Boulenger, 1894 Type locality: Kyushu, Japan

Holotype: 1946.9.4.1

Distribution: Japan: Hokkaido, Honshu, Schikoku, Kyushu, Oki Islands, Yaku, Sado, Tsushima, Nakanoshima, Tokara group

Takydromus tachydromoides oldi Walley, 1958

Takydromus tachydromoides oldi Walley, 1958: 203

Type locality: from an abandoned building, Tajon, South Korea

Holotype: USNM 129618

Distribution: central South Korea

Takydromus toyamai Takeda & Ota, 1996

Takydromus toyamai Takeda & Ota, 1996: 79

Type locality: Hirara, Miyakoyima, Ryukyu Archipelago, Japan

Holotype: KUZ 18818

Distribution: Japan: southern Ryukyu Archipelago: Miyako group: Miyakojima and Irabujima

Takydromus viridipunctatus Lue & Lin, 2008

Takydromus viridipunctatus Lue & Lin, 2008: 383

Type locality: Su-ao (24°35'23.4"N, 121°51'37.5"E), Yilan County, Taiwan

Holotype: NMNS 4431

Distribution: northeastern Taiwan

Takydromus wolteri Fischer, 1885

Takydromus wolteri Fischer, 1885: 82

Type locality: Chemulpo, North Korea

Holotype: ZMH, lost

Synonyms:

Takydromus wolteri cupreus Pavlov, 1932 Type locality: Man Keou, Manchuria, China

Holotype: TNZ ?

Takydromus wolteri manchuriensis Pavlov, 1932 Type localities: Soun and Man Keou, Manchuria, China Syntypes: TNZ ?

Distribution: China: Liaoning, Jilin, Heilongjiang, Fujian, Jiangxi, Jiangsu, Anhui, Hubei, eastern Sichuan, Hunan Provinces ; Russia: southernmost Ussuri area; Korea

Teira Gray, 1838

Gray, J.E. 1838. Catalogue of the Slender-tongued Saurians, with Descriptions of many new Genera and Species. Annals and Magazine of Natural History 1: 274-283, 388-394.

Type species: *Teira punctata* Gray, 1838



Teira dugesii dugesii A&S Troidl

Teira dugesii (Milne-Edwards, 1829)

Teira dugesii dugesii (Milne-Edwards, 1829)

Lacerta dugesii Milne-Edwards, 1829: 71

Type locality: Madeira Island, Portugal

Holotype: BMNH 1946.9.3.1

Distribution: Portugal: La Gartixa de Madeira

Teira dugesii selvagensis (Bischoff, Osenegg & Mayer, 1989)

Podarcis dugesii selvagensis Bischoff, Osenegg & Mayer, 1989: 252

Type locality: Ilhas Selvagens: Selvagem Grande, Portugal

Holotype: ZFMK 41561

Distribution: Portugal: Selvagens Islands

Teira dugesii jogeri (Bischoff, Osenegg & Mayer, 1989)

Podarcis dugesii jogeri Bischoff, Osenegg & Mayer, 1989: 255

Type locality: Madeira Archipelago, Porto Santo, Portugal

Holotype: ZFMK 44193

Distribution: Portugal: Madeira Archipelago: Porto Santo Island

Teira dugesii maui (Mertens, 1938)

Lacerta dugesii maui Mertens, 1938: 287

Type locality: Deserta Grande Island, near Madeira, Portugal

Holotype: SMF 25001

Distribution: Portugal: Desertas Islands: Ilhéu Chão, Deserta Grande and Bugio

Timon Tschudi, 1836 Tschudi, J. J. 1836. Über ein neues subgenus von *Lacerta* Cuv. *Isis von Oken* 29: columns 546–551 Type species: *Lacerta lepidus* Daudin, 1802



Timon kurdistanicus R Sindaco

Timon kurdistanicus (Suchow, 1936)

Lacerta princeps kurdistanica Suchow, 1936: ?

Type locality: Beydarvaz (35°16'N, 46°09'E), Iran

Holotype: ZISP 11441:b

Distribution: southeastern Turkey, northeastern Syria, northern Iraq, northwestern Iran

Timon lepidus (Daudin, 1802)

Timon lepidus lepidus (Daudin, 1802)

Lacerta lepidus Daudin, 1802: 204

Type locality: vicinity of Montpellier, Languedoc, France

Holotype: MNHNP, not located

Synonym:

Lacerta jamaicensis Daudin, 1802 Type locality: Gibraltar Holotype: MNHNP ?

Distribution: Spain (except the southeast); Andorra; Portugal; southern France; northwestern Italy

Timon lepidus ibericus (López-Seoane, 1884)

Lacerta ocellata iberica López-Seoane, 1884: 19

Type locality: Spain

Lectotype: SMF 6038

Distribution: northwestern Spain, northwestern Portugal

Timon lepidus oterorum (Castroviejo & Mateo, 1998)

Lacerta lepidus oteroi Castroviejo & Mateo, 1998: 7

Type locality: Isla de Salvora, outskirts of Santa Eugenia de Ribeira, prov. de La Coruña, Spain

Holotypes: EBD 16729

Distribution: Spain: Sálvora island off the mouth of the Arousa river

Timon nevadensis (Buchholz, 1963)

Lacerta lepidus nevadensis Buchholz, 1963: 155

Type locality: Sierra Nevada, southern Spain, north slope of des Pico Veleto

Holotype: ZFMK 3581
Distribution: southeastern Spain

Timon pater (Lataste, 1880)

[Lacerta ocellata pater Lataste, 1880: 306](#)

Type localities: Lambaesa, Setif, El Guerah, Edough, Algiers and Batna, Algeria

Syntypes: BMNH 1946.9.3.86-87, 1946.8.6.55, 1946.9.2.19-26, MCZ R-4613

Distribution: Algeria, Tunisia

Timon princeps (Blanford, 1874)

[Lacerta princeps Blanford, 1874: 31](#)

Type locality: southern Iran

Holotype: ZSI 3351

Distribution: Iran: central Zagros Mountains

Timon tangitanus (Boulenger, 1889)

[Lacerta ocellata tangitana Boulenger, 1889: 303](#)

Type locality: Tangiers, Morocco

Syntypes: BMNH 1946.9.2.11-16, 1946.8.6.4-5,

Distribution: Morocco

Tropidosaura Fitzinger, 1826 Fitzinger, L. 1826. Neue Classification der Reptilien nach ihren natürlichen



Verwandtschaften nebst einer Verwandtschafts-Tafel und einem Verzeichnisse der Reptilien-Sammlung des K. K. Zoologischen Museums zu Wien. J. G. Heubner, Wien, 66 pp. Type species: *Lacerta montanus* Gray, 1831

Tropidosaura cottrelli Hewitt

Tropidosaura cottrelli (Hewitt, 1925)

[Basutosaura cottrelli Hewitt, 1925: 356](#)

Type locality: Nemahedi Camp, nr. Mont aux Sources, Basutoland, South Africa

Holotype: PEM R16093

Distribution: South Africa: Drakensberg: Free State, KwaZuluNatal Province; Lesotho

Tropidosaura essexi Hewitt, 1927

[Tropidosaura essexi Hewitt, 1927: 378](#)

Type locality: Mont aux Sources, Basutoland, South Africa

Syntypes: BMNH 1946.8.4.48, 1946.9.4.20; PEM R16094–98

Distribution: South Africa: Drakensberg: Lesotho; South Africa: southeastern Free State, southwestern KwaZuluNatal Provinces

Tropidosaura gularis Hewitt, 1927

Tropidosaura gularis Hewitt, 1927: 387

Type locality: Table Mountain, Cape Province, South Africa

Holotype: SAM 17723

Distribution: South Africa: southern Western Cape, southwestern Eastern Cape Provinces

Tropidosaura montana (Gray, 1831)

Tropidosaura montana montana (Gray, 1831)

Lacerta montanus Gray, 1831: 35

Type locality: Cape of Good Hope, South Africa

Syntypes: BMNH 1946.8.4.46., 1946.8.4.49

Distribution: South Africa: Western Cape, southwestern Eastern Cape Provinces

Tropidosaura montana natalensis Fitzsimons, 1947

Tropidosaura montana natalensis Fitzsimons, 1947: ?

Type locality: ?

Holotype: TM 21281

Distribution: South Africa: southern KwaZuleNatal Province

Tropidosaura montana rangeri Hewitt, 1926

Tropidosaura montana rangeri Hewitt, 1926: 485

Type locality: Farm Gleniffer, near Kei Rd, southeastern Cape Province

Syntype: BMNH 1946.8.4.47

Distribution: South Africa: Eastern Cape Province

Vhembelacerta Edwards, Herrel, Vanhooydonck, Measey, Tolley & Branch 2013. Edwards, S.,



W.R. Branch, B. Van Hooydonck, A. Herrel, G.J. Measey, K.A. Tolley. 2013. Taxonomic adjustments in the systematics of the southern African lacertid lizards (Sauria: Lacertidae). *Zootaxa* 3669 (2): 101–114 Type species: *Lacerta rupicola* FitzSimons, 1933

Vhembelacerta rupicola Sebastian Kirchof

Vhembelacerta rupicola (Fitzsimons, 1933)

Lacerta rupicola Fitzsimons, 1933: 278

Type locality: Lake Fundusi, Zoutpansberg Mtns., Limpopo Province, South Africa

Holotype: TMP 13989

Distribution: South Africa: Soutpansberg, Limpopo Province

Zootoca Wagler, 1830

Wagler, J. 1830. Natürliches System der Amphibien, mit vorangehender



Classification der Säugthiere und Vögel. Ein Beitrag zur vergleichenden Zoologie. Munich, Stuttgart and Tübingen: J.G. Cotta. vi + 354 pp. + one plate. Type species: *Lacerta vivipara* Lichtenstein, 1823

Zootoca vivipara vivipara ♂ Gebhart

Zootoca vivipara (De Jacquin, 1787)

Zootoca vivipara vivipara (De Jacquin, 1787)

Lacerta vivipara Jacquin, 1787: 33

Type locality: Mount Scheeberg, near Vienna, Austria

Holotype: Icontype: Jacquin 1787: Tab II

Synonyms:

Lacertus cinereus Lacépède, 1788 Type locality: ? Holotype: ?

Lacerta oedura Sheppard, 1804 Type locality: England Holotype: ?

Lacerta montana Mikan, 1805 Type locality: Riesengebirge, Germany Holotype: Icontype: Sturni, 1805: 112, fig A

Lacerta nigra Wolf, 1805 Type locality: Wenger Alpe, Canton Bern, Switzerland. Holotype: Icontype: Sturni, 1805: 112, fig B

Lacerta crocea Wolf, 1805 Type locality: vicinity of Nuremberg, Germany Holotype: Icontype: ? Sturni, 1805

Lacerta unicolor Kuhl, 1820 Type locality: unknown Holotype: ?

Lacerta ptychodes Kuhl, 1820 Type locality: ? Holotype: ?

Lacerta pyrrhogaster Merrem, 1820 Type locality: Germany Holotype: ?

Lacerta schreibersiana Milne-Edwards, 1829 Type locality: Vienna basin, Austria Holotype: MNHNP0.51

Lacerta schreibersiana fusca Gachet, 1832 Type locality: vicinity of Bordeaux, France Holotype: ?

Lacerta schreibersiana lutea Gachet, 1832 Type locality: vicinity of Bordeaux, France Holotype: ?

Lacerta chrysogastra Andrzejowski, 1832 Type locality: valleys around Cremeniec, Ukraine Holotype: ?

Zootoca guerini Cocteau, 1835 Type locality: Foret d'Eu, Departement Seine-Maritime, Upper Normandy, France Holotype: MNHNP 0.47

Lacerta vivipara pallida Fatio, 1872 Type locality: Swiss Alps and in the Jura Mountains, Switzerland Holotype: ?

Lacerta vivipara melanogastra Prazák, 1898 Type locality: Bohemia, Czech Republic Holotype: ?

Lacerta vivipara barabensis Kashenko, 1902 Type locality: ? Holotype: ?

Distribution: Albania, western Austria, Belarus, Belgium, Bosnia and Herzegovina, Bulgaria, China, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Germany, Ireland, Italy, Japan, Kosovo, Latvia, Liechtenstein, Lithuania, Luxembourg, Macedonia, Moldova, Mongolia, Montenegro, Netherlands, Norway, northern Poland, Russia, Serbia, Sweden, Switzerland, Ukraine, United Kingdom, Greece

Zootoca vivipara louislantzi Arribas, 2009

Zootoca vivipara louislantzi Arribas, 2009: 140

Type locality: Pla de Beret, Vall d'Aràn, Lleida, Spain

Holotype: MNCN 44290

Distribution: Spain: Pyrenees Mountains, Basque Mountains, Cantabrian Mts., Xistral Mountains; France: Landes Department

Zootoca vivipara pannonica (Lac & Kluch, 1968)

Lacerta vivipara pannonica Lac & Kluch, 1968: 8

Type locality: Boľany, Kapu-sansky Forest in eastern Slovakia

Holotype: ?

Distribution: eastern Austria, Slovakia, Hungary, Romania

Zootoca vivipara sachalinensis (Pereleshin & Terentjev, 1963)

Lacerta vivipara sachalinensis Pereleshin & Terentjev, 1963: 4

Type locality: Sakhalin Island, Russia

Holotype: ?

Distribution: Poland, southern Belarus: Carpathian Mountains; southeastern Russia, incl. Sakhalin island; Japan: northern Hokkaido Island

Zootoca vivipara carniolica (Mayer, Böhme, Tiedemann & Bishoff, 2000)

Lacerta vivipara carniolica Mayer, Böhme, Tiedemann & Bishoff, 2000: 65

Type locality: Mt. Sneznik: 8 km SE Masun village, Slovenia.

Holotype: ZFMK 68438

Distribution: Slovenia, northeastern Italy, southern Austria, northwestern Croatia

ACKNOWLEDGEMENTS

This work could not have been completed without the help of numerous individuals. We thus acknowledge the help and advice of Werner Conradie (PEM), Annamaria Nistri (MZUF), Patrick Campell (BMNH), Boris Kryštufek (ML) and others

The author expresses his to the photographers who have contributed to this work

If any of the readers have any of the missing data (designated by ?) [there is a lot in this catalog], I would appreciate if they would contact me

(HDJTNP.gambelia@gmail.com)

REFERENCES

- Agarwal, I., A. Khandekar, U. Ramakrishnan, R. Vyas and V. B. Giri. 2018.** Two New Species of the *Ophisops microlepis* (Squamata: Lacertidae) Complex from northwestern India with A Key to Indian Ophisops. *Journal of Natural History*, 52: 1-29
- Ahl, E. 1930.** Beiträge zur Lurch- und Kriechtierfauna Kwangsi Eidechse . *Sitzungsberichte der Gesellschaft naturforschender Freunde zu Berlin*, 1930: 310-332
- Ahmadzadeh, F, M. Flecks, M.A. Carretero, O. Mozaffari, W. Böhme, D.J. Harris, S. Freitas & D.Rödder. 2013.** Cryptic Speciation Patterns in Iranian Rock Lizards Uncovered by Integrative Taxonomy. *PLoS ONE* 8(12): e80563. doi:10.1371/journal.pone.0080563
- Alcock, A. W. & F Finn. 1897.** An account of the Reptilia collected by Dr. F.P. Maynard, Captain A.H. McMahon, C.I.E., and the members of the Afghan-Baluch Boundary Commission of 1896. *Journal of the Asiatic Society of Bengal* 65: 550-566
- Anderson, S.C. & A.E. Leviton. 1967.** A new species of *Eremias* (Reptilia: Lacertidae) from Afghanistan. *Occasional Papers of the California Academy of Sciences* (64): 1-4
- Andrzejowski, A. 1832.** Reptilia inprimis Volhyniae. Podoliae, et Gubernii Chersonensis. *Nouveaux Mémoires de la Société Impériale des Naturalistes de Moscou*, 2 : 320-346 Taf. XXII-XXIV
- Angel, F. 1936.** Reptiles et batraciens de Syrie et de Mésopotamie récoltés (part.) M.P. Pallary. *Bulletin De l'Institut d'Egypte* 18: 107-116
- Angel, M.F. 1936.** Description d'une nouvelle espèce du genre *Philochortus* provenant des confins Saharo-Soudanais, et tableau synoptique des espèces du genre. *Bulletin de la Société zoologique de France* (Paris) 61: 100-105
- Arillo, A., E. Balletto & S. Spanò. 1965.** II e III spedizione Scortecci in Migiurtinia: il genere *Eremias* Wiegmaan (Reptilia, Lacertidae). *Bollettino dei Musei e degli Istituti Biologici della Università di Genova*, vol. 33, p. 85-109.
- Arnold, E.N. 1972.** Lizards with northern affinities from the mountains of Oman. *Zoologische Mededelingen* (Leiden) 47: 111-128
- Arnold, E.N. 1980.** The scientific results of the Oman flora and fauna survey 1977 (Dhofar). The reptiles and amphibians of Dhofar, southern Arabia. *Journal of Oman Studies*, Special Report (No. 2): 273-332
- Arnold E N 1986.** A new spiny-footed lizard (*Acanthodactylus*: Lacertidae) from Saudi Arabia. *Fauna of Saudi Arabia* 8: 378-384
- Arnold, E.N. 1989.** Towards a phylogeny and biogeography of the Lacertidae: relationships within an Old-World family of lizards derived from morphology. *Bulletin of the British Museum (Natural History), Zoology*, 55, 209-257.
- Arnold, E.N., O. Arribas & S. Carranza. 2007.** Systematics of the Palearctic and Oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with descriptions of eight new genera. *Zootaxa* 1430: 1-86
- Arribas, O. J. 1993.** Estatus específico para *Lacerta* (*Archaeolacerta*) *monticola bonnali* Lantz, 1927 (Reptilia, Lacertidae). *Boletin de la Real Sociedad Espanola de Historia Natural* (Sec. Biol.) 90 (1-4): 101-112
- Arribas, O. 1994.** Una nueva especie de Lagartija de los Pirineos Orientales: *Lacerta* (*Archaeolacerta*) *aurelioi* sp. nov. (Reptilia: Lacertidae). *Bollettino del Museo regionale di Scienze naturali*, Torino 12 (1): 327-351

- Arribas, O.J. 1996.** Taxonomic revision of the Iberian 'Archaeolacertae' I: A new interpretation of the geographical variation of '*Lacerta monticola* Boulenger 1905 and '*Lacerta cyreni* Müller & Hellmich 1937 (Squamata: Sauria: Lacertidae). *Herpetozoa* 9: 31-56
- Arribas, O. J. 1997.** Morfologia, filogenia y bibliografía de las lagartijas de alta montaña de los Pirineos . Tesis doctoral - Universitat Autònoma de Barcelona. *Publicaciones de la Universitat Autònoma de Barcelona*, 08193 Bellaterra (Barcelona), 353 pp
- Arribas, O.J. & S. Carranza & G. Odierna. 2006.** Description of a new endemic species of mountain lizard from Northwestern Spain: *Iberolacerta galani* sp. nov. (Squamata: Lacertidae). *Zootaxa*, 1240: 1-55.
- Arribas, O.J. 2009.** Morphological variability of the Cantabro-Pyrenean populations of *Zootoca vivipara* (Jacquin, 1787) with description of a new subspecies. *Herpetozoa* 21 (3/4): 123-146
- Arribas, O.J. , C. Ilgar, Y. Kumlutas, S.H. Durmus, A. Avci & N. Üzüüm. 2013.** External morphology and osteology of *Darevskia rudis* (Bedriaga, 1886), with a taxonomic revision of the Pontic and Small-Caucasus populations (Squamata: Lacertidae). *Zootaxa*, 3626 (54): 401-428.
- Arribas, O.J., P. Galan, N. Remon & H. Naveira. 2014.** A new mountain lizard from Montes de León (NW Iberian Peninsula): *Iberolacerta monticola astur* ssp. nov. (Squamata: Lacertidae) *Zootaxa* 3796 (2): 201–236
- Arribas, O., Ç. Ilgaz, Y. Kumlutaş. 2018.** Reevaluation of the intraspecific variability in *Darevskia parvula* (Lantz & Cyren, 1913): an integrated approach using morphology, osteology and genetics (Squamata: Lacertidae). *Zootaxa* 4472(1): 71–99
- Arillo, A., E. Balletto & S. Spanò. 1965.** II e III Spedizione Scotecchi in Migiurtinia: il genere *Eremias* Wiegmann (Reptilia, Lacertidae). *Bollettino dei Musei e degli Istituti Biologici Dell'Università di Genova*, 33 (204): 85-109.
- Arillo, A., E. Balletto & S. Spano. 1967.** Il genere *Latastia* Bedriaga in Somalia. *Bollettino dei Musei e degli Istituti Biologici dell'Università di Genova*, 35: 105-145
- Audouin, J. V. 1827.** Explication sommaire des planches de reptiles (supplément) ... offrant un exposé des caractères naturels des genres, avec la distinction des espèces. . In: Savigny, M. J. C. L. de., *Description d'Égypte*, Vol. I. Historie Naturelle. Paris: Imprimerie impériale/Imprimerie royale
- Baha El Din, S.M. 2007.** A new lizard of the *Acanthodactylus scutellatus* group (Squamata: Lacertidae) from Egypt. *Zoology in the Middle East* 40: 21-32
- Baig, K.J. and R. Masroor 2006.** A New Species of *Eremias* (Sauria: Lacertidae) from Cholistan Desert, Pakistan. *Russian Journal of Herpetology* 13 (3): 167-174
- Bakradze, M.A. 1976.** New subspecies, *Lacerta portschinskii nigrita* ssp.n., from the Eastern Transcaucasia.- *Vestnik zoologii, Journal of the Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine*, Kiev, 1976 (4): 54-57.
- Baloutch, M. 1976.** Une nouvelle espèce de *Lacerta* (Lacertilia, Lacertidae) de sud-est de l'Iran. *Bulletin du Muséum National d'histoire naturelle, Paris*, 417: 1379-1384
- Baran, I. & A. Budak. 1978.** A new form of *Ophisops elegans* (Lacertidae, Rept.) from Anatolia. *Ege Üniversitesi Fen Fakültesi Dergisi, Bornova-Izmir (Ser.B), C II (S. 2): 185-192.*
- Baran, I., Y. Kumlutas, B. Lanza, R. Sindaco, Ç. Ilgaz & A. Avci & P. Crucitti. 2005.** *Acanthodactylus harranensis*, A New Species of Lizard from Southeastern Turkey (Reptilia: Sauria: Lacertidae). *Bollettino del Museo Regionale di Scienze Naturali Torino* 23 (1): 323-341
- Barbour, T. 1914.** Some new reptiles. *Proceedings of the New England Zoölogical Club* 4: 95-98

- Barrbadillo, L.J., J.I. Lacomba, V. Pérez-Mellado, V. Sancho & L.F. López-Jurado. 1999.** *Anfibios y Reptiles de la Peninsula Ibérica, Baleares y Canarias*. Barcelona (Editorial Planeta), 419 pp.
- Bartenjew, D. & Rjesnikowa. 1931.** Neue Lacerta-Formen (Reptilia) aus dem Kaukasischen Staatsnaturschutzgebiete (West-Kaukasus). *Zoologischer Anzeiger* 96: 9-10.
- Basoglu, M. & W. Hellmich. 1968.** Eine neue Eremias-Form aus Ost-Anatolien (Reptilia, Lacertidae). *Ege Üniversitesi Fen Fakültesi İlmî Raportlar Serisi, Bornova-İzmir* 67: 1-9
- Bauer, A.M., J.L. Childers, C. Broeckhoven and P.F.N. Mouton. 2019.** A New *Nucras* Gray, 1838 (Squamata: Lacertidae) from the Strandveld of the Western Cape, South Africa. *Zootaxa*. 4560(1); 149–163
- Beddome, R. H. 1870.** Descriptions of new reptiles from the Madras Presidency. *Madras Monthly Journal of Medical Science*, 2: 169-176
- Bedriaga, J. v. 1876.** *Die Faraglione-Eidechse und die Entstehung der Farben bei den Eidechsen*. Heidelberg: Carl Winter: 1-21
- Bedriaga, J. von 1879.** Herpetologische Studien. *Archiv für Naturgeschichte* XLIV: 259-320
- Bedriaga, J. von 1879.** Memoire sur les variétés européennes du lézard des murailles. *Bulletin de la Société zoologique de France*, Paris, 4: 194-228
- Bedriaga, J. v. 1881.** Über die Auffassung und Anwendung der Begriffe: Species, Subspecies und Varietas. *Zoologischer Anzeiger* 4: 66-71
- Bedriaga, J. von. 1882.** Die Amphibien und Reptilien Griechenlands. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 56:43—103
- Bedriaga, J. von 1884.** Die neue Lacertiden-Gattung *Latastia* und ihre Arten. *Annali del Museo Civico di Storia Naturale Giacomo Doria* 20: 307-324
- Bedriaga, J. von 1886.** Beiträge zur Kenntnis der Lacertiden-Familie (*Lacerta*, *Algiroides*, *Tropidosaura*, *Zerzumia*, *Bettaia*). *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* (Frankfurt) 14: 17-444
- Bedriaga, J. von 1907[1906].** Verzeichnis der von der Central-Asiatischen Expedition unter Stabs-Kapitän W. Roborowski in den Jahren 1893-1895 gesammelten Reptilien. *Annuaire Musée Zoologique de l'Académie Impériale des Sciences de St.-Pétersbourg* , 10 (3-4): 159-200
- Bedriaga, J. von 1912.** *Amphibien und Reptilien*. Wissenschaftliche Resultate der von N.M. Przewalski nach Central-Asien unternommenen Reisen. *Zoologischer Theil*, Bd. 3, Abth. 1, Lfrg. 4, S. 503 – 769.
- Berger-Dell'mour, H. A.E. & W. Mayer. 1989.** On the parapatric existence of two species of the *Pedioplanis undata* group (Reptilia: Sauria: Lacertidae) in the central Namib Desert (Southwest Africa) with description of the new species *Pedioplanis husabensis*. *Herpetozoa* 1 (3-4): 83-95
- Berthold, A.A. 1840.** Ueber verschiedene neue oder seltene Amphibienarten. *Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen*, 1: 47-72
- Bird, C. G. 1936.** The distribution of reptiles and amphibians in Asiatic Turkey with notes on a collection from the Vilayets of Adana, Gaziantep and Malatya. *Annals and Magazine of Natural History* (10) 18: 257-281.
- Bibron, M. & J.B. Bory de Saint-Vincent. 1833.** Vertébrés a sang froid. Reptiles et poissons. Reptiles. In: Geoffroy & Geoffroy, *Expédition Scientifique de Morée*, Tome III. 1re partie, Zoologie: pp. 57-76

- Bischoff, W. 1982.** Zur Kenntnis der innerartlichen Gliederung der Artwiner Eidechse *Lacerta derjugini*. *Zoologische Abhandlungen Staatliches Museum für Tierkunde in Dresden* 1 (38): 1-52
- Bischoff, W. 1982.** Die innerartliche Gliederung von *Gallotia galloti* (Duméril & Bibron 1839) (Reptilia: Sauria: Lacertidae) auf Teneriffa, Kanarische Inseln. *Bonner zoologische Beiträge*, 33 (2-4): 363-382.
- Bischoff, W. 1984.** Bemerkungen zur innerartlichen Gliederung und zur Verbreitung der Artwiner Eidechse (*Lacerta derjugini* Nikol'sky, 1898) an den Südhängen des Grossen Kaukasus (Sauria: Lacertidae). *Salamandra* 20 (2/3): 101-111
- Bischoff, W. 1985.** Vorläufige Beschreibung der Eidechse der Kanareninsel Fuerteventura als *Gallotia atlantica mahoratae* ssp. nov. *Salamandra* 21 (1): 46-54
- Bischoff, W., K. Osenegg and W. Mayer. 1989.** Untersuchungen zur subspezifischen Gliederung der Madeira-Mauereidechse, *Podarcis dugesii* (Milne-Edwards, 1829). *Salamandra* 25 (3/4): 237-259
- Bischoff, W. & J. Müller. 1999.** Revision des levantinischen *Lacerta laevis/kulzeri*-Komplexes: 2. Die Petra-Eidechse *Lacerta kulzeri petraea* ssp.n. *Salamandra* 35 (4): 243-254
- Blanford, W.T. 1870.** *Observations of the geology and zoology of Abyssinia, made 1867-68.* McMillan (London), xii + 487 pp
- Blanford, W.T. 1874.** Descriptions of new Reptilia and Amphibia from Persia and Baluchistan. *Annals and Magazine of Natural History* (4) 14: 31-35
- Blanford, W.T. 1875.** List of Reptilia and Amphibia collected by the late Dr. Stoliczka in Kashmir, Ladák, eastern Turkestan, and Wakhán, with descriptions of new species. *Journal of the Asiatic Society of Bengal*, Calcutta, new ser., 44 (pt. 2, no. 3): 191-196
- Blanford, W. T. 1881.** Notes on the lizards collected in Socotra by Professor I. Bayley Balfour. *Proceedings of the Zoological Society of London* 1881: 464-469
- Blyth, E. 1854.** Notices and descriptions of various reptiles, new or little-known. Part I. *Journal of the Asiatic Society of Bengal*, 22 [1853]: 639-655.
- Bobrov, V.V. 2013.** A new species of Grass Lizard (Reptilia, Sauria, Lacertidae, *Takydromus*) from Southern Vietnam. *Current Studies in Herpetology*, 13 (3/4): 97-100.
- Bocage, J.V. du B. 1866.** Lista dos reptis das possessões portuguezas d'Africa occidental que existem no Museu Lisboa. *Jornal de Sciencias Mathematicas, Physicas e Naturaes, Academia Real das Sciencias de Lisboa* 1: 37-56
- Bocage, J.V. du B. 1867.** Descriptions of two new Saurians from Mossamedes (West Africa). *Annals and Magazine of Natural History* (3) 20: 225-228
- Bocage, J.V. du B. 1867.** Segunda lista dos reptis das possessões portuguezas d'Africa occidental que existem no Museu de Lisboa. *Jornal de Sciencias Mathematicas, Physicas e Naturaes, Academia Real das Sciencias de Lisboa* (3): 217-228
- Bodenheimer, F.S. 1944.** Introduction into the knowledge of the Amphibia and Reptilia of Turkey. *Revue de la Faculté des Sciences de l'Universite d'Istanbul*, Ser. B. 9: 1-78
- Böhme, W. & A. Budak. 1977.** Über die rudis-Gruppe des *Lacerta saxicola*-Komplexes in der Türkei, II (Reptilia: Sauria: Lacertidae). *Salamandra* 13 (3/4): 141-149

- Bohme, W. & N.N. Szczerbak. 1991.** Ein neuer Wustenrenner aus dem Hochland Afghanistans, *Eremias (Eremias) afghanistanica* sp. n. (Reptilia: Sauria: Lacertidae). *Bonner Zoologische Beiträge* 42(2): 137-41
- Bonaparte, C.L. 1838.** *Iconografia della Fauna Italica* II – Anfibi. Salvineci & Filippo, Roma 1832–41.
- Bons, J. 1960.** Description d'un nouveau lézard du Sahara: *Eremias pasteuiri* sp. nov. (Lacertidés). *Comptes rendus des séances mensuelles, Société des Sciences Naturelles et Physiques du Maroc*, 4: 69-71
- Börner, A-R. 1974.** Resultate der Rhodos-Exkursion, April 1974. Teil 1. Eine neue Unterart der Eidechse *Lacerta danfordi* (Günther) 1876 von den südlichen Sporaden. *Miscellaneous Articles in Saurology* 3: 1-11
- Boscá, E. 1916.** Un género nuevo para la fauna herpetológica de España y especie nueva o poco conocida. *Boletín de la Sociedad Española de Historia Natural* 16: 294-297
- Boettger, O. 1878.** Reptilien und Amphibien aus Syrien. *Bericht über die Senckenbergische Naturforschende Gesellschaft in Frankfurt am Main*, vol. 1878: 57-84
- Boettger, O. 1880.** Die Reptilien und Amphibien von Syrien, Palaestina und Cypem. *Bericht über die Senckenbergische Naturforschende Gesellschaft in Frankfurt am Main*, 1879-1880: 132-219
- Boettger, O. 1881.** Diagnoses Reptilium novorum Maroccanorum. *Zoologischer Anzeiger* 4: 570-572
- Boettger, O. 1883.** Die Reptilien und Amphibien von Marocco. II. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* 13: 93-146
- Boettger, O. 1888.** Aufzählung einiger neu erworbener Reptilien und Batrachier aus Ost-Asien. *Bericht der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt* 1888: 187-190
- Boettger, O. 1892.** Wissenschaftliche Ergebnisse der Reise Dr. Jean Valentins im Sommer 1890 – I. Kriechtiere der Kaukasusländer, gesammelt durch die Radde-Valentin'sche Expedition nach dem Karabagh und durch die Herren Dr. J. Valentin und P. Reibisch. *Berichte über die Senckenbergische naturforschende Gesellschaft in Frankfurt am Main* 1892: 131-150.
- Boettger, O. 1893.** Übersicht der von Prof. C. Keller anlässlich der Ruspoli'schen Expedition nach den Somaliländern gesammelten Reptilien und Batrachier. *Zoologischer Anzeiger* 16 (416): 113-119
- Boettger, O. & L. Müller. 1914.** Preliminary notes on the local races of some Canarian lizards. *Annals and Magazine of Natural History* (8) 14: 67-78
- Bolkay, St. J. 1919.** Prinosi herpetologiji zapadnoga dijela balkanskog poluostrva [Additions to the herpetology of the western Balkan peninsula]. *Glasnik zemaljskog Muzeja u Bosni i Hercegovini*, Sarajevo, 31, 1-38.
- Bons, J. & B. Girot. 1964.** Révision de l'espèce *Acanthodactylus scutellatus* (Lacertidé-Saurien). *Bulletin de la Société des Sciences Naturelles du Maroc* 42: 311-334
- Boscá, E. 1883.** Exploración herpetológica de la isla de Ibiza. *Anales de la Sociedad Española de Historia Natural* Madrid 12: 241-250
- Boscá, E. 1916.** Dos observaciones a propósito de la *Lacerta muralis* en España. *Boletín de la Real Sociedad Española de Historia Natural* 16: 327-330

- Boulenger, G.A. 1878.** Sur les espèces d'*Acanthodactylus* des bords de la Méditerranée. *Bulletin de la Société zoologique de France* 3: 179-197
- Boulenger, G.A. 1887.** Descriptions of new reptiles and batrachians in the British Museum (Natural History), Part iii. *Annals and Magazine of Natural History* (5) 20: 50-53
- Boulenger, G. A. 1887.** *Catalogue of the Lizards in the British Museum* (Nat. Hist.) III. Lacertidae, Gerrhosauridae, Scincidae, Anelytropsidae, Dibamidae, Chamaeleontidae. London: 575pp
- Boulenger, G.A. 1887.** On a collection of reptiles and batrachians made by Mr. H. Pryer in the Loo Choo Islands. *Proceedings of the Zoological Society of London* 1887: 146-150
- Boulenger, G. A. 1889.** On the reptiles and batrachians obtained in Morocco by M. Henry Vaucher. *Annals and Magazine of Natural History* (6) 3: 303-307
- Boulenger, G.A. 1891.** On some Reptiles collected by Sig. L. Bricchetti Robecchi in Somaliland. *Annali del Museo Civico de Storia Naturale di Genova* (2) 12: 5-15
- Boulenger, G.A. 1894.** Description of a new lizard and a new fish obtained in Formosa by Mr. Holst. *Annals and Magazine of Natural History* (6) 14: 462-463
- Boulenger, G.A. 1894.** Second report on additions to the lizard collection in the Natural History Museum. *Proceedings of the Zoological Society of London* 1894: 722-736
- Boulenger, G.A. 1895.** An account of the reptiles and batrachians collected by Dr. A. Donaldson Smith in western Somaliland and the Galla Country. *Proceedings of the Zoological Society of London* 1895: 530-540.
- Boulenger, G. A. 1897.** Description of a new lizard from Obok. *Annals and Magazine of Natural History* (6) 19: 467-468
- Boulenger, G. A. 1898.** On a second collection of reptiles made by Mr. E. Lord-Phillips in Somaliland. *Annals and Magazine of Natural History* (7) 2: 130-133
- Boulenger, G. A. 1899.** Description of two new lizards from the interior of British East Africa. *Proceedings of The Zoological Society of London* 1899: 96-98
- Boulenger, G.A. 1902.** A List of the Fishes, Batrachians, and Reptiles collected by Mr. J. ffolliott Darling in Mashonaland, with Descriptions of new Species. *Proceedings of the Zoological Society of London* , 1902: 13-18
- Boulenger, G.A. 1903.** Descriptions of two new Lizards discovered by Mr. E. Degen in his Journey to Abyssinia. - *The Annals and magazine of natural history*, 7 (11): 54-55
- Boulenger, G. A. 1905.** A contribution to the knowledge of the varieties of the Wall-Lizard (*Lacerta muralis*) in western Europe and North Africa. *Transactions of the Zoological Society of London*, 17: 351-420
- Boulenger, G.A. 1905.** On a new variety of the Wall Lizard (*Lacerta muralis* var. *breviceps*). *Annuario del Museo Zoologico della Università di Napoli* 1 (29): 1-2.
- Boulenger, G.A. 1906.** Additions to the herpetology of British East Africa. *Proceedings of The Zoological Society of London* 2: 570-573

- Boulenger, G.A. 1907.** Descriptions of two new African lizards of the genus *Latastia*. *Annals and Magazine of Natural History* (7) 19: 392-394
- Boulenger, G.A. 1907.** Descriptions of new lizards in the British Museum. *Annals and Magazine of Natural History* (7) 19: 486-489
- Boulenger, G.A. 1908.** Description of a new species of *Lacerta* from Persia. *Proceedings of The Zoological Society of London* 1907: 934-936
- Boulenger, G.A. 1909.** Description of a new lizard of the genus *Acanthodactylus* from Syria. *Annals and Magazine of Natural History* (8) 4: 188-189
- Boulenger, G.A. 1913.** Second contribution to our knowledge of the varieties of the wall-lizard (*Lacerta muralis*). *Proceedings of The Zoological Society of London* 1913: 135-230
- Boulenger, G.A. 1916.** Description of a new genus of the family Lacertidae from Central Africa. *Annals and Magazine of Natural History* (8) 18: 112-114
- Boulenger, G.A. 1917.** Descriptions of new lizards of the family Lacertidae. *Annals and Magazine of Natural History* (8) 19: 277-279
- Boulenger, G.A. 1917.** Description of a new South African lizard of the genus *Eremias*. *Annals of the South African Museum* 13 (4): 217-219
- Boulenger, G.A. 1917.** A revision of the lizards of the genus *Tachydromus*. *Memoirs of the Asiatic Society of Bengal, Calcutta.*, 5 (6): 207-235
- Boulenger, G.A. 1918.** Sur les lézards du genre *Acanthodactylus* Wieg. *Bulletin de la Société zoologique de France* 43: 143-155
- Boulenger, G.A. 1918.** Description of a new lizard of the genus *Acanthodactylus* from Mesopotamia. *Journal of the Bombay Natural History Society* 25: 373-374
- Boulenger, G.A. 1918.** On the varieties of the lizard *Ophisops elegans* Ménétries. *Annals and Magazine of Natural History* (9) 2: 158-161
- Boulenger, G.A. 1919.** On a new variety of *Acanthodactylus boskianus* Daud. from the Euphrates. *Annals and Magazine of Natural History* (9) 3: 549-550
- Boulenger, G.A. 1919.** On a new lizard of the genus *Latastia* from southern Rhodesia. *Transactions of the Royal Society of South Africa* 8 (1): 39-40
- Boulenger, G.A. 1921.** Monograph of the Lacertidae. Vol. II. *British Museum (Natural History). Department of Zoology*. London. 451
- Braun, M. 1877.** *Lacerta lilfordi* und *Lacerta muralis*. *Arbeiten aus dem Zoologisch-Zootomisch Institut zu Würzburg* 4: 1-64
- Brelih, S. 1961.** Seven new races of the species *Lacerta (Podarcis) sicula* Rafinesque from the Rovinj-Porec Region. *Biološki Vestnik* 9: 71-91
- Brelih, S. 1963.** Prispevek k poznavanju kvarnerskih kuščaric (Reptilia, Lacertidae). *Biološki Vestnik* 11: 107-113

- Brizzi, R. & B. Lanza. 1975.** The natural history of the Macinaggio Islets (northeastern Corsica) with particular reference to their herpetofauna. *Natura - Societa italiana di scienze naturali e del Museo civico di storia naturale di Milano* 66 (1-2): 53-72
- Broadley, D.G. 1967.** A new species of *Ichnotropis* (Sauria: Lacertidae) from the Botswana-Caprivi border. *Arnoldia* 3 (24): 1-5
- Broadley, D.G. 1972.** A review of the *Nucras tessellata* group (Sauria: Lacertidae). *Arnoldia* 20: 1-35
- Buchholz, K.F. 1954.** Zur Kenntnis der Rassen von *Lacerta pityusensis* Bosca (Rept., Lacertidae). *Bonner zoologische Beiträge* 5 (1/2): 69-88
- Buchholz, K. F. 1960.** Zur Kenntnis von *Lacerta peloponnesiaca* (Reptilia: Lacertidae). *Bonner zoologische Beiträge* 11: 87-107
- Buchholz, K.F. 1962.** Die Mauereidechse von Ananes, Milos-Archipel. *Bonner zoologische Beiträge*, 13 (1/3): 216-218
- Buchholz, K.F. 1963.** Die Perleidechse der Sierra Nevada (Reptilia: Lacertidae). *Bonner zoologische Beiträge*, 14 (1/2): 151-156.
- Buchholz, K.F. 1965.** Zur Kenntnis des Genus *Algyroides* (Reptilia: Lacertidae). *Bonner zoologische Beiträge* 15: 239-246
- Burt, C.E. & M.D. Burt. 1931.** South American lizards in the collection of the American Museum of Natural History. *Bulletin of the American Museum of Natural History*. 61 (7): 227-395
- Busack, S.D., A. Salvador & R. Lawson. 2006.** Two new species in the genus *Psammotromus* (Reptilia: Lacertidae) from the Iberian Peninsula. *Annals of Carnegie Museum*, 75 (1): 1-10.
- Camerano, L. 1877.** Considerazioni sul genere *Lacerta* Linn. e descrizione di due nuove specie. *Atti della Accademia delle scienze di Torino* 13: 79-98
- Camerano, L. 1885.** Monografia dei Sauri italiani. *Zoologischer Anzeiger* 8: 417-419
- Capolongo, D. 1979.** Nota preliminare su di una nuova forma microinsulare di *Podarcis sicula* (Rafinesque) dell'isola di Dino nel mar Tirreno. *Bollettino della Società dei naturalisti in Napoli*, 87, 387-392
- Capolongo, D. 1984.** Note sull'erpetofauna Pugliese. *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, 125: 189-200
- Cara, A. 1872.** *Monografia della Lucertola comune di Sardegna*. Timon (Cagliari). 41 pp
- Castroviejo, J. & J.A. Mateo & E. Collado. 1985.** Sobre la sistemática de *Gallotia atlantica* (Peters y Doria, 1882). - *Publicacion Ocasional, Doñana, Acta Vertebrata*, Sevilla, 1985: 1-85.
- Castroviejo, J. & J.A. Mateo. 1998.** Una nueva subespecie de *Lacerta lepida* Daudin, 1802 (Sauria, Lacertidae) para la Isla de Salvora (España). *Publicaciones Asociación Amigos de Doñana*, 12: 1-21.
- Chabanaud, P. 1917.** Énumération des reptiles non encore étudiés de l'Afrique occidentale, appartenant aux collections du Muséum, avec la description des espèces nouvelles. *Bulletin du Muséum National d'Histoire Naturelle*. Paris 23: 83-105

Chabanaud, P. 1918. Étude complémentaire de deux Agama de l'Afrique occidentale et description de quatre espèces nouvelles de reptiles de la même région. *Bulletin du Museum National d'Histoire Naturelle*. Paris 24: 104-112

Chabanaud, P. 1918. Etude d'une collection de reptiles de l'Afrique occidentale française. *Bulletin du Museum National d'Histoire Naturelle*. Paris 24: 160-166

Chabanaud, P.M. 1919. Énumération des Reptiles et des Batraciens de la péninsule Balkanique envoyés au Muséum par le Dr. Rivet, de 1917 à 1919, avec la description d'une variété nouvelle. *Bulletin du Muséum d'Histoire Naturelle, Paris*. vol. 25, no 1, p. 21-26

Chou, W-H., N.Q. Truong and O.S. G. Pauwels. 2001. A new species of *Takydromus* (Reptilia: Lacertidae) from Vietnam. *Herpetologica* 57 (4):497-508

Cirer, A.M. 1980. Descriptió de dues subespecies noves de *Podarcis pityusensis*. *Butlletí de la Institució Catalana d'Historia Natural*, 45: 121-126

Cocteau, T. 1835. Sur un genre peu connu de Lézards vivipares (*Zootoca*, Wagler), et sur une nouvelle espèce de ce genre. *Magasin de Zoologie*, 1835. 25 pp. Taf. 9.

Conradie, W., G.J. Measey, W.R. Branch & K.A. Tolley. 2012. Revised phylogeny of African sand lizards (*Pedioplanis*), with the description of two new species from south-western Angola. *African Journal of Herpetology* 61 (2): 91-112

Cope, E.D. 1862. On *Lacerta echinata* and *Tiliqua dura*. *Proceedings of the Academy of Natural Sciences of Philadelphia* 1862: 189-191

Costa, O.G. 1828. Fauna di Aspromonte e sue adiacenze. *Atti della Reale Accademia delle Scienze, sezione della Società Reale Borbonica*, 4: 61-84.

Costa, O.G., 1839. Fauna di Aspromonte e sue adiacenze. Letta nella tornata de' 12 Febrajo 182_8. *Atti dell'Accademia delle scienze fisiche e matematiche di Napoli*, 4: 61-173, 4 tav. f.t.

Cuvier, G. 1829. *Le règne animal distribué d'après son organisation, pour servir de base à l'Histoire naturelle des Animaux et d'introduction à l'Anatomie Comparée*. Nouvelle Edition [second edition]. Vol. 2. Les Reptiles. Déterville, Paris, i-xvi, 1-406

Cyrén, O. 1933. Lacertiden der Südöstlichen Balkanhalbinsel. *Izvestiia na Tsarskitia prirodonauchni instituti v Sofiia*, 6: 219-240

Cyren. O. 1941. Beiträge zur Herpetologie der Baikanhalbinsel. *Mitteilungen aus den Königlichen Naturwissenschaftlichen Instituten in Sofia* 14, 36-152.

Daudin, F. M. 1802. *Histoire Naturelle, générale et particulière des reptiles*, ouvrage faisant suite, à l'histoire naturelle, générale et particulière composée par Lelerc De Buffon, et redigée par C. S. Sonnini, vol. 3. F. Dufart, Paris.

Darevsky, I.S. 1953. About regular position of Transcaucasian multi-coloured *Eremias arguta* (Pallas) (Reptilia, Sauria). *Report of the Armenian CCP Academy of Sciences* 16(4): 117 - 121

Darevsky, I.S. 1957. Systematics and ecology of rock lizards (*Lacerta saxicola* Eversmann) in Armenia. *Zoology sb. AN Armenia*, Yerevan 10: 27-57

- Darevsky, I.S. 1966.** Natural parthogenesis in a polymorphic group of Caucasian Rock lizards related to *Lacerta saxicola* Eversmann. *Journal of the Ohio Herpetological Society* 5 (4): 115-152
- Darevsky, I. S. 1967.** Rock lizards of the Caucasus: Systematics, ecology and phylogenesis of the polymorphic groups of Caucasian rock lizards of the the subgenus *Archaeolacerta* (Skal'nye Yashcheritsy Kavkasa; *Izdat elst'vo Nauka, Leningrad*). Leningrad, Nauka (English translation 1978): Ind. Nat. Sci. Doc. Centre; New Delhi, 276 pp
- Darevsky, I.S. & J. Eiselt. 1967.** Ein neuer Name für *Lacerta saxicola mehelyi*, Lantz und Cyrén, 1936. *Annalen des Naturhistorischen Museums in Wien*, 70: 107
- Darevsky, I. & V. Vedmederja. 1977.** A new species of rock lizard *Lacerta saxicola* Eversmann group from northeastern Turkey and adjoining regions of Adjaria. *Trudy Zoologicheskogo Instituta Akademii Nauk SSSR*, 74: 50-54
- Darevsky, I.S. and F.D. Danielyan. 1977.** *Lacerta uzzelli* sp. nov. (Sauria: Lacertidae) - a new parthenogenetic species of rock lizard from eastern Turkey. *Trudy Zoologicheskogo Instituta Akademii Nauk SSSR*, Leningrad 76: 55-59
- Darevsky, I. S. and J. Eiselt. 1980.** Neue Felseneidechsen (Reptilia: Lacertidae) aus dem Kaukasus und aus der Türkei. *Amphibia-Reptilia* 1: 29-40
- Darevsky, I.S. & N.N. Shcherbak. 1978.** *Eremias andersoni*, a new lizard (Reptilia, Lacertilia, Lacertidae) from Iran. *Journal of Herpetology* 12 (1): 13-15
- Darevsky, I.S. & B.S. Tuniyev. 1997.** A new species from *Lacerta saxicola* group - *Lacerta dryada* sp. nov. (Sauria: Lacertidae) and some comments relative to *Lacerta clarkorum* Darevsky & Vedmederja 1977. *Russian Journal of Herpetology* 4 (1): 1-7
- Darevsky, I. S. & E. S. Roitberg 1999.** A new subspecies of the rock lizard *Lacerta caucasica* (Sauria, Lacertidae) from the south-east of Chechen Republic on the Caucasus. *Russian Journal of Herpetology* 6 (3): 209-214
- De Betta, E. 1857.** Erpetologica delle Provincie Venete e del Tirole meridionale. *Accademia di agricoltura, arti e commercio di Verona*. 18; 383 pp
- De Filippi, F. 1863.** Nuove o poco note specie di animali vertebrati raccolte in un viaggio in Persia. *Archivio per la Zoologia, l'Anatomia e la Fisiologia* (Genova) 2: 377-394
- De Filippi, F. 1865.** *Note di viaggio in Persia nel 1862*. Milan, 1, 1-396
- De Jacquin, J.F. 1787.** *Lacerta vivipara*, observatio. *Nova Acta Helvetica*. I: 33-34
- Deraniyagala, P.E.P. 1953.** A Colored Atlas of some vertebrates from Ceylon, Tetrapod Reptilia Vol. II. *National Museums of Sri Lanka*: 101
- Deraniyagala, P. 1971.** A new lizard from Ceylon. *Spolia Zeylanica*, 32: 103-105
- De Witte, G.-F. 1942.** Description d'un Lacertidae nouveau du Congo Belge. *Bulletin Musee royal d'Histoire naturelle de Belgique* XVIII (41), 4pp
- Di Palma, M. 1980.** La Lucertola del faraglione "La Canna" nelle Isole Eolie: *Podarcis sicula cucchiarai* subsp. nova (Reptilia, Lacertidae). *Il Naturalista Siciliano*, Ser. IV, 4: 3-12.
- Doi, H. 1920.** *Takydromus kwangakuensis* sp. nov. from Korea. *Chosen Iho*, 1920: 70-75.

Doi, H. 1929. A new species of *Takydromus* from Corea, *Takydromus auroralis* sp. nov. *Journal of the Chosen Natural History Society*, 9: 17-19

Doronin, I.V. 2011. A description of a new subspecies of Rock Lizard *Darevskia brauneri myusserica* ssp. nov. from the Western Transcaucasia (Abkhazia), with comments on systematics of *Darevskia saxicola* complex. *Proceedings of the Zoological Institute*, 315 (3): 242-262

Doumergue, F. 1901. *Essai sur la faune erpétologique de l'Oranie* avec des tableaux analytiques et des notations pour la détermination de tous les reptiles et batraciens du Maroc, de l'Algerie et de la Tunisie. L. Fouque, Oran, 404 pp

Dugès, A. 1829. Mémoire sur les espèces indigènes du genre *Lacerta*. *Annales des Sciences naturelles*, 16: 337-491

Duméril, A.M.C. and G. Bibron. 1839. *Erpétologie Générale ou Histoire Naturelle Complète des Reptiles*. Vol.5. Roret/Fain et Thunot, Paris, 871 pp

Dvigubsky, I.A. 1832. ?

Eichwald, E. 1831. *Zoologia specialis*, quam expositis animalibus tum vivis, tum fossilibus potissimum rossiae in univ. et poloniae in specie, in usum lectionum publicarum in Universitate Caesarea Vिल्nensi. Vol. 3 Josophi Zawadski 428pp

Eimer, G.H.T. 1872. Über eine neue Eidechse von Capri. *Verhandlungen der Physikalisch-medicinischen gesellschaft zu Würzburg* 3: 9-11

Eimer, T. 1874. Zoologische Studien auf Capri. II. *Lacerta muralis coerulea*, ein Beitrag zur Darwinschen Lehre. - Leipzig

Eimer, T. 1881. *Untersuchungen über das Variieren der Mauereidechse*. Berlin, R. Stricker, 281 pp. + 3 plates

Eiselt, J. 1979. Ergebnisse zoologischer Sammelreisen in der Türkei *Lacerta cappadocica* Werner 1902 (Lacertidae, Reptilia). *Annalen des Naturhistorischen Museums in Wien* 82: 387-422

Eiselt, J. & J.F. Schmidtler. 1986. Der *Lacerta danfordi*-Komplex (Reptilia: Lacertidae). *Spixiana*, München, 9 (3): 289-328.

Eiselt, J. & I. Darevsky. 1991. *Lacerta rudis chechenica* ssp.n. aus dem Kaukasus (Reptilia: Lacertidae). *Annalen des Naturhistorischen Museums in Wien* 92: 15-29

Eiselt, J., I.S. Darevsky, J.F. Schmidtler. 1992. Untersuchungen an Felseidechsen (*Lacerta saxicola* - Komplex; Reptilia: Lacertidae) in der östlichen Türkei. 1. *Lacerta valentini* Boettger. *Annalen des Naturhistorischen Museums in Wien*, Serie B, 93: 1-18

Eiselt, J., J.F. Schmidtler. & I.S. Darevsky. 1993. Untersuchungen an Felseidechsen (*Lacerta saxicola* Komplex) in der östlichen Türkei. 2. Eine neue Unterart der *Lacerta raddei* Boettger, 1892 (Squamata: Sauria: Lacertidae). *Herpetozoa* 6 (1/2): 65-70

Eiselt, J. 1995. Ein Beitrag zur Kenntnis der Archaeolacerten (sensu Mehely, 1909) des Iran (Squamata: Sauria: Lacertidae). *Herpetozoa* 8 (1-2): 59-72

Eisentraut, M. 1928. Vorläufige Diagnosen einiger neuer Rassen der balearischen Inseleidechse *Lacerta lilfordi* Gthr. *Das Aquarium* 1928: 121-124

Eisentraut, M. 1928. Weitere neue Rassen der balearischen Inseleidechse *Lacerta lilfordi* Gthr. *Mitteilungen aus dem Zoologischen Museum in Berlin*, 14: 465-468

Eisentraut, M. 1930. Beitrag zur Eidechsenfauna der Pityusen und Columbreten. *Mitteilungen aus dem Zoologischen Museum in Berlin*, 16: 397-410

Eremchenko, V., A.M. Panfilov & E.I. Tsarinenko. 1992. *Conspectus on cytogenetic and taxonomic research on some Asiatic species of Scincidae and Lacertidae.* Ylym Publ., Bishket, Kiev, 182 pp.

Eremchenko, V. & A. Panfilov. 1999. Taxonomic position and geographic relations of a lacertid lizard *Eremias velox* from the Issyk-Kul lake depression, Tien Shan mountains, Kyrgyzstan. *Science and New Technologies* (1): 119-125

Eversmann, E. 1834. Lacertae Imperii Rossici variis in itineribus meis observatae. *Nouveaux mémoires de la Société impériale des naturalistes de Moscou* 3: 339-358.

Fatio, V. 1872. **Faune des Vertébrés de la Suisse:** Histoire naturelle des reptiles et des batraciens, J.-B. Baillièere et fils 1872, Geneva

Fejervary-Langh, A.M. 1924. Preliminary notes to a monograph of the lacertian fauna of the Maltese Island. *Biologica Hungarica*, Budapest, 1 (5): 1-14

Ferreira, J.B. 1892. Revisão dos réptis e batrachios de Portugal. *Jornal de Ciências, Mathemáticas, Physicas e Naturaes.* Lisboa 3: 19-27

Fischer, J. G. 1885. Ichthyologische und herpetologische Bemerkungen. V. Herpetologische Bemerkungen. *Jahrbuch der Hamburgischen Wissenschaftlichen Anstaltent.* 2: 82-121

Fischer, J. G. 1886. Herpetologische Notizen. *Abhandlungen des Naturwissenschaftlichen Vereins in Hamburg* 9: 1-19

Fischer, J. von 1887. Über die Kieleidechsen (*Zerzunia Blanci* Lataste und *Tropidosaura algira* Linné) und eine Varietät der letzteren, *Trop. Alg. Vor. Nollii.* v. *F Zoologischer Garten* , 28 (3): 65-74

Fitze, P.S., V. Gonzalez-Jimena, L.M. San-José, D. San Mauro & R. Zardoya. 2012. A new species of sand racer, *Psammmodromus* (Squamata: Lacertidae), from the Western Iberian Peninsula. *Zootaxa* 3205: 41–52

FitzSimons, V. 1933. Description of five new lizards from the Transvaal and Southern Rhodesia. *Annals of the Transvaal Museum.* 15 (2): 273-280.

Fitzsimons, V.F.M. 1947. Descriptions of new species and subspecies of reptiles and amphibians from Natal, together with notes on some other little known species. *Annals of the Natal Museum* (Pietermaritzburg) 11 (1): 111-137

Fitzinger, L. 1826. Neue Classification der Reptilien nach ihren natürlichen Verwandtschaften. *Nebst einer Verwandtschafts-Tafel und einem Verzeichnisse der Reptilien-Sammlung des K. K. Zoologischen Museums zu Wien.* J. G. Heubner, Wien, 66 pp

Fitzinger, L. 1843. *Systema Reptilium*, Fasciculus Primus, Amblyglossae. Vienna: Braumüller & Seidel. 106 pp

Fuhn, I.E. and R. Mertens. 1959. Studien an *Lacerta trilineata* aus Rumänien mit Beschreibung einer neuen Unterart. *Senckenbergiana Biologica* 40 (1/2): 25-42

- Fuhn, I.E., & S. Vancea. 1964.** Die innerartliche Gliederung der Zauneidechse (*Lacerta agilis*) in Rumänien. *Senckenbergiana Biologica* 45 (3/4): 469-489
- Gachet, H. 1832.** Notice sur le lézard de Schreibers (*Lacerta Schreibersiana* Milne-Edwards, *annal des sc. Nat.*, t. 16, 1829, P. 68, 83, pl. 5, fig. 5). *Actes de la Société Linnéenne de Bordeaux*, 5: 233-242.
- Galán, F. 1931.** Batracios y reptiles del Marruecos Español. *Boletín de la Real Sociedad Española de Historia Natural*, 31: 361-367
- Gans C., R.F. Laurent & H. Pandit. 1965.** Notes on a herpetological collection from the Somali Republic. *Annales du Musée Royal de l'Afrique Centrale* (8vo), Sciences Zoologiques, Tervuren 134: 1-93
- Geniez, P. & A. Foucart 1995.** Un nouvel acanthodactyle en Algérie: *Acanthodactylus taghitensis* n. sp. (Reptilia, Sauria, Lacertidae). *Bulletin du Muséum National d'Histoire Naturelle* (Sec. A Zool. Biol. Ecol. Anim.) 17 (1-2): 3-9.
- Geniez, P. & P. Sá-Sousa, C.P. Guillaume, A. Cluchier & P-A. 2014.** Systematics of the *Podarcis hispanicus* complex (Sauria, Lacertidae) III: valid nomina of the western and central Iberian forms. - *Zootaxa*, 3794 (1): 1–51
- Gistel, J. 1868.** Die Lurche Europas. Ein Beitrag zur Lehre von der geographischen Verbreitung derselben. In: *Blicke in das Leben der Natur und des Menschen*. Leipzig: Ed Wartig, 274 pp
- Gmelin, J.F. 1789.** Caroli a Linné *Systema naturae*. 13. ed., Tom 1 Pars 3. G. E. Beer, Lipsiae. 1033-1516 pp.
- Gray, J. E. 1831.** A synopsis of the species of Class Reptilia. In: Griffith, E & E. Pidgeon: *The animal kingdom arranged in conformity with its organisation by the Baron Cuvier* with additional descriptions of all the species hitherto named, and of many before noticed. V Whittaker, Treacher and Co., London: 481 + 110 pp
- Gray, J. E. 1838.** Catalogue of the slender-tongued saurians, with descriptions of many new genera and species. Part 1. *Annals and Magazine of Natural History* (1) 1: 274-283
- Gray, J.E. 1845.** *Catalogue of the specimens of lizards in the collection of the British Museum*, London. London: Trustees of the British Museum. (Edward Newman, printer). xxviii + 289 pp. pp. 36.
- Gray, J.E. 1864.** Description of two new genera of lizards (*Holaspis* and *Poriodogaster*, A. Smith, MS). *Annals and Magazine of Natural History* (3) 13: 101-104
- Gray, J. E. 1864.** Notes on some new lizards from south—eastern Africa, with the descriptions of several new species. *Proceedings of The Zoological Society of London* 1864: 58—62.
- Greenbaum, E., C.O. Villanueva, C. Kusamba, M.M. Aristote and W.R. Branch. 2011.** A molecular phylogeny of Equatorial African Lacertidae, with the description of a new genus and species from eastern Democratic Republic of the Congo. *Zoological Journal of the Linnean Society* 163: 913–942
- Greenbaum, E. & Dowell Beer, S. & Hughes, D.F. & Wagner, P. & Anderson, C.G. & Villanueva, C.O. & Malonza, P.K. & Kusamba, C. & Muninga, W.M. & Aristote, M.M. & Branch, W.R. 2018.** Phylogeography of Jackson's Forest Lizard *Adolfus jacksoni* (Sauria: Lacertidae) Reveals Cryptic Diversity in the Highlands of East Africa. *Herpetological Monographs*, 32:51–68
- Gressitt, J. L. 1938.** A new burrowing frog and a new lizard from Hainan Island. *Proceedings of the Biological Society of Washington* 51: 127-130.
- Gruber, U. 1986.** *Podarcis gageae* - Skyros-Mauereidechse. In: Böhme, W. (ed.), *Handbuch der Reptilien und Amphibien Europas*, Band 2/II., Echsen III (Podarcis). Aula-Verlag Wiesbaden, pp. 65-70
- Guillaume, C.P. & P. Geniez. 1986.** Description d'une sous-espèce de *Podarcis hispanica* (Sauria: Lacertidae): *Podarcis hispanica cebennensis* Guillaume et Geniez in Fretey. *Bulletin de la Société Herpétologique de France*, 39: 1-15

- Gulia, G. 1914.** *Lacerta generalensis*. In: Despott, G. 1914 : The Reptiles on the Maltese Islands. *The Zoologist* (London), vol. 891, no 19,p. 321-327
- Günther, A. 1859.** On the reptiles and fishes collected by the Rev. H.B. Tristram in northern Africa. *Proceedings of The Zoological Society of London* 1859: 469-470
- Günther, A. 1864.** *The Reptiles of British India*. London (Taylor & Francis), xxvii + 452 pp
- Günther, A. 1864.** Report on a collection of reptiles and fishes from Palestine. *Proceedings of The Zoological Society of London* 1864: 488-493
- Günther, A. 1872.** Description of three new species of *Eremias*. *Annals and Magazine of Natural History* (4) 9: 381-382
- Günther, A. 1872.** On some new species of reptiles and fishes collected by J. Brenchley, Esq. *Annals and Magazine of Natural History* (4) 10 (60): 418-426.
- Günther, A. 1874.** Description of a new European species of *Zootoca*. *Annals and Magazine of Natural History* (4) 14: 158-159
- Günther, A. 1876.** Description of a new species of lizard from Asia Minor. *Proceedings of The Zoological Society of London* 1876: 818
- Günther, A. 1888.** On a collection of reptiles from China. *Annals and Magazine of Natural History* (6) 1: 165-172
- Günther, A. 1903.** Reptiles from Rio de Oro, Western Sahara. *Novitates Zoologicae*. 10: 298-299
- Haas, G. 1951.** Remarks on the status of the lizard *Eremias olivieri* in Palestine, and a description of a new subspecies. *Copeia* 1951 (4): 274-276
- Haas, G. 1957.** Some amphibians and reptiles from Arabia. *Proceedings of the California Academy of Natural Sciences* (3): 47-86
- Haas, G. and Y.L.Werner. 1969.** Lizards and snakes from Southwestern Asia, collected by Henry Field. *Bulletin of the Museum of Comparative Zoology*, Harvard 138: 327-406
- Hallowell, E. 1857.** Notes of a collection of reptiles from the Gaboon country, West Africa, recently presented to the Academy of Natural Sciences of Philadelphia, by Dr. Henry A. Ford. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 9: 48-72.
- Hartmann, M. 1953.** Die Rassenaufspaltung der Balearischen Inseleidechsen. *Zoologische Jahrbücher. Abteilung für Systematik, Geographie und Biologie der Tiere*, Jena., 64: 86-96
- Hecht, G. 1930.** Die märkische Smaragdeidechse, *Lacerta viridis* Laur. subsp. *Brandenburgiensis* subsp. nov. *Das Aquarium* 1930: 62-63
- Heidari, N., N. Rastegar-Pouyani, E. Rastegar-Pouyani & M. Rajabizadeh. 2013.** A new species of *Acanthodactylus* Fitzinger 1834 (Sauria: Lacertidae) from southern Iran. *Zootaxa* 3722 (3): 333-346
- Hewitt, J. 1925.** On some new species of Reptiles and Amphibians from South Africa. *Records of the Albany Museum* (Grahamstown) 3: 343-370

Hewitt, J. 1926. Some new or little-known reptiles and batrachians from South Africa. *Annals of the South African Museum* 20: 473-490

Hewitt, J. 1927. Further descriptions of reptiles and batrachians from South Africa. *Records of the Albany Museum* (Grahamstown) 3 (5): 371-415

Hewitt, J. 1935. Some new forms of batrachians and reptiles from South Africa. *Records of the Albany Museum*, 1935: 283-357

Hutterer, R. 1985. Neue Funde von Rieseneidechsen (Lacertidae) auf der Insel Gomera. *Bonner zoologische Beiträge*, Bonn . 36: 365-394

Jerdon, T.C. 1870. Notes on Indian Herpetology. *Proceedings of the Asiatic Society of Bengal* March 1870: 66-85

Joger, U. & W. Mayer. 2002. A new species of *Mesalina* (Reptilia: Lacertidae) from Abd al-Kuri, Socotra Archipelago, Yemen, and a preliminary molecular phylogeny for the genus *Mesalina*. *Fauna of Arabia* 19: 497-505

Karaman, M. S. 1928. Contribution a l'herpetologie de la Jugoslavija. *Bulletin de la Société Scientifique de Skoplje* 4 (1): 129-143

Kashchenko, N.F. 1902. Obzor gadov Tomskago kraya. *Tomsk (MN Kononov)* 1-24

Kattinger, E. 1942. Makedonische Reptilien. IV. Die Taurische Eidechse. *Blätter für Aquarien- und Terrarienkunde*, 39: 59-60

Kessler, K.F. 1878. Transcaucasian voyage. *Gesellschaft Naturforscher St. Petersburg* 8: 143-163

Keymar, P.F. 1986. Die Amphibien und Reptilien der ionischen Region (Griechenland). analyse ihrer rezenten Verbreitungsmuster und Überlegungen zu ihrer Ausbreitungsgeschichte. *ÖHG-Nachrichten* (Wien) 8/9: 8-44

Kirchhof, S., J. Penner, M.-O. Rödel, J. Müller. 2017. Resolution of the types, diagnostic features, and distribution of two easily confused Sand Lizards, *Pedioplanis laticeps* (Smith, 1845) and *P. burchelli* (Duméril & Bibron, 1839) (Squamata: Lacertidae). *Zootaxa* 4318(1): 82-109

Klemmer, K. 1956. Eine neue Eidechsenrasse von Marettimo (Ägäische Inseln). *Senckenbergiana biologica*, Frankfurt/Main, 37: 377-380

Klemmer, K. 1959. Systematische Stellung und Rassengliederung der Spanischen Mauereidechse, *Lacerta hispanica*. *Senckenbergiana biologica*, Frankfurt/Main, 40 (5/6): 245-250.

Knoepffler, P. & E. Sochurek. 1956. Amphibien und Reptilien zwischen Banyuls und Mentone. *Aquarien und Terrarium* 3: 147-151

Kolombatovic, J. 1886. Imenik kraljesnjaka Dalmacije. II. dio: Dvoživci, gmazovi i ribe. *God. izv. Velike realke u Splitu* 1885/86 (Split): 1-20

Kuhl, H. 1820. *Beiträge zur Zoologie und vergleichenden Anatomie*. Hermannsche Buchhandlung, Frankfurt, 152 pp.

Lác, J. & E. Kluch. 1968. Die Bergeidechse der Ostslowakischen Tiefebene als selbständige Unterart *Lacerta vivipara pannonica* n. subsp. *Zoologické listy*, Brno, czech republic, 17 (2): 157-173

- Lantz, L.A. & O. Cyrén. 1913.** Eine neue Varietät der Felseneidechse *Lacerta saxicola* Eversmann *parvula* nov. var. *Tiflis Mitt. Kaukasus Museum Mus.* 7: 163-168
- Lantz, L.A. & O. Cyrén. 1919.** On *Lacerta praticola*, Eversm. *Annals & Magazine of natural History* (9) 3: 28-31
- Lantz, L.A. & O. Cyrén. 1920.** Notes sur les *Lacerta viridis* du Caucase. *Bulletin de la Société zoologique de France* 45: 33-37
- Lantz, L.A. 1927.** Quelques observations nouvelles sur l'herpétologie des Pyrénées Centrales. *Extrait de la Revue d'Histoire naturelle appliquée* Première partie, No. E et 2: 1-14. (13-11)
- Lantz, L.A. 1928.** Les *Eremias* de l'Asia Occidentale. *Bulletin du Musée de Géorgie*, Tbilisi vols. 4/5: 136 pp
- Lantz, L.A. & G.F. Suchow. 1934.** *Apathya cappadocica urmiana* susp. nov., eine neue Eidechsenform aus dem persischen Kurdistan. *Zoologischer Anzeiger* 106: 294.
- Lantz, L.A. & O. Cyrén. 1936.** Contribution à la connaissance de *Lacerta saxicola* Eversmann. *Bulletin de la Société zoologique de France* (Paris) 61: 159-181
- Lanza, B. 1952.** Note critiche su alcune lucertole italiane e diagnosi preliminare di una nuova razza insulare. *Natura*, Milano 43 (3-4): 69-82
- Lanza, B. 1954.** Notizie su alcune lucertole italiane e descrizione di una nuova razza insulare del golfo di Salerno. *Bolletino di zoologia* (Torino) 21: 133-143
- Lanza, B. 1956.** Contributo alla migliore conoscenza di alcune forme italiane di *Lacerta muralis* (Laurenti) e descrizione di una nuova razza dell'Arcipelago Toscano. *Monitore Zoologico Italiano* 63: 259-284.
- Lanza, B. 1958.** Notizie su due popolazioni insulari di *Lacerta muralis* e sulla *Vipera ursinii* in Italia. *Annali del Museo Civico di Storia Naturale « G. Doria »*, Genova 70: 305-322
- Lanza, B. 1967.** Su due nuove razze insulari di *Lacerta sicula* e di *Lacerta tiliguerta*. *Archivio Zoologico Italiano*, Napoli 51: 511-522 [1966]
- Lanza, B., B. Adriani & M. Romiti. 1971.** Eine neue blaue Inselrasse der *Lacerta sicula* aus Süditalien. *Salamandra* 7: 5-8.
- Lanza, B. & D. Capolongo. 1972.** Die blaue Ruineneidechse der tyrrhenischen Insel Licosa (Salerno). *Salamandra* 8 (1): 21-26.
- Lanza, B. 1972.** The natural history of the Cerbicale Islands (southeastern Corsica) with particular reference to their herpetofauna. *Natura - Società italiana di scienze naturali e del Museo civico di storia naturale di Milano* 63 (4): 345-407
- Lanza, B. & R. Brizzi. 1974.** On two new Corsican microinsular subspecies of *Podarcis tiliguerta* (Gmelin, 1789). *Natura - Società italiana di scienze naturali e del Museo civico di storia naturale di Milano*, 65 (3/4): 155-193
- Lanza, B. & M. Poggesi. 1975.** On a new *Eremias* from central Somalia. *Monitore Zoologico Italiano*, Firenze (n.s.) Suppl. 6: 305-312
- Lanza, B. 1976.** On a new Wall-lizard from Corsica, with notes on the Sanguinarie Islands (Reptilia Lacertidae). *Natura - Società italiana di scienze naturali e del Museo civico di storia naturale di Milano* 67 (3-4): 185-202

- Lataste, F. 1880.** Diagnoses de reptiles nouveaux d'Algérie. *Le Naturaliste* 1. *Zerzoumia blanci* n.sp.; 2. *Pterodactylus*; 3. *Lacerta ocellata pater* n.subsp : 99-325
- Lataste, F. 1881.** Diagnoses de reptiles nouveaux d'Algerie. *Le Naturaliste* 1881: 357-359
- Lataste, F. 1885.** Les Acanthodactyles de Barbarie et les autres espèces du genre. Description d'une nouvelle espèce, du pays des Çomalis (*Acanthodactylus Vaillanti*). *Annali del Museo Civico di Storia Naturale di Genova*, ser. 2A, vol. 2, p. 476–516
- Laurent, R.F. 1952.** Batraciens et reptiles récemment acquis par la Musée du Congo belge. *Revue de Zoologie et de Botanique Africaines*, Bruxelles, 45 (3-4): 198-203
- Laurent, R.F. 1964.** Reptiles et batraciens de l'Angola (troisième note). *Companhia de Di amantes de Angola (Diamang), Serviços Culturais, Museu do Dundo (Angola)*, No. 67, 165 pp.
- Laurenti, J.N. 1768.** *Specimen medicum, exhibens synopsis reptilium emendatam cum experimentis circa venena et antidota reptilium Austriacorum.* pp. [1-2], 1-214, [1], Tab. I-V [= 1-5]. Viennæ. (Trattner).
- Lacepède, B.G.E. 1788.** *Histoire naturelle des quadrupèdes ovipares et des serpens.* Vol. I . 18+651 pp. Paris
- Lehrs, P. 1902.** Zur Kenntnis der Gattung Lacerta und einer verkannten Form: *Lacerta ionica*. *Zoologischer Anzeiger* 25: 225-237
- Lehrs, P. 1910.** Ueber eine *Lacerta* aus dem hohen Libanon (*L. fraasi* n. sp.) und andere Montanformen unter den Eidechsen. In: *Festschrift für R. Hertwig*, Bd. 2. Jena, G. Fischer, pp. 225-238
- Lehrs, P. 1914.** Description of a new lizard from the Canary Islands. *Proceedings of the Zoological Society of London*, 1914: 681-684.
- Leviton, A.E. & S.C. Anderson. 1967.** Survey of the reptiles of the Sheikdom of Abu Dhabi, Arabian Peninsula. Part II. Systematic account of the collection of reptiles made in the Sheikdom of Abu Daby by John Gasperetti. *Proceedings of the California Academy of Natural Sciences* (4) 39: 157-192
- Lichtenstein, M.H.C. 1823.** Verzeichniss der Doubletten des zoologischen Museums der Königl. Universität zu Berlin nebst Beschreibung vieler bisher unbekannter Arten von Säugethieren, Vögeln, Amphibien und Fischen. *Königlichen Akademie der Wissenschaften zu Berlin*. T. Trautwein, Berlin. x, 118 pages
- Lichtenstein, M.H. 1856.** Nomenclator reptilium et amphibiorum musei zoologici Berolinensis. *Königlichen Akademie der Wissenschaften*, Berlin. 48 pp.
- Lin, J.Y. & H.Y. Cheng. 1981.** A new species of *Takydromus* (Sauria: Lacertidae) from Taiwan. *Bulletin of the Institute of Zoology Academia Sinica (Taipei)*, 20 (1): 43-47.
- Linnaeus, C. 1758.** *Systema naturæ per regna tria naturæ, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis.* Tomus I. Editio decima, reformata. Laurentii Salvii, Holmiæ. 10th Edition: 824 pp
- Ljubisavljevic, K., O. Arribas, G. Duki & S. Carranza. 2007.** Genetic and morphological differentiation of Mosor rock lizards, *Dinarolacerta mosorensis* (Kolombatović, 1886), with the description of a new species from the Prokletije Mountain Massif (Montenegro) (Squamata: Lacertidae). *Zootaxa* 1613: 1–22

- Lönnberg, E. 1907.** Reptilia and Batrachia. In: Sjöstedt, Y. (Hrsg.): *Wissenschaftliche Ergebnisse der Schwedischen Zoologischen Expedition nach dem Kilimandjaro, dem Meru und den umgebenden Massaisteppen* 1905-1906. - Upsala, No. 4, 1-28
- López-Jurado, L.F. 1989.** A new Canarian lizard subspecies from Hierro Island (Canarian archipelago). *Bonner zoologische Beiträge* 40: 265-272
- Loveridge, A. 1922.** New reptiles from Tanganyika Territory. *Proceedings of The Zoological Society of London* 1922: 313-315
- Loveridge, A. 1936.** A new lacertid (*Bedriagaia moreaui*) from Tanganyika Territory. *Proceedings of the New England Zoölogical Club* 15: 67-70
- Lue, K-Y. and S-M. Lin 2008.** Two New Cryptic Species of *Takydromus* (Squamata: Lacertidae) from Taiwan. *Herpetologica* 64 (3): 379-395
- Lukina, G.P. 1963.** Sistematiceskoje položenije i biologija skalnoj jascericy *Lacerta saxicola* Eversmann na sewerizoadbih granice areala wida w predjelach Kawkaza. *Izv. Akademia Nauk Azerbajdzan* 6: 53-61
- Lutz, D., W. Bischoff & W. Mayer. 1986.** Chemosystematische Untersuchungen zur Stellung von *Lacerta jayakari* Boulenger, 1887, sowie der Gattungen *Gallotia* Boulenger und *Psammodromus* Fitzinger (Sauria; Lacertidae). *Zeitschrift für Zoologische Systematik und Evolutionsforschung* 24: 144-157
- Lymberakis, P., N. Poulakakis, A. Kaliontzopoulou, E. Valakos & M. Mylonas. 2008.** Two new species of *Podarcis* (Squamata; Lacertidae) from Greece. *Systematics and Biodiversity* 6 (3): 307–318
- Makin, J.C. 1805.** *Lacerta montana* In: Sturm, J.: *Deutschlands Fauna in Abbildungen nach der Natur mit Beschreibungen* 3 Abtheilung 4: 17-19
- Martín, A. 1985.** Los lagartos de los roques del norte de Tenerife. *Bonner zoologische Beiträge* 36: 517-528
- Marx, H. 1956.** A new lacertid lizard from Angola. *Fieldiana: Zoology* 39: 5-9
- Mateo, J.A., M. Garcia-Marquez, L.F. Lopez-Jurado F. Barahona. 2001.** Descripción del lagarto gigante de La Palma (Islas Canarias) a partir de restos subfósiles. *Revista Española de Herpetología* 15: 53-59
- Matschie, P. 1893.** Über einige von Herrn Oscar Neumann bei Aden gesammelte und beobachtete Säugethiere, Reptilien und Amphibien. *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin* (1893): 24-31
- Mayer, W. & W. Böhme. 2000.** A note on the validity and distribution of *Zootoca vivipara sachalinensis*. *Casopis Narodniho Muzea Rada Prirodovedna* 169(1-4):123-124.
- Méhely, L. 1904.** Eine neue *Lacerta* aus Ungarn. *Annales historico-naturales Musei nationalis Hungarici* (Budapest) 2: 362-367
- Méhely, L. 1909.** Materialien zu einer Systematik und Phylogenie der muralis-ähnlichen Lacerten. *Annales historico-naturales Musei nationalis Hungarici*, Budapest, 7 (2): 409-621
- Ménétries, E. 1832.** Catalogue raisonné des objets de zoologie recueillis dans un voyage au caucase et jusqu'aux frontières actuelles de la Perse. *L'Académie Impériale des Sciences, St. -Pétersbourg* 1-271
- Merrem, B. 1820.** Versuch eines Systems der Amphibien I (*Tentamen Systematis Amphibiorum*). J. C. Krieger, Marburg, 191 pp.

- Mertens, R. 1921.** Zur Kenntnis der Reptilienfauna von Malta. *Zoologischer Anzeiger* 53: 236-240
- Mertens, R. 1921.** Eine neue Eidechse von den Pityusen. *Senckenbergiana* 3 (5): 142-144
- Mertens, R. 1922.** *Lacerta strigata wolterstorffi* subsp. nov. *Archiv für Naturgeschichte*, Berlin, 88 A (3): 193-195.
- Mertens, R. 1926.** Zwei neue Inselrassen der Gattung *Lacerta*. *Zoologischer Anzeiger* 68: 319-322
- Mertens, R. 1927.** Herpetologische Mitteilungen. XVI. Eine neue Rasse von *Lacerta bedriagae* Camerano. *Senckenbergiana* 8 (3/4): 178-180
- Mertens, R. 1927.** Über die Rassen der Pityuseneidechse (*Lacerta pityusensis* Bosca). *Zoologischer Anzeiger* 69: 299 - 304
- Mertens, R. 1929.** Zur Kenntnis der Eidechsenfauna Nordwest-Spaniens. *Senckenbergiana*, Frankfurt/Main, 11: 282-289.
- Mertens, R. 1932.** Zur Verbreitung und Systematik einiger *Lacerta* — Formen der Apenninischen Halbinsel und der Tyrrhenischen Inselwelt. *Senckenbergiana* 14: 235-259
- Mertens, R. 1937.** Neues über die Eidechsen-Fauna Istriens. *Zoologischer Anzeiger* 119: 332-336
- Mertens, R. 1938.** Über eine herpetologische Sammlung aus dem Gebiete des Njarasa-Grabens Ost-Afrikas. *Veröffentlichungen aus dem Deutschen Kolonial- und Übersee-Museum in Bremen* 2 (1): 1-9
- Mertens, R. 1938.** Zur Kenntnis der Reptilien-Fauna der Lüderitz-Bucht, Südwest-Afrika. *Veröffentlichungen aus dem Deutschen Kolonial- und Übersee-Museum in Bremen* 2: 10-16
- Mertens, R. 1938.** Eine melanistische Rasse der Madeira-Eidechse. *Senckenbergiana* 20 (3/4): 287-290
- Mertens, R. & L. Müller. 1940.** *Podarcis pityusensis* - Die Amphibien und Reptilien Europas. Zweite Liste, nach dem Stand vom 1. Januar 1940. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, 451: 36-39
- Mertens, R. 1942.** Die Mauereidechse von Isola Bella. *Zoologischer Anzeiger* 138: 44-47
- Mertens, R. 1952.** Neue Eidechsenrassen von den Liparischen Inseln. *Senckenbergiana* 32: 309-314
- Mertens, R. 1954.** Neue Eidechsen aus Südwest-Afrika. *Senckenbergiana*, 34 (4/6): 175-183
- Mertens, R. 1955.** Die Amphibien und Reptilien Südwestafrikas. Aus den Ergebnissen einer im Jahre 1952 ausgeführten Reise. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* (Frankfurt), 490: 1-172, Taf. 1-24.
- Mertens, R. 1955.** Die Mauereidechsen der Liparischen Inseln, gesammelt von Dr. Antonino Trischitta. *Senckenbergiana Biologica* 36: 25-40
- Mertens, R. 1959.** Zur Kenntnis der Lacerten auf der Insel Rhodos. *Senckenbergiana biologica* 40: 15-24
- Mertens, R. 1961.** Die Inseleidechsen des Golfes von Salerno. *Senckenbergiana biologica* 42 (1/2): 31-40.
- Mertens, R. 1967.** Die Reptilien der Pontinischen Inseln. *Senckenbergiana Biologica* 48 (2): 125-144

- Milne-Edwards, M.H. 1829.** Recherches zoologiques pour servir à l'histoire des lézards, extraites d'une monographie de ce genre. *Annales des science naturelles : zoologie* Paris 16: 50-89
- Mocquard, F. 1888.** Sur une Collection de Reptiles et de Batraciens rapportes de pays Comalis et de Zanzibar par R.G. Revoil. *Mémoires publiés par la Société philomathique à l'occasion du centenaire de sa fondation 1788-1888.* Société philomathique de Paris 5: 190-206
- Modrý, D., P. Necas, L. Rifai, W. Bischoff, N. Hamidan & Z. Amr. 2013.** Revision of the Levantine “*Lacerta*” *laevis/kulzeri*- Complex: 3. The Rock Lizard of Wadi Ramm, *Phoenicolacerta kulzeri khazaliensis* ssp. n. *Vertebrate Zoology* 63(3):307–312
- Monard, A. 1940.** Résultats de la mission du Dr. Monard en Guinée Portugaise 1937 – 1938. *Arquivos do Museum Bocage*, Lisbon 11: 147-182
- Monard, A. 1949.** Vertébrés nouveaux du Cameroun. *Revue Suisse de Zoologie* 56 (38): 731-745
- Moravec, J. 1994.** A new lizard from Iran, *Eremias (Eremias) laezharica* sp. n. (Reptilia: Lacertilia: Lacertidae). *Bonner zoologische Beiträge*. 45: 61-66
- Moravec, J., S.B. El Din, H. Seligmann, N. Sivan & Y.L. Werner. 1999.** Systematics and distribution of the *Acanthodactylus pardalis* group (Lacertidae) in Egypt and Israel. *Zoology in the Middle East* 17: 21-50
- Mozaffari, O. & J.F. Parham. 2007.** A new species of racerunner lizard (Lacertidae: *Eremias*) from Iran. *Proceedings of the California Academy of Sciences* 58 (23-30): 569-574
- Mozaffari, O. & F. Ahmadzadeh & J.F. Parham. 2011.** *Eremias papenfussi* sp. nov., a new lacertid lizard (Sauria: Lacertidae) from Tehran Province, Iran. *Zootaxa* 3114: 57-62
- Müller, L. 1927.** Beitrag zur Kenntnis der Rassen von *Lacerta lilfordi* Gthr. *Zoologischer Anzeiger* 73 (11/12): 257-269
- Müller, L. 1927.** Zwei weitere Rassen der *Lacerta lilfordi* Gthr. *Zoologischer Anzeiger* 74 (7/10): 185-194
- Müller, L. 1928.** Zweiter Beitrag zur Kenntnis der Rassen von *Lacerta lilfordi* Gthr. *Zoologischer Anzeiger*, 78: 261-273
- Müller, L. & O. Wettstein. 1932.** Über eine neue *Lacerta*-Form aus dem Libanon. *Zoologischer Anzeiger* 98: 218-223
- Müller, L. 1935.** Über die Smaragdeidechse der Kykladen-Insel Milos. *Zoologischer Anzeiger* 109: 225 – 236
- Müller, L. & W. Hellmich. 1937.** Mitteilungen über die Herpetofauna der Iberischen Halbinsel. II. Zur Kenntnis der *Lacerta monticola*. *Zoologischer Anzeiger* 117 (3/4): 65-73
- Müller, L. 1938.** Über die Mauereidechse der Insel Gerakunia. *Blätter für Aquarien und Terrarienkunde*, 1938 (3): 38-40.
- Munkhbayar, K. & L.J. Borkin. 2010.** A new subspecies of the lizard *Eremias multiocellata tsaganbogensis* subsp. nov. (Lacertidae) from southern Mongolia. *Izvestiya Samarskogo Nauchnogo Centra Rossiskaya Akademii Nauk*, 12(1)(No 33): 122-124
- Neumann, O. 1900.** Description of a new lizard of the genus *Nucras* from Usoga, British East Africa. *Annals and Magazine of Natural History* (7) 5: 56

- Nikolsky, A.M. 1897.** Les reptiles, amphibiens et poissons recueillis (part.) Mr. N. Zaroudny dans la Perse orientale *Annuaire Musée Zoologique de l'Académie Impériale des Sciences de St.-Petersbourg* 2: 306-348
- Nikolsky, A. M. 1898.** Dwa nowych wida jasceric iz Rossii [On two new lizards from Russia]. *Annuaire Musée Zoologique de l'Académie Impériale des Sciences de St.-Petersbourg* 3: 284-288
- Nikolsky, A. M. 1905.** Herpetologia rossica. *Mémoires Académie Impériale des Sciences, Physico-Mathématique Classe*, St. Pétersbourg, XVII (1): 1-518
- Nikolsky, A.M. 1915.** *Faune de la Russie et des Pays Limitrophes*. Reptiles, Vol. I (Chelonia et Sauria). Petrograd. 532 pp.
- Nilson, G., N. Rastegar-Pouyani, E. Rastegar-Pouyani and C. André. 2003.** Lacertas of South and Central Zagros Mountains, Iran, with descriptions of two new taxa. *Russian Journal of Herpetology*, 10 (1): 11-24.
- Nouria, S. & C.P. Blanc. 1998.** Description d'une nouvelle espece d'acanthodactyle de Tunisie: *Acanthodactylus mechriguensis*. *Atti e Memorie Ente Fauna Siciliana* 5: 103-110
- Orlova, V.F., N.A. Poyarkov, M.A. Chirikova, R.A. Nazarov, M. Munkhbaatar, K. Munkhbayar, K. Terbish. 2017.** MtDNA differentiation and taxonomy of Central Asian racerunners of *Eremias multiocellata-E. przewalskii* species complex (Squamata, Lacertidae). *Zootaxa* 4282, No 1: 1–42
- Oudart, P-L., P.B. Webb & S. Berthelot. 1839.** *Histoire naturelle des Iles Canaries*. Vol. 2 (2): pl. 1, Paris Béhune, éditeur, 1836-1850
- Palacios, F. & J. Castroviejo. 1975.** Descripción de una nueva subespecie de lagarto ágil (*Lacerta agilis garzoni*) de los Pirineos. *Doñana, Acta Vertebrata*, 2(1): 5-24
- Pallas, P.S. 1771.** Reise durch verschiedene Provinzen des Russischen Reiches, vol. 1. *Kaiserlichen Akademie der Wissenschaften*, St. Petersburg, 504pp
- Pallas, P.S. 1773.** Reise durch verschiedene Provinzen des Russischen Reiches, vol. 2. *Kaiserlichen Akademie der Wissenschaften*, St. Petersburg, 744pp
- Pallas, P.S. 1814.** Zoographia Rosso-Asiatica, sistens omnium in extenso Imperio Rossico , et adjacentibus maribus observatorum recensionem, domicilia, mores et descriptiones, anatomen atque icones plurimorum. Vol. 3., *Caesarea Academie Scientiarum*; Petropoli. 428pp
- Pamstruch, J.W. & O. Swartz, 1808.** Grä Ödla. *Svensk Zoologi*, Vol 1: 27-32
- Parker, H.W. 1932.** Scientific results of the Cambridge expedition to the east African lakes, 1930-31. 5. Reptiles and amphibians. *The Journal of the Linnean Society of London. Zoology*, 38: 213-229.
- Parker, H.W. 1935.** Two new lizards from Somaliland. *Annals and Magazine of Natural History* (10) 16: 525
- Parker, H.W. 1942.** The lizards of British Somaliland. *Bulletin of the Museum of Comparative Zoology at Harvard* 91: 1—101
- Pavlov, P. 1932.** Liste des sauriens et serpents des collections du Musée Huang ho Pai Tien Tsin. *Publications du Musée Huang ho Pai Tien Tsin*, Tianjin, 12: 1-27
- Pereleshin, S. D. and P.V. Terentjev. 1963.** Materials on herpetofauna of Sakhalin Island and Kuril Islands. *Trudy Sakhalinsk KNII* 14: 3 – 29

Peracca, M. G. 1903. Descrizione di una nuova specie del Gen. *Lacerta* L. di Sardegna. *Bollettino dei Musei di Zoologia ed Anatomia comparata della R. Università di Torino*, 18 (458): 1-3

Peracca, M.G. 1917. Sopra una nuova specie di Lacertidae del gen. *Algiroides* dell'Uganda. *Atti Della R. Accademia Delle Scienze Di Torino. Classe Di Scienze Fisiche, Matematiche E Naturali* 52: 351–354.

Pérez-Mellado, V. 1981. La lagartija de bocage, *Podarcis bocagei* (Seoane, 1884). Primeros datos sobre su distribución, colorido y ecología. *Amphibia-Reptilia* 1: 253-268

Pérez-Mellado, V. & A. Salvador. 1988. The Balearic lizard *Podarcis lilfordi* (Günther, 1874) (Sauria, lacertidae) of Menorca. *Arquivos do Museu Bocage*, 1(10): 127-195

Peters, G. 1958. Die Zauneidechse des Kleinen Kaukasus als besondere Unterart - *Lacerta agilis brevicaudata* ssp.n. *Zoologische Jahrbücher. Abteilung für Systematik, Geographie und Biologie der Tiere*, Jena, 86b (1/2): 127-138.

Peters, G. 1960. Die Grusinische Zauneidechse *Lacerta agilis grusinica* nomen novum. *Zoologischer Anzeiger*, Jena, 165: 279-289.

Peters, G. 1964. Studien zur Taxonomie, Verbreitung und Ökologie der Smaragdeidechsen III. Die orientalischen Populationen von *Lacerta trilineata*. *Mitteilungen aus dem Zoologischen Museum in Berlin*, 40 (2): 186-250.

Peters, G. & T.A. Muskhelishvili. 1968. *Lacerta agilis ioriensis* – eine neue Subspecies der kaukasischen Zauneidechsen. *Zoologische Jahrbücher. Abteilung für Systematik, Geographie und Biologie der Tiere*, Jena, 95: 213-228.

Peters, W.C.H. 1854. Diagnosen neuer Batrachier, welche zusammen mit der früher (24. Juli und 17. August) gegebenen Übersicht der Schlangen und Eidechsen mitgeteilt werden. *Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich Preussische Akademie der Wissenschaften zu Berlin*. 1854: 614-628

Peters, W.C.H. 1854. Corrigenda. *Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich Preussische Akademie der Wissenschaften zu Berlin* 1854 (Dezember): 726

Peters, W.C.H. 1863. Über eine neue Schlangengattung, *Styporhynchus*, und verschiedene andere Amphibien des zoologischen Museums. *Monatsberichte der Königlich Preussische Akademie des Wissenschaften zu Berlin*. 1863 (October): 399-413

Peters, W.C.H. 1869. Eine Mittheilung über neue Gattungen und Arten von Eidechsen. *Monatsberichte der Königlich Preussische Akademie des Wissenschaften zu Berlin*, 1869: 57-66

Peters, W.C.H. 1869. Förtekning pa de af J. Wahlberg i Damaralandet insamlade Reptilierna. *Öfversigt af Kongl. Vetenskaps-akademiens Forhandlingar*, 26 (7): 657-662.

Peters, W.C.H. 1869. Über neue Gattungen und neue oder weniger bekannte Arten von Amphibien (*Eremias*, *Dicrodon*, *Euprepes*, *Lygosoma*, *Typhlops*, *Eryx*, *Rhynchonyx*, *Elapomorphus*, *Achalinus*, *Coronella*, *Dromicus*, *Xenopholis*, *Anoplodipsas*, *Spilotes*, *Tropidonotus*). *Monatsberichte der Königlich Preussische Akademie des Wissenschaften zu Berlin*, 1869: 432-447

Peters, W.C.H. 1870. Report on a collection made at Hantam (Oorlogsrevier, S.W. Africa). *Monatsberichte der Königlich Preussische Akademie des Wissenschaften zu Berlin*, 1870: 110-115

- Peters, W.C.H. 1874.** Über einige neue Reptilien (*Lacerta*, *Eremias*, *Diploglossus*, *Euprepes*, *Lygosoma*, *Sepsina*, *Ablepharus*, *Simotes*, *Onychocephalus*). *Monatsberichte der Königlich Preussische Akademie des Wissenschaften zu Berlin*. 1874 (Juni): 368-377
- Peters, W.C.H. 1878.** Über die von Hrn. J. M. Hildebrandt während seiner letzten ostafrikanischen Reise gesammelten Säugethiere und Amphibien. *Monatsberichte der Königlich Preussische Akademie des Wissenschaften zu Berlin* 1878: 194-209
- Peters, W.C.H. 1881.** Einige herpetologische Mittheilungen. Ueber eine neue Art von *Tachydromus* aus dem Amurlande. *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin*, 1881: 71-72
- Peters, W.C.H. and G. Doria 1882.** Note erpetologiche e descrizione di una nuova specie di *Lacerta* delle Isole Canarie. *Annali del Museo Civico de Storia Naturale di Genova*. (ser. 1) 18: 431-434
- Pope, C. 1929.** Notes on reptiles from Fukien and other Chinese provinces. *Bulletin of the American Museum of Natural History*, 58: 335-487
- Prazak, J.P. 1898.** Systematische Übersicht der Reptilien und Batrachier Bohmens. *Zoologische Jahrbücher, Abteilung für Systematik Geographie und Biologie der Tiere* 11 (3): 1873-234
- Radovanovic, M. 1956.** Rassenbildung bei den Eidechsen auf adriatischen Inseln. *Österreichische Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Klasse, Denkschriften* 110: 1-82.
- Radovanovic, M. 1959.** Zum Problem der Speziation bei Inseleidechsen. *Zoologische Jahrbücher. Abteilung für Systematik, Geographie und Biologie der Tiere*, Jena, 86 (4/5): 395-436.
- Radovanović, M. 1970.** Herpetoloska proucavanja na Olibu i okolnim ostrvima. *Prirod. istrazivanja JAZU, Zagreb* 38 (Acta biologica 6): 59-74
- Rafinesque Schmaltz, C. S. 1810.** *Caratteri di alcuni nuovi generi e nuove specie di animali e piante della Sicilia*. Sanfilippo, Palermo, 105 pp.
- Rastegar-Pouyani, N. & G. Nilson 1998.** A new species of *Eremias* (Sauria: Lacertidae) from Fars Province, South-Central Iran. *Russian Journal of Herpetology* 4 (2): 94-101
- Rastegar-Pouyani, N. & G. Nilson. 1998.** A new species of *Lacerta* (Sauria: Lacertidae) from the Zagros mountains, Esfahan Province, West-Central Iran. *Proceedings of the California Academy of Sciences* 50 (10): 267-277
- Rastegar-Pouyani, N. and E. Rastegar-Pouyani. 2001.** A new species of *Eremias* (Sauria: Lacertidae) from highlands of Kermanshah Province, western Iran. *Asiatic Herpetological Research*. 9: 107-112
- Rastegar-Pouyani, N., H. Gholikami, M. Rajabzadeh, S. Shaffifei & S.C. Anderson. 2008.** Annotated Checklist of Amphibians and Reptiles of Iran. *Iranian Journal of Animal Biosystematics* 4 (1): 7-30
- Rastegar-Pouyani, N., R. Karamiani, H. Oraei, A. Khosrawani, E. Rastegar-Pouyani. 2011.** A New Subspecies of *Darevskia raddei* (Boettger, 1892) (Sauria: Lacertidae) from the West Azerbaijan Province, Iran. *Asian Herpetological Research* 3 (3): 216-222
- Rastegar-Pouyani, E., S. Hosseinian, S. Rafiee, H.G. Kami, M. Rajabzadeh & M. Wink. 2016.** A new species of the genus *Eremias* Fitzinger, 1834 (Squamata: Lacertidae) from Central Iran, supported by mtDNA sequences and morphology. *Zootaxa* 4132 (2): 207-220

Reuss, A. 1834. Zoologische Miscellen, Reptilien. Saurier. Batrachier. *Museum Senckenbergianum, Frankfurt am Main* 1 (6): 27-62

Risso, A. 1826. *Histoire Naturelle des principales productions de l'Europe Méridionale* et particulièrement de celles des Environs de Nice et des Alpes Maritimes – III. Levraut, Paris.

Roux, J. 1907. Beiträge zur Kenntnis der Fauna von Süd-Afrika. Ergebnisse einer Reise von Prof. Max Weber im Jahre 1894. VII. Lacertilia (Eidechsen). *Zoologische Jahrbücher. Abteilung für Systematik, Geographie und Biologie der Tiere*, Jena, 25: 403-444

Rykena, S., H-K. Nettmann and W. Mayer 2001. *Lacerta viridis guentherpetersi* ssp. nov. eine neue Unterart der Smaragdeidechse aus Griechenland. In: Kerstin Elbing and Hans-Konrad Nettmann (eds.) Beiträge zur Naturgeschichte und zum Schutz der Smaragdeidechsen (*Lacerta* s. str.). *Contributions to Mertensiella* 13: 89-97

Sánchez-Vialas, A., M. Calvo-Revuelta, J.L. Rubio, F. Palacios, M. García-París. 2019. Taxonomic and nomenclatural status of Iberian *Algyroides* (Lacertidae). *Amphibia-Reptilia* 40: 91–102

Salvador, A. 1979. Materiales para una „Herpetofauna balearica“. 2. Taxonomia de las lagartijas baleares del archipiélago de Cabrera. *Bonner zoologische Beiträge* 30, 1/2: 176-191

Salvador, A. 1979. Una nueva subespecie melánica de lagartija balear (*Lacerta lilfordi*). *Boletín de la Real Sociedad Española de Historia Natural*, 77: 491-492

Salvador, A 1982. A revision of the lizards of the genus *Acanthodactylus* (Sauria: Lacertidae). *Bonner zoologische Monographien* (No. 16) 1982: 1-167

Sauvage, H.-E. 1884. Sur quelques Reptiles de la collection du Muséum d'Histoire Naturelle. *Bulletin de la Société philomathique de Paris*, (7) 8: 142-146

Schenkel, E. 1901. Achter Nachtrag zum Katalog der herpetologischen Sammlung des Basler Museums. *Verhandlungen der Naturforschenden Gesellschaft in Basel* 13:142-199

Schinz, K.R. 1833. *Naturgeschichte und Abbildungen der Reptilien*. Wiedmann, Leipzig, 240 pp.

Schlegel, H. 1838. Reptilia elaborantibus. Saurii et batrachii. In: Temminck, C. J. & H. Schlegel: *Fauna Japonica* sive Descriptio animalium, quae in itinere per Japonianum, jussu et auspiciis superiorum, qui summum in India. Batavorum Imperium tenent. Pp. 85-144

Schmidt, K.P. 1919. Contributions to the Herpetology of the Belgian Congo based on the Collection of the American Congo Expedition, 1909-1915. Part I: turtles, crocodiles, lizards, and chamaeleons. *Bulletin of the American Museum of Natural History* 39 (2): 385-624

Schmidt, K.P. 1925. New Chinese amphibians and reptiles. *American Museum Novitates* (175): 1-3

Schmidt, K.P. 1939. Reptiles and amphibians from Southwestern Asia. *Field Museum of Natural History Publications, Zoology Ser.*, 24: 49-92

Schmidt, K.P. & H. Marx. 1957. Results of the namru-3 southeastern Egypt Expedition, 1954. 2. Reptiles and amphibians. *Bulletin, Zoological Society of Egypt*, Cairo, 13: 16-27

Schmidtler, J.F. 1975. Zur Taxonomie der Riesen-Smaragdeidechsen (*Lacerta trilineata* Bedriaga) Süd-Anatoliens. *Veröffentlichungen der Zoologischen Staatssammlung München* 18: 45-68.

Schmidtler, J.F. 1986. Orientalische Smaragdeidechsen: 1. Zur Systematik und Verbreitung von *Lacerta viridis* in der Türkei. *Salamandra* 22 (1): 29-46

Schmidler, J. F., J. Eiselt, I.S. Darevsky. 1994. Untersuchungen an Feldeidechsen (*Lacerta-saxicola*-Gruppe) in der östlichen Türkei: 3. Zwei neue parthenogenetische Arten. *Salamandra* 30 (1): 55-70

Schmidler, J. F. & W. Bischoff. 1999. Revision des levantinischen *Lacerta laevis/kulzeri*-Komplexes: 1. Die Felseneidechse *Lacerta cyanisparsa* sp.n. *Salamandra* 35 (3): 129-146

Schreiber, E. 1891. Ueber die typischen Exemplare von *Lacerta mosorensis* Kolomb. mit 1 Tafel. *Annalen des k.k. naturhistorischen Hofmuseums in Wien* 7 (3): 235-240

Schreiber, E. 1912. *Herpetologia Europaea* (2. Aufl.). Fischer (Jena), 960 pp

Scortecci, G. 1934. Descrizione preliminare di una nuova specie del genere *Philochortus* (*Philochortus zolii*) della zona di Gat (Missione della reale Societa Geografica). *Atti della Società Italiana di Scienze Naturali* 73: 305-308

Segoli, M., T. Cohen, Y.L. Werner. 2002. A new lizard of the genus *Mesalina* from Mt. Sinai, Egypt (Reptilia: Squamata: Sauria: Lacertidae). *Faunistische Abhandlungen* (Dresden) 23 (1):157-176

Seoane, V.L. 1885. Identidad de *Lacerta schreiberi* (Bedriaga) y *Lacerta viridis* var. *Gadovii* (Boulenger), é investigaciones. *Herpetologicas de Galicia. La Coruña* [1884]: 19 pp.

Sheppard, R. 1804. Description of British Lizards, and a new British species of viper. *Transactions of the Linnean Society of London*, 7:49-56

Sindaco, R., M. Simó-Riudalbas, R. Sacchi, S. Carranza. 2018. Systematics of the *Mesalina guttulata* species complex (Squamata: Lacertidae) from Arabia with the description of two new species. *Zootaxa* 4429 (3): 513–547

Šmíd, J., J. Moravec, V. Gvoždík, J. Štundl, D. Frynta, P. Lymberakis, P. Kapli, T. Wilms, A. Schmitz, M. Shobrak, S.H. Yousefkhani, E. Rastegar-Pouyani, A.M. Castilla, J. Els & W. Mayer. 2017. Cutting the Gordian Knot: Phylogenetic and ecological diversification of the *Mesalina brevirostris* species complex (Squamata, Lacertidae). *Zoologica Scripta*. 2017:1–16.

Smith, A. 1838. Contributions to the Natural History of Southern Africa. Art. VIII. *Magazine of Natural History*, London, 2 (14): 92-94

Smith, A. 1845. *Illustrations of the zoology of South Africa*, Reptilia. Smith, Elder, and Co., London

Smith, A. 1849. *Tropidosaura dumerilii* - *Illustrations of the zoology of South Africa*, III, Appendix: 7.

Sobolewsky, N. I. 1930. A new form of the genus *Lacerta* (Reptilia) from Hungary. *Studies of the Association of Research Institutes at the Physics and Mathematics Faculty of Moscow State University* 3, 1A: 3—9

Sobolewsky, N. I. 1930. ? . *Studies of the Association of Research Institutes at the Physics and Mathematics Faculty of Moscow State University*, 3 2a: 129.

Steindachner, F. 1870. Herpetologische Notizen (II). Reptilien gesammelt Während einer Reise in Sengambien. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien*, 62: 326-348

Steindachner, F. 1882. Über eine neue *Eremias*-Art aus dem Tale des Limpopo-(Krokodilflusses) in Transvaal. (E. holubi). *Sitzungsberichte der Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Classe*. Wien, 86: 83-85

Steindachner, F. 1889. Über eine von Prof. O. Simony auf den Roques del Zalmor bei Hierro, Canarische inseln, entdeckte neue Eidechsenart. *Anzeiger der Akademie der Wissenschaften in Wien* 1889: 260-262

Steindachner, F. 1891. Ueber neue und seltene Lacertiden aus der herpetologischen Sammlung des k. k. naturhistorischen Hofmuseums. *Annalen des Naturhistorischen Museums in Wien* 6: 371-378

Stejneger, L. 1893. On some collections of Reptiles and Batrachians from East Africa and the adjacent islands, received from Dr. W.L. Abbott and Mr. William Astor Chanler, with descriptions of new species. *Proceedings of the United States National Museum*, 16: 711-741.

- Stejneger, L. 1904.** A new species of lizard from the Riu Kiu Archipelago, Japan. *Smithsonian Miscellaneous Collections*, 47: 294-295.
- Stejneger, L.H. 1924.** Herpetological novelties from China. *Occasional papers of the Boston Society of Natural History*, 5: 119-121.
- Stemmler, O. 1962.** Ein Beitrag zur Biologie und Systematik von *Lacerta bedriagae*. *Aquaria* 1962: 5-12
- Sternfeld, R. 1911.** Zur Herpetologie Südwestafrikas. *Mitteilung aus dem Zoologischen Museum in Berlin*, 5: 393-411
- Sternfeld, R. 1912.** IV. Zoologie II. Lfg. Reptilia. In: Schubotz., R. (ed.): *Wissenschaftliche Ergebnisse der Deutschen Zentral-Afrika Expedition 1907- 1908*, unter Führung A. Friedrichs, Herzogs zu Mecklenburg. Klinkhard und Biermann, Leipzig: 197-279
- Stoliczka, F. 1872.** Notes on reptiles collected by surgeon F. Day in Sind. *Proceedings of the Asiatic Society of Bengal* 1872: 85-92
- Stoliczka, F. 1872.** Notes on some new species of Reptilia and Amphibia collected by Dr. W. Waagen in north-Western Punjab. *Proceedings of the Asiatic Society of Bengal* 1872: 124-131
- Strauch, A. 1867.** Bemerkungen über die Eidechsen-Gattung *Scapteira* (Fitz.). *Mélanges biologiques tirés du Bulletin de l'Académie impériale des sciences de St. Pétersbourg* 4: 403-426
- Strauch, A.A. 1876.** Reptiles. In: Przewalski, N.M. 1875. *Mongolia, the Tangut country, and the solitudes of northern Tibet, being a narrative of three years' travel in eastern high Asia*. - London. Sampson Low, Marston, Searle & Rivington. 1876. vol. 1 and 2.
- Suchov, G.F. 1927.** Die Zauneidechse aus der Krim (*Lacerta agilis tauridica* subsp. nov.). *Mémoires de la Classe des Sciences Physiques et Mathématiques de l'Académie des Sciences de l'Ukraine*. Tome IV: 83-87.
- Suchov, G.F. 1929.** Description of a new species of lizards from the environs of Vladicaucasus (*Lacerta boemica* sp.nov.) *Academie des Sciences de l'Ukraine, Mémoires de la Classendes Sciences Physiques et Mathématiques*. Tome XIII. Livr. 1: 117-119.
- Suchow, G. F. 1936.** Eine neue Unterart der Eidechse aus dem persischen Kurdistan. *Travaux de l'Institut Zoologique de l'Academie des Sciences URSS*, 3:303-308, 2 figs
- Szczerbak, N.N. 1962.** O sistematike skalnych jasceric *Lacerta saxicola* Evers mann Kryma i Sewernowo Kawkaza. *Zoologicheskij Zhurnal, Moskwa*, 11: 1374-1385
- Szczerbak, N.N. 1970.** New subspecies *Eremias przewalskii tuvensis* sp. n. (Sauria, Reptilia) from the Tuva Autonomous Soviet Specialist Republic and data on the species systematics as a whole. *Vestnik Zoologii* 1970(3):31-36
- Szczerbak, N.N. 1972.** New subspecies of *Eremias strauchi*-*Eremias strauchi kopetdaghica* ssp. nova (Sauria, Reptilia) from Turkmenia. *Vestnik Zoologii* 1972 (6): 83-86
- Szczerbak, N.N. & E.V. Nashetko. 1973.** New subspecies – *Eremias scripta pherganensis* ssp. n. (Reptilia, Sauria) from Uzbekistan. - *Vestnik zoologii, Journal of the Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine, Kiev*, 1973 (1): 84-86.

- Szczerbak, N.N. 1973.** New subspecies of *Eremias multiocellata* – *Eremias multiocellata bannikowi* ssp. n. (Reptilia, Sauria) from Tuva and north-western Mongolia. *Vestnik zoologii, Journal of the Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine, Kiev*, 1973 (3): 84-87
- Taddei, A. 1949.** Le Lacerte (Archaeolacerte e Podarcis) dell'Italia peninsulare e delle Isole. *Commentationes Pontificia Academia Scientiarum*, 13 (4): 197-274
- Taddei, A. 1949.** Le Lacerte (*Podarcis*) delle Isole dell'Archipelago Toscano. *Monitore Zoologico Italiano*, Firenze, 57: 12-34
- Taddei, A. 1950.** Le Lacerte (*Archaeolacerta* e *Podarcis*) dell'Istria e della Dalmazia. *Pontificia Academia Scientiarum Commentationes* 14: 137-166
- Taddei, A. 1950.** Le Lacerta (*Lacerta*) in Italia. *Pontificia Academia Scientiarum Commentationes*, Roma 14 (5): 197-219
- Taddei, A. 1953.** Contributo allo studio delle lucertole del Monte Argentario (Toscana). *Monitore Zoologico Italiano*, Firenze, 60: 84-90
- Takeda, N. & H. Ota. 1996.** Description of a new species of *Takydromus* from the Ryukyu Archipelago, Japan, and a taxonomic redefinition of *T. smaragdinus* Bouleger, 1887 (Reptilia: Lacertidae). *Herpetologica* 52 (1): 77-88
- Tamar, K., P. Geniez, J.C. Brito & P-A. Crochet. 2017.** Systematic revision of *Acanthodactylus busacki* (Squamata: Lacertidae) with a description of a new species from Morocco. *Zootaxa*, 4276 (3): 357-386.
- Tornier, G. 1900.** Neue Liste der Crocodile, Schildkröten und Eidechsen Deutsch-Ost-Afrikas. *Zoologische Jahrbücher. Abteilung für Systematik, Geographie und Biologie der Tiere*, Jena 13: 579-618
- Tornier, G. 1902.** Herpetologisch Neues aus Ostafrika. *Zoologischer Anzeiger* 25: 700-704
- Tornier, G. 1905.** Schildkröten und Eidechsen aus Nord-Ost Afrika und Arabien. Aus Carlo v. Erlanger's und Oscar Neumann's Forschungsreise. *Zoologische Jahrbücher. Abteilung für Systematik, Ökologie und Geographie der Tiere* 22: 365-388
- Trape, J.F., S. Trape & L. Chirio. 2012.** *Lézards, crocodiles et tortues d'Afrique occidentale et du Sahara*. IRD Orstom, 503 pp
- Tsaruk, O.J. 1986.** Variability of the head pholidosis and intraspecific taxonomy of *Eremias arguta* (Pallas, 1773). *Trudy Zoologicheskogo Instituta Akademii Nauk SSSR, Leningrad*, 157: 203-214.
- Tschernow, N. 1934** ?
- Tschudi, J. J. 1836.** Über ein neues subgenus von *Lacerta* Cuv. *Isis von Oken* 29: columns 546–551
- Tuniyev, S. B. and B.S. Tuniyev. 2008.** Intraspecific Variation of the Sand Lizard (*Lacerta agilis*) from the Western Caucasus and Description of a New Subspecies *Lacerta agilis mzymtensis* ssp. nov. (Reptilia: Sauria). *Russian Journal of Herpetology* 15 (1): 55-66
- Tuniyev, S.B., I.A. Doronin, A.A. Kidov, B.S. Tuniyev. 2011.** Systematic and Geographical Variability of Meadow Lizard, *Darevskia praticola* (Reptilia: Sauria) in the Caucasus. *Russian Journal of Herpetology* 18 (4): 295–316

- Tuniyev, B.S., I. Doronin, B.Tuniyev, A. Aghasyan, A. Kidov, L. Aghasyan. 2013.** New Subspecies of Meadow Lizard, *Darevskia praticola loriensis* ssp. nov. (Reptilia: Sauria) from Armenia. *Russian Journal of Herpetology* 20 (3): 223-237
- Tuniyev, B.S., T.V. Petrova. 2019.** A new lizard species of the genus *Darevskia* Arribas, 1997 from Southern Armenia. *Proceedings of the Zoological Institute of the Russian Academy of Sciences*. 323 (2):. 136–148
- Vaillant, M.L. 1882.** Reptiles et Batraciens. In: Révoil: *Mission G. Révoil aux pays Comalis, Faune et Flore*. Faune et Flore des Pays Comalis, Afr. orient., Paris. 1-25, 3 plates
- Valverde, J.A. 1958.** Una nueva lagartija del género *Algiroides* Bribron procedente de la Sierra de Cazorla (Sur de España). *Archivos del Instituto de Aclimatación de Almería* , Almeria 16: 127-134
- Van Denburgh, J. 1909.** New and previously unrecorded species of reptiles and amphibians from the island of Formosa. *Proceedings of the California Academy of Sciences* (4) 3 (3): 49-56.
- Van Denburgh, J., 1912.** Concerning certain species of reptiles and amphibians from China, Japan, the Loo Choo Islands, and Formosa. *Proceedings of the California Academy of Sciences (Series 4)* 3 (10): 187-258
- Vasvári, M. 1926.** Contributions to the knowledge of the form-group of *Lacerta viridis* Laur. *Állattani Közlemények* 23: 34–66, 129–132
- Vicente, L.A. 1985.** Description d'une nouvelle sous-espèce de *Podarcis bocagei* (Seoane, 1884) (Sauria, Lacertidae) de l'île de Berlenga: *Podarcis bocagei berlengensis*. *Bulletin du Muséum national d'histoire naturelle, Paris*, 4e sér. 7 (section A): 267-274.
- Vogt, T. 1914.** Südchinesische Amphibien und Reptilien. *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin*, 1914: 96-102
- Wagler, J. 1830.** Natürliches System der Amphibien, mit vorangehender Classification der Säugthiere und Vögel. *Ein Beitrag zur vergleichenden Zoologie*. Munich, Stuttgart and Tübingen: J.G. Cotta. vi + 354 pp. + one plate
- Wagner, P., E. Greenbaum, P. Malonza & B. Branch. 2014.** Resolving sky island speciation in populations of East African *Adolfus alleni* (Sauria, Lacertidae). *Salamandra* 50 (1): 1-17
- Walley, H.D. 1958.** A new lacertid lizard from Korea. *Herpetologica* 14: 203-205
- Wang, Y-Y, S-P Gong, P. Liu & X. Wang. 2017.** new species of the genus *Takydromus* (Squamata: Lacertidae) from Tianjingshan Forestry Station, northern Guangdong, China. *Zootaxa* 4338 (3): 401–440
- Werner, F. 1891.** Beiträge zur Kenntnis der Reptilien und Amphibien von Istrien und Dalmatien. *Verhandlungen der kaiserlich-königlichen zoologisch-botanischen Gesellschaft in Wien* 41: 751-768
- Werner, F. 1896.** Über einige Reptilien aus Usambara (Deutsch-Ostafrika). *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien*, 45:190—194
- Werner, F. 1897.** *Die Reptilien und Amphibien Oesterreich-Ungarns und der Occupationsländer*. Verlag von A. Pichler's Witwe & Solin, Wien: 161 pp
- Werner, F. 1900.** Beschreibung einer bisher noch unbekanntes Eidechse aus Kleinasien *Lacerta anatolica*. *Anzeiger der Akademie der Wissenschaften in Wien* 25: 269-271
- Werner, F. 1902.** Ueber Reptilien und Batrachier aus West-Asien (Anatolien und Persien). *Zoologische Jahrbücher Abteilung für Systematik, Geographie und Biologie der Tiere*. 19: 329-345
- Werner, F. 1902.** Reptilien. In: E. Gaivigni: Beiträge zur Kenntniss der Fauna einiger dalmatinischer Inseln. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien* 52: 381-388

- Werner, F. 1904.** Zur Kenntnis der *Lacerta danfordi* Guthr. und der oxycephalen Eidechsen überhaupt. *Zoologischer Anzeiger* 27: 254-259
- Werner, F. 1904.** Beschreibung neuer Reptilien aus den Gattungen *Acanthosaura*, *Calotes*, *Gastropholis* und *Typhlops*. *Zoologischer Anzeiger* 27: 461-464
- Werner, F. 1908.** Die zoologische Reise des naturwissenschaftlichen Vereines nach Dalmatien im April 1906, B. Spezieller Teil. Bearbeitung des gesammelten Materials. 4. Reptilien und Batrachier. *Mitteilungen des naturwissenschaftlichen Vereines an der Universität Wien* 6 : 44-53. Tafel 1.
- Werner, F. 1913.** Neue oder seltene Reptilien und Frösche des Naturhistorischen Museums in Hamburg. Reptilien der Ostafrika-Expedition der Hamburger Geographischen Gesellschaft 1911/12. Leiter: Dr. E. Obst. Reptilien und Amphibien von Formosa. *Jahrbuch der Hamburgischen Wissenschaftlichen Anstalten* , 30 [1912], 2. Beiheft: 1-39, 40-45, 45-51
- Werner, F. 1917.** Über einige neue Reptilien und einen neuen Frosch des Zoologischen Museums in Hamburg. *Mitteilungen aus dem Zoologischen Museum Hamburg*, 2. Beiheft zum Jahrbuch der Hamburgischen Wissenschaftlichen Anstalten 34: 31-36
- Werner, F. 1929.** Beiträge zur Kenntnis der Fauna von Syrien und Persien. *Zoologischer Anzeiger* 81 (7/10): 238-245
- Werner, F. 1929.** Wissenschaftliche Ergebnisse einer zoologischen Forschungsreise nach Westalgerien und Marokko. *Sitzungsberichte der Akademie der Wissenschaften in Wien Mathematisch-naturwissenschaftliche Klasse*, Abt. 1, 138; 1-34
- Werner, F. 1930.** Contribution to the knowledge of the reptiles and amphibians of Greece, especially the Aegean islands. *Occasional papers of the Museum of Zoology, University of Michigan* (211): 1-46
- Werner, F. 1931.** Ergebnisse einer zoologischen Forschungsreise nach Marokko. III. Unternommen 1930 mit Unterstützung der Akademie der Wissenschaften in Wien von Franz Werner und Richard Ebner. III. Amphibien und Reptilien. *Sitzungsberichte der Akademie der Wissenschaften in Wien Mathematisch-naturwissenschaftliche Klasse* Sect. 1, 140: 272-318
- Werner, F. 1933.** Ergebnisse einer zoologischen Studien- und Sammelreise nach den Inseln des Ägäischen Meeres. *Sitzungsberichte der Akademie der Wissenschaften in Wien Mathematisch-naturwissenschaftliche Klasse* 142 (3/4): 103-133
- Werner, F. 1935.** Reptilien der Ägäischen Inseln. *Akademie der Wissenschaften in Wien, Sitzungsberichte, Mathematisch-naturwissenschaftliche Klasse* 144: 81-117
- Werner, F. 1936.** Reptiles from Mount Troodos, Cyprus. *Proceedings of the Zoological Society of London* 1936: 655-658
- Werner, F. 1938.** Ergebnisse der achten zoologischen Forschungsreise nach Griechenland (Euboea, Tinos, Skiathos, Thasos usw.). *Sitzungsberichte der Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*. Wien, Abteilung 1, 147:151-173.
- Werner, Y. L. 2004.** A new species of the *Acanthodactylus pardalis* group (Reptilia: Lacertidae) from Jordan. *Zoology in the Middle East* 32: 39-46

- Werner, Y.L. & S. Ashkenazi. 2010.** Notes on some Egyptian Lacertidae, including a new subspecies of *Mesalina*, involving the Seligmann effect. *Turkish Journal of Zoology* 34: 123-133
- Wettstein, O. 1926.** Beitrag zur Systematik der adriatischen Inseleidechsen und Aufzählung der von P. Kammerer gesammelten Reptilien und Amphibien. In: P. Kammerer: *Die Artenwandel auf Inseln*. Deuticke, Wien und Leipzig
- Wettstein, O. 1931.** Neue oder wenig bekannte Eidechsen. *Zoologischer Anzeiger*, Leipzig (1931) 95: 280-291
- Wettstein, O. von. 1937.** Über Balearen-Eidechsen. *Zoologischer Anzeiger*, 117: 293-297
- Wettstein, O. 1937.** Vierzehn neue Reptilienrassen von den südlichen Ägäischen Inseln. *Zoologischer Anzeiger* 118: 79—90
- Wettstein, O. 1940.** Eidechsen aus dem Tien-Shan-Gebiet. *Zoologischer Anzeiger* 130 (3/4): 83-89
- Wettstein, O. 1952.** Dreizehn neue Reptilienrassen von den Ägäischen Inseln. *Anzeiger der Mathematisch-Naturwissenschaftliche Klasse, Österreichische Akademie der Wissenschaften*, Wien, 89:251-256.
- Wettstein, O. 1956.** Zwei neue Eidechsenrassen von den Kykladen. *Anzeiger der Mathematisch-Naturwissenschaftliche Klasse, Österreichische Akademie der Wissenschaften*, Wien 1956 (11): 1-2; 134-135.
- Wettstein, O. von. 1965.** Ergebnisse der von Dr. O. Paget und Dr. E. Kritscher auf Rhodos durchgeführten zoologischen Exkursionen. *Annalen des Naturhistorischen Museums in Wien*, 68: 635-640.
- Wiegmann, A.F.A. 1834.** *Herpetologica Mexicana, seu descriptio amphibiorum Novae Hispaniae*. Pars Prima, Saurorum Species, Berlin: C.G. Lüderitz. vi + 54 pp. + Plates I-X.
- Wiegmann, A.F.A. 1835.** Ueber *Amystes*, eine neue Gattung der Lacerten, ohne Augenlieder. *Archiv für Naturgeschichte* 1 (2): 1-5
- Witte, G.-F. de & R.F. Laurent. 1942.** Liste des Lacertidae du Congo belge et description d'une espèce nouvelle. - *Revue de Zoologie et de Botanique Africaines*, Bruxelles, 36 (2): 165-180
- Wolf, J. 1805.** *L. nigra*, *L. viridis*, *L. crocea*. In: Sturm, J.: *Deutschlands Fauna in Abbildungen nach der Natur mit Beschreibungen*. III. Abtheilung. Die Amphibien, 4. Heft. Nürnberg
- Yalçinkaya, D. & B. Göçmen. 2012.** A new subspecies from Anatolia, *Acanthodactylus schreiberi* Boulenger, 1879 *ataturi* n. ssp. (Squamata: Lacertidae). *Biharian Biologist* 6 (1): 19-31
- Zarevsky, S.T. 1918.** Reptiles et amphibiens, collectionnés par V.J. Lazdin dans la Boukharie méridionale et orientale ainsi que dans la partie N.-E. de la Province Transcaspienne en été 1915. *Annuaire Musée Zoologique de l'Académie Impériale des Sciences de St.-Pétersbourg*, 22: 79-90.
- Ziegler, T. & W. Bischoff. 1999.** *Takydromus (Platyplacopus) kuehnei vietnamensis* ssp. n., eine neue Schnellläufereidechsen-Unterart aus Vietnam (Reptilia: Squamata: Lacertidae). *Salamandra* 35 (4): 209-226

TAXON INDEX

A

abchastica (see *Darevskia derjugini*)
acrolampra (see *Podarcis muralis*)
acutirostris (see *Eremias*)
addayae (see *Podarcis lilfordi*)
adjarica (see *Darevskia*)
adolffjordansi (see *Podarcis milensis*)
adramitana (see *Mesalina*)
adriaticus (see *Podarcis siculus*)
aegaea (see *Anatololacerta anatolica*)
aegyptius (see *Acanthodactylus*)
aemiliani (see *Podarcis siculus*)
aenofuscus (see *Takydromus sexlineatus*)
aeoli (see *Podarcis melisellensis*)
affinis (see *Podarcis pityusensis*)
afghanistanica (see *Eremias*)
africanus (see *Adolfus*)
ahmaddisii (see *Acanthodactylus*)
ahorcadosi (see *Podarcis pityusensis*)
aghasyani (see *Darevskia*)
agilis (see *Lacerta*)
albanica (see *Podarcis muralis*)
albiventris (see *Podarcis siculus*)
algae (see *Podarcis pityusensis*)
algirus (see *Psammodromus*)
alleni (see *Adolfus*)
alpina (see *Darevskia*)
alvearioi (see *Podarcis raffonei*)
amorgensis (see *Podarcis erhardii*)
amparoae (see *Podarcis siculus*)
amurensis (see *Takydromus*)
anatolica (see *Anatololacerta*)
anchietae (see *Meroles*)
annulifera (see *Pedioplanis lineocellata*)
antoninoi (see *Podarcis raffonei*)
appenninica (see *Podarcis muralis*)
arabicus (see *Acanthodactylus*)
aranica (see *Iberolacerta*)
aria (see *Eremias*)

andersoni (see *Eremias*)
andersonii (see *Latastia longicaudata*)
andreanskyi (see *Atlantolacerta*)
arenicola (see *Latastia boscai*)
argus (see *Eremias*)
argus (see *Lacerta agilis*)
arguta (see *Eremias*)
armeniaca (see *Darevskia*)
arnoldi (see *Mesalina*)
aporosceles (see *Eremias acutirostris*)
asper (see *Acanthodactylus boskianus*)
aspera (see *Pedioplanis lineocellata*)
astorgae (see *Podarcis siculus*)
astur (see *Iberolacerta monticola*)
asukului (see *Congolacerta*)
ataturi (see *Acanthodactylus schreiberi*)
atlantica (see *Gallotia*)
atlanticus (see *Acanthodactylus erythrurus*)
atratus (see *Podarcis liolepis*)
auaritae (see *Gallotia*)
audouini (see *Acanthodactylus scutellatus*)
aurantiaca (see *Nucras*)
aurelioi (see *Iberolacerta*)
aureus (see *Acanthodactylus*)
auroralis (see *Takydromus*)
australis (see *Australolacerta*)
austroarabica (see *Mesalina*)
ayunensis (see *Meroles*)

B

bagnolensis (see *Podarcis siculus*)
bahaeldini (see *Meroles*)
baldasseronii (see *Podarcis muralis*)
balearica (see *Podarcis lilfordi*)
balfouri (see *Mesalina*)
bannikowi (see *Eremias multiocellata*)
barabensis (see *Zootoca vivipara*)
barani (see *Darevskia derjugini*)
barbouri (see *Eremias argus*)
basoglui (see *Ophisops elegans*)
beccarii (see *Podarcis muralis*)
beddomei (see *Ophisops*)
bedriagae (see *Archaeolacerta*)
bedriagai (see *Acanthodactylus*)
beershebensis (see *Acanthodactylus*)
belli (see *Acanthodactylus erythrurus*)
bendimahiensis (see *Darevskia*)
benguellensis (see *Pedioplanis*)
benuensis (see *Acanthodactylus guineensis*)
berlengensis (see *Podarcis carbonelli*)
bernoullii (see *Mesalina brevirostris*)

biinsulicola (see *Podarcis erhardii*)
bileki (see *Anatololacerta danfordi*)
bilineata (see *Lacerta*)
bilkewitschi (see *Eremias scripta*)
bischoffi (see *Darevskia rudis*)
bithynica (see *Darevskia*)
bivittata (see *Ichnotropis*)
bivittata (see *Ophisops jerdonii*)
blanci (see *Acanthodactylus*)
blanci (see *Psammodromus*)
blanfordii (see *Acanthodactylus*)
blanfordi (see *Ophisops elegans*)
bocagei (see *Acanthodactylus erythrurus*)
bocagei (see *Podarcis*)
boehmei (see *Darevskia derjugini*)
boemica (see *Lacerta agilis*)
bokicae (see *Podarcis melisellensis*)
bolei (see *Podarcis siculus*)
bolkardaghica (see *Darevskia rudis*)
bonnali (see *Iberolacerta*)
boristhenis (see *Eremias arguta*)
borkini (see *Eremias velox*)
borromeica (see *Podarcis muralis*)
boscai (see *Latastia*)
boskianus (see *Acanthodactylus*)
bosnica (see *Lacerta agilis*)
boueti (see *Acanthodactylus*)
boulengeri (see *Congolacerta vauereselli*)
boulengeri (see *Nucras*)
brachydactyla (see *Eremias przewalskii*)
brandenburgiensis (see *Lacerta viridis*)
brandtii (see *Iranolacerta*)
brauneri (see *Darevskia*)
brauni (see *Podarcis lilfordi*)
bravoana (see *Gallotia*)
bredoi (see *Latastia johnstonii*)
brenchleyi (see *Eremias*)
brenneri (see *Pseuderemias*)
brevicaudata (see *Lacerta agilis*)
breviceps (see *Pedioplanis*)
breviceps (see *Podarcis muralis*)
brevirostris (see *Mesalina*)
brongniardii (see *Podarcis muralis*)
brueggemanni (see *Podarcis muralis*)
brunnea (see *Ophisops leschenaultii*)
budaki (see *Anatololacerta*)
buechneri (see *Eremias*)
burchelli (see *Pedioplanis*)
buchholzi (see *Podarcis erhardii*)

burii (see *Latastia boscai*)
busacki (see *Acanthodactylus*)

C

caeruleopunctata (see *Latastia*)
caesicaudata (see *Nucras*)
caesaris (see *Gallotia*)
calabra (see *Podarcis siculus*)
calabresiae (see *Podarcis siculus*)
calaesaladae (see *Podarcis pityusensis*)
campana (see *Podarcis siculus*)
campestris (see *Podarcis siculus*)
canaretensis (see *Podarcis pityusensis*)
canensis (see *Podarcis pityusensis*)
cantabrica (see *Iberolacerta monticola*)
cantoris (see *Acanthodactylus*)
capensis (see *Ichnotropis*)
capensis (see *Meroles ctenodactylus*)
cappadocica (see *Apathya*)
caporiacoi (see *Podarcis siculus*)
caprina (see *Podarcis melisellensis*)
caragolensis (see *Podarcis pityusensis*)
carbonelli (see *Podarcis*)
carbonerae (see *Podarcis lilfordi*)
cariensis (see *Lacerta trilineata*)
carlkochi (see *Podarcis pityusensis*)
carniolica (see *Zootoca vivipara*)
caspica (see *Darevskia*)
castilliana (see *Iberolacerta cyreni*)
catanensis (see *Podarcis siculus*)
cattaroi (see *Podarcis siculus*)
caucasia (see *Eremias velox*)
caucasica (see *Darevskia*)
caucasica (see *Darevskia daghestanica*)
cazzae (see *Podarcis siculus*)
cebennensis (see *Podarcis liolepis*)
centralanatoliae (see *Ophisops elegans*)
cerbolensis (see *Podarcis siculus*)
cettii (see *Podarcis siculus*)
chabanaudi (see *Scelarcis perspicillata*)
chaldoranensis (see *Darevskia raddei*)
chalybdea (see *Darevskia armeniaca*)
chaperi (see *Ophisops jerdonii*)
chapini (see *Ichnotropis*)
characae (see *Podarcis pityusensis*)
chechenica (see *Darevskia rudis*)
cherchii (see *Latastia*)
chersonensis (see *Lacerta agilis*)
chlorogaster (see *Darevskia*)

chloronota (see *Lacerta bilineata*)
chlorosecunda (see *Lacerta bilineata*)
cholistanica (see *Eremias*)
chrysogastra (see *Zootoca vivipara*)
ciclopica (see *Podarcis siculus*)
ciliciensis (see *Lacerta media*)
cinereus (see *Zootoca vivipara*)
citrovittata (see *Lacerta trilineata*)
clarkorum (see *Darevskia*)
coccinea (see *Eremias velox*)
codrellensis (see *Podarcis lilfordi*)
coeruleo-ocellata (see *Eremias velox*)
coeruleus (see *Podarcis siculus*)
colchica (see *Lacerta agilis*)
colomi (see *Podarcis lilfordi*)
colosii (see *Podarcis muralis*)
columbretensis (see *Podarcis liolepis*)
conejerae (see *Podarcis lilfordi*)
contii (see *Podarcis tiliguerta*)
cottrelli (see *Tropidosaura*)
cretensis (see *Podarcis*)
cruenta (see *Eremias velox*)
ctenodactylus (see *Meroles*)
cuneirostris (see *Meroles*)
cucchiarai (see *Podarcis raffonei*)
cupreus (see *Takydromus wolteri*)
curatorum (see *Mesalina bahaeldini*)
curzolensis (see *Podarcis melisellensis*)
cyanisparsa (see *Phoenicolacerta*)
cyanura (see *Omanosaura*)
cyreni (see *Iberolacerta*)

D

daghestanica (see *Darevskia*)
dahli (see *Darevskia*)
danfordi (see *Anatololacerta*)
darevskii (see *Darevskia braueri*)
darevskii (see *Eremias arguta*)
defilippii (see *Darevskia*)
degeni (see *Philochortus hardeggeri*)
delibesi (see *Gallotia atlantica*)
derjugini (see *Darevskia*)
deserti (see *Acanthodactylus maculatus*)
deserti (see *Acanthodactylus pardalis*)
deserti (see *Eremias arguta*)
digenea (see *Podarcis melisellensis*)
dilepis (see *Lacerta agilis*)

diplochondrodes (see *Lacerta trilineata*)
dobrogica (see *Lacerta trilineata*)
doriae (see *Psammodromus algirus*)
doriai (see *Latastia*)
dorsalis (see *Acanthodactylus tristrami*)
dorsalis (see *Takydromus*)
dryada (see *Darevskia clarkorum*)
dugesii (see *Teira*)
dumerelii (see *Ichnotropis capensis*)
dumerilii (see *Acanthodactylus*)
dupinici (see *Podarcis siculus*)
dzungarica (see *Eremias*)

E

echinata (see *Gastropholis*)
edwardsii (see *Pseuderemias brenneri*)
edwardsianus (see *Psammodromus*)
ehrenbergii (see *Ophisops elegans*)
eiselti (see *Podarcis tiliguerta*)
eisentrauti (see *Gallotia galloti*)
elaphonisii (see *Podarcis cretensis*)
elbaensis (see *Ophisops*)
elegans (see *Nucras tessellata*)
elegans (see *Ophisops*)
elegans (see *Podarcis siculus*)
emini (see *Nucras boulengeri*)
esfahanica (see *Iranolacerta brandtii*)
espalmadoris (see *Podarcis pityusensis*)
espartellensis (see *Podarcis pityusensis*)
espongicola (see *Podarcis lilfordi*)
ercolinii (see *Mesalina*)
erhardii (see *Podarcis*)
erythrosticta (see *Pseuderemias*)
erythrurus (see *Acanthodactylus*)
essexi (see *Tropidosaura*)
estelicola (see *Podarcis lilfordi*)
euphraticus (see *Acanthodactylus boskianus*)
euxinica (see *Lacerta agilis*)
exigua (see *Lacerta agilis*)
exiguus (see *Acanthodactylus dumerilii*)

F

fahrae (see *Podarcis lilfordi*)
fasciata (see *Eremias*)
fasciatus (see *Podarcis siculus*)
felicis (see *Acanthodactylus*)
fejervaryi (see *Lacerta bilineata*)

fenni (see *Podarcis lilfordi*)
ferrerae (see *Acanthodactylus bedriagae*)
fieldi (see *Mesalina brevirostris*)
filfolensis (see *Podarcis*)
finikensis (see *Anatololacerta pelasgiana*)
fitzingeri (see *Algyroides*)
fiumana (see *Podarcis melisellensis*)
fiumanoideus (see *Podarcis siculus*)
flavigulus (see *Podarcis siculus*)
fordii (see *Poromera*)
formenterae (see *Podarcis pityusensis*)
formosa (see *Pedioplanis lineocellata*)
formosanus (see *Takydromus*)
fraasii (see *Parvilacerta*)
fragilis (see *Zootoca vivipara*)
frailensis (see *Podarcis pityusensis*)
fraseri (see *Acanthodactylus grandis*)
fridericianus (see *Adolfus africanus*)
fusca (see *Zootoca vivipara*)

G

gaerdesi (see *Pedioplanis*)
gaigeae (see *Podarcis*)
galani (see *Iberolacerta*)
galatiensis (see *Lacerta trilineata*)
gallensis (see *Podarcis siculus*)
galloti (see *Gallotia*)
galvagnii (see *Podarcis melisellensis*)
garambensis (see *Heliobolus nitidus*)
gardoensis (see *Pseuderemias striatus*)
garzoni (see *Lacerta agilis*)
gastabiensis (see *Podarcis pityusensis*)
generalensis (see *Podarcis filfolensis*)
gerakuniae (see *Podarcis milensis*)
gigliolii (see *Podarcis lilfordi*)
gigantea (see *Podarcis melisellensis*)
gigas (see *Podarcis melisellensis*)
gomeræ (see *Gallotia caesaris*)
gongrorhynchatus (see *Acanthodactylus*)
gorrae (see *Podarcis pityusensis*)
granchii (see *Podarcis tiliguerta*)
grandiceps (see *Ichnotropis*)
grandis (see *Acanthodactylus*)
grandisonae (see *Podarcis tiliguerta*)
gracilis (see *Darevskia daghestanica*)
gracilis (see *Eremias velox*)
graeca (see *Hellenolacerta*)
grammica (see *Eremias*)
grossae (see *Podarcis pityusensis*)

grueni (see *Podarcis pityusensis*)
grumgrzmailoi (see *Eremias lineolata*)
grusinica (see *Lacerta agilis*)
guadarramae (see *Podarcis*)
guentheri (see *Holaspis*)
guentherpetersi (see *Lacerta viridis*)
guerin (see *Zootoca vivipara*)
guichenoti (see *Mesalina olivieri*)
guineensis (see *Acanthodactylus*)
gularis (see *Tropidosaura*)
guttulata (see *Mesalina*)

H

haackei (see *Pedioplanis*)
haasi (see *Acanthodactylus*)
hadzii (see *Podarcis siculus*)
hani (see *Takydromus*)
hansschweizeri (see *Lacerta trilineata*)
hardeggeri (see *Philochortus*)
hardyi (see *Acanthodactylus*)
harranensis (see *Acanthodactylus*)
hartmanni (see *Podarcis lilfordi*)
haughtonianus (see *Takydromus*)
hedwigkammerae (see *Podarcis pityusensis*)
hesperica (see *Podarcis hispanicus*)
heterolepis (see *Pedioplanis haackei*)
hidalgoi (see *Algyroides*)
hieroglyphicus (see *Podarcis siculus*)
hispanicus (see *Podarcis*)
hispanicus (see *Psammodromus*)
holsti (see *Takydromus tachydromoides*)
holubi (see *Nucras*)
hortae (see *Podarcis pityusensis*)
horvathi (see *Iberolacerta*)
hospitalis (see *Podarcis lilfordi*)
hsuehshanensis (see *Takydromus*)
hungarica (see *Darevskia praticola*)
huntleyi (see *Pedioplanis*)
husabensis (see *Pedioplanis*)
hyrcanica (see *Darevskia praticola*)

I

ibagnezi (see *Gallotia atlantica*)
ibericus (see *Timon lepidus*)
ibrahimi (see *Anatololacerta pelasgiana*)
imperialensis (see *Podarcis lilfordi*)
inaequalis (see *Acanthodactylus boskianus*)
infrapunctata (see *Lacerta viridis*)
inocellata (see *Pedioplanis lineocellata*)

inornata (see *Acanthodactylus dumerilii*)
inornata (see *Pedioplanis*)
insulanagae (see *Gallotia galloti*)
insulanica (see *Podarcis muralis*)
insularus (see *Podarcis siculus*)
intermedia (see *Eremias*)
intermedia (see *Gallotia*)
intermedius (see *Acanthodactylus maculatus*)
intermedius (see *Philochortus*)
intermedius (see *Takydromus*)
intertexta (see *Nucras*)
ionicus (see *Podarcis tauricus*)
ioriensis (see *Lacerta agilis*)
irritans (see *Eremias arguta*)
isaurica (see *Lacerta media*)
isfahanica (see *Eremias*)
israelica (see *Lacerta media*)
istriensis (see *Lacerta viridis*)

J

jacksoni (see *Adolfus*)
jamaicensis (*Timon lepidus*)
japonicus (see *Takydromus tachydromoides*)
jayakari (see *Omanosaura*)
jeanneae (see *Psammodromus algirus*)
jerdonii (see *Ophisops*)
jidulae (see *Podarcis melisellensis*)
joachinimurati (see *Podarcis siculus*)
jogeri (see *Teira dugesii*)
johnstonii (see *Latastia*)
jordansi (see *Podarcis lilfordi*)
judaica (see *Phoenicolacerta laevis*)

K

kamii (see *Darevskia*)
kammereri (see *Podarcis melisellensis*)
kameriana (see *Podarcis pityusensis*)
kavirensis (see *Eremias*)
kephallithacius (see *Algyroides nigropunctatus*)
kessleri (see *Eremias przewalskii*)
ketamensis (see *Psammodromus algirus*)
khamirensis (see *Acanthodactylus*)
khasiensis (see *Takydromus*)
khattensis (see *Acanthodactylus boskianus*)
khazaliensis (see *Phoenicolacerta kulzeri*)
kibonotensis (see *Adolfus*)
kidwelli (see *Latastia johnstonii*)
kieselbachi (see *Podarcis filfolensis*)

kilosae (see *Nucras boulengeri*)
kinarensis (see *Podarcis erhardii*)
klemmeri (see *Podarcis siculus*)
knoxii (see *Meroles*)
kokshaaliensis (see *Eremias*)
kolombatovici (see *Podarcis siculus*)
kopetdaghica (see *Darevskia*)
kopetdaghica (see *Eremias*)
kornatica (see *Podarcis melisellensis*)
kozlowi (see *Eremias multiocellata*)
ksourensis (see *Acanthodactylus erythrurus*)
kuehnei (see *Takydromus*)
kuligae (see *Podarcis lilfordi*)
kulzeri (see *Phoenicolacerta*)
kuri (see *Mesalina*)
kutchensis (see *Ophisops*)
kwangakuensis (see *Takydromus*)
kwangsiensis (see *Takydromus sexlineatus*)

L

laevis (see *Holaspis*)
laevis (see *Phoenicolacerta*)
laganjensis (see *Podarcis siculus*)
lais (see *Podarcis peloponnesiacus*)
lalandii (see *Nucras*)
lalezharica (see *Eremias*)
langi (see *Gastropholis echinata*)
lankae (see *Ophisops leschenaultii*)
lantzicyreni (see *Darevskia valentini*)
lanzai (see *Latastia longicaudata*)
lanzai (see *Podarcis siculus*)
lasdini (see *Eremias scripta*)
latastei (see *Acanthodactylus maculatus*)
latastei (see *Podarcis siculus*)
latasti (see *Mesalina olivieri*)
laticeps (see *Pedioplanis*)
laurae (see *Gallotia atlantica*)
laurentiimuelleri (see *Podarcis filfolensis*)
lepidus (see *Timon*)
leschenaultii (see *Ophisops*)
leucosticta (see *Eremias velox*)
leukaorii (see *Podarcis cretensis*)
levendis (see *Podarcis*)
levithensis (see *Podarcis erhardii*)
lhotei (see *Philochortus zolii*)
lichtensteini (see *Mesalina rubropunctata*)
lilfordi (see *Podarcis*)
lindholmi (see *Darevskia*)

lineolata (see *Eremias*)
lineomaculatus (see *Acanthodactylus erythrurus*)
lineocellata (see *Pedioplanis*)
liolepis (see *Podarcis*)
liscabiancae (see *Podarcis siculus*)
lissana (see *Podarcis melisellensis*)
livida (see *Nucras*)
livadiaca (see *Podarcis erhardii*)
longicaudata (*Latastia*)
longipes (see *Acanthodactylus*)
longipes (see *Ichnotropis capensis*)
loriensis (see *Darevskia praticola*)
louislantzi (see *Zootoca vivipara*)
lugubris (see *Heliobolus*)
lupa (see *Podarcis melisellensis*)
lusitanicus (see *Podarcis guadarramae*)
lutea (see *Zootoca vivipara*)
lutzei (see *Gastropholis vittata*)
luyeanus (see *Takydromus*)

M

machadoi (see *Gallotia simonyi*)
macrodactylus (see *Ophisops elegans*)
macromaculata (see *Darevskia rudis*)
maculatus (see *Acanthodactylus*)
maculiventris (see *Podarcis muralis*)
madaensis (see *Takydromus*)
mahoratae (see *Gallotia atlantica*)
magnifica (see *Lacerta viridis*)
major (see *Lacerta trilineata*)
makariaisi (see *Podarcis erhardii*)
maltensis (see *Podarcis filfolensis*)
maluquerorum (see *Podarcis pityusensis*)
mandchuriensis (see *Takydromus wolteri*)
manuelae (see *Psammodromus algirus*)
marchi (see *Algyroides hidalgoi*)
marcuccii (see *Podarcis muralis*)
maresi (see *Podarcis tiliguerta*)
marettimensis (see *Podarcis waglerianus*)
margaritae (see *Acanthodactylus*)
martensi (see *Latastia doriai*)
martini (see *Mesalina*)
masavaensis (see *Adolfus*)
masirae (see *Acanthodactylus*)
massinei (see *Podarcis siculus*)
martinezricai (see *Iberolacerta*)
mathewsensis (see *Adolfus*)
mauli (see *Teira dugesii*)
mauritanicus (see *Acanthodactylus erythrurus*)
mechriguensis (see *Acanthodactylus maculatus*)

media (see *Lacerta*)
megalophthenae (see *Podarcis erhardii*)
meizolepis (see *Ophisops elegans*)
melanogastra (see *Zootoca vivipara*)
melisellensis (see *Podarcis*)
medemi (see *Podarcis siculus*)
mediofasciatus (see *Podarcis siculus*)
meridionalis (see *Lacerta viridis*)
merremia (see *Podarcis muralis*)
mertensi (see *Podarcis siculus*)
microdactylus (see *Psammodromus*)
microlepidota (see *Ichnotropis capensis*)
microlepidota (see *Ichnotropis*)
microlepis (see *Mesalina brevirostris*)
microlepis (see *Ophisops*)
micropholidotus (see *Meroles*)
micropholis (see *Acanthodactylus*)
miguelensis (see *Podarcis pityusensis*)
mikavicae (see *Podarcis melisellensis*)
milensis (see *Podarcis*)
minor (see *Ophisops*)
mirabilis (see *Darevskia rudis*)
mixta (see *Darevskia*)
monaconensis (see *Podarcis siculus*)
montana (see *Eremias*)
montana (see *Tropidosaura*)
montana (see *Zootoca vivipara*)
montenegrina (see *Dinarolacerta*)
monticola (see *Iberolacerta*)
monticola (see *Ophisops beddomei*)
moreaui (see *Gastropholis prasina*)
moreoticus (see *Algyroides*)
mosorensis (see *Dinarolacerta*)
mostoufii (see *Darevskia chlorogaster*)
mucronata (see *Pseuderemias*)
muellerlorenzi (see *Podarcis muralis*)
muhtari (see *Apathya cappadocica*)
mülleri (see *Podarcis lilfordi*)
multiocellata (see *Eremias*)
muelleri (see *Podarcis siculus*)
muradae (see *Podarcis pityusensis*)
muralis (see *Podarcis*)
mykonensis (see *Podarcis erhardii*)
myusserica (see *Darevskia brauneri*)
mzyntensis (see *Lacerta agilis*)

N

nairensis (see *Darevskia*)
namaquensis (see *Pedioplanis*)
natalensis (see *Tropidosaura montana*)
naxensis (see *Podarcis erhardii*)
nebulosa (see *Mesalina rubropunctata*)
negrae (see *Podarcis pityusensis*)
nevadensis (see *Timon*)
neumanni (see *Heliobolus*)
neumanni (see *Philochortus*)
nicefori (see *Psammodromus algirus*)
nictans (see *Ophisops minor*)
niethammeri (see *Algyroides hidalgoi*)
nigeriensis (see *Acanthodactylus boskianus*)
nigrescens (see *Ichnotropis capensis*)
nigerrima (see *Podarcis lilfordi*)
nigrita (see *Darevskia parvula*)
nigriventris (see *Podarcis muralis*)
nigrocellata (see *Eremias*)
nigrolateralis (see *Eremias persica*)
nigropunctatus (see *Algyroides*)
nikolici (see *Podarcis siculus*)
nikolskii (see *Eremias*)
nilsoni (see *Acanthodactylus*)
nitidus (see *Heliobolus*)
nollii (see *Psammodromus algirus*)
novo (see *Eremias montana*)

O

obscura (see *Darevskia rudis*)
occidentalis (see *Ophisops*)
occidentalis (see *Podarcis muralis*)
occidentalis (see *Psammodromus*)
ocellatus (see *Takydromus sexlineatus*)
oedura (see *Zootoca vivipara*)
oertzeni (see *Anatololacerta anatolica*)
oldi (see *Takydromus tachydromoides*)
olivieri (see *Mesalina*)
opheodurus (see *Acanthodactylus*)
ophidusae (see *Podarcis erhardii*)
oranensis (see *Acanthodactylus savignyi*)
orientalis (see *Acanthodactylus*)
oristanensis (see *Podarcis siculus*)
orlowae (see *Darevskia derjugini*)
ornata (*Latastia*)
ornata (see *Nucras*)
oteroorum (see *Timon lepidus*)
overlaeti (see *Ichnotropis capensis*)
oxycephala (see *Dalmatolacerta*)

oxyrrhina (see *Eremias intermedia*)

P

pachiae (see *Podarcis erhardii*)

paessleri (see *Archaeolacerta bedriagae*)

pallida (see *Ichnotropis bivittata*)

pallida (see *Zootoca vivipara*)

palmae (see *Gallotia galloti*)

palmarolae (see *Podarcis siculus*)

pamphylica (see *Lacerta*)

pannonica (see *Zootoca vivipara*)

panousei (see *Acanthodactylus longipes*)

papenfussi (see *Eremias*)

paradoxa (see *Lacerta agilis*)

pardalis (see *Acanthodactylus*)

pardalis (see *Mesalina guttulata*)

pardaloides (see *Mesalina guttulata*)

pardii (see *Podarcis tiliguerta*)

parkeri (see *Podarcis muralis*)

parva (see *Parvilacerta*)

parvula (see *Darevskia*)

pasquinii (see *Podarcis siculus*)

pasteuri (see *Mesalina*)

pater (see *Timon*)

patrizii (see *Podarcis siculus*)

paulae (see *Podarcis siculus*)

paulinii (see *Podarcis muralis*)

pelagosae (see *Podarcis siculus*)

pelasgiana (see *Anatololacerta*)

pellegrini (see *Scelarcis perspicillata*)

peloponnesiacus (see *Podarcis*)

pentanisiensis (see *Anatololacerta pelasgiana*)

pequensis (see *Meroles knoxii*)

persica (see *Darevskia defilippii*)

persica (see *Eremias*)

persicus (see *Ophisops elegans*)

perspicillata (see *Scelarcis*)

petersiana (see *Latastia*)

petraea (see *Phoenicolacerta kulzeri*)

pherganensis (see *Eremias scripta*)

phillipsi (see *Philochortus*)

phryne (see *Podarcis peloponnesiacus*)

phytiusae (see *Podarcis erhardii*)

pirosoensis (see *Podarcis siculus*)

pityusensis (see *Podarcis*)

planae (see *Podarcis lilfordi*)

planiceps (see *Eremias multiocellata*)

plicata (see *Darevskia praticola*)

plutonis (see *Podarcis melisellensis*)

pohlibensis (see *Podarcis siculus*)
polenci (see *Podarcis siculus*)
polylepidota (see *Lacerta trilineata*)
pomoensis (see *Podarcis melisellensis*)
pontica (see *Darevskia*)
porrosicola (see *Podarcis lilfordi*)
portschinskii (see *Darevskia*)
potanini (see *Eremias arguta*)
prasina (see *Gastropholis*)
praticola (see *Darevskia*)
premudensis (see *Podarcis siculus*)
pretneri (see *Podarcis siculus*)
princeps (see *Timon*)
przewalskii (see *Eremias*)
ptychodes (see *Zootoca vivipara*)
puercosensis (see *Podarcis pityusensis*)
pulchella (see *Pedioplanis lineocellata*)
pulchra (see *Pedioplanis lineocellata*)
punctigularis (see *Podacris cretensis*)
punctulata (see *Mesalina guttulata*)
purroigensis (see *Podarcis pityusensis*)
pushkarensis (see *Ophisops*)
pustulata (see *Eremias arguta*)
pylzowi (see *Eremias vermiculata*)
pyrrhogaster (see *Zootoca vivipara*)

Q

quadrifrons (see *Eremias*)
quadrangularis (see *Pedioplanis namaquensis*)
quadrinasalis (see *Heliobolus nitidus*)
quanttaylori (see *Anatololacerta danfordi*)
quinquevittata (see *Lacerta strigata*)
quinquevittata (see *Pedioplanis burchelli*)

R

raddei (see *Darevskia*)
radovanovici (see *Podarcis siculus*)
raffonei (see *Podarcis*)
ragusae (see *Podarcis siculus*)
rangeri (see *Tropidosaura montana*)
ranzii (see *Podarcis tiliguerta*)
ratae (see *Podarcis pityusensis*)
rechingeri (see *Podacris cretensis*)
redonae (see *Podarcis pityusensis*)
regeli (see *Eremias*)
reticulata (see *Eremias multiocellata*)
reticulatus (see *Meroles*)
revoili (see *Latastia longicaudata*)
riveti (see *Podarcis erhardii*)

roberti (see *Podarcis siculus*)
roborowskii (see *Eremias velox*)
robustus (see *Acanthodactylus*)
rodriquezi (see *Podarcis lilfordi*)
rostombekowi (see *Darevskia*)
rubens (see *Pedioplanis*)
rubropunctata (see *Mesalina*)
rubriventris (see *Podarcis muralis*)
rudolfensis (see *Philochortus*)
rudolphisimonii (see *Podarcis tiliguerta*)
rugiceps (see *Heliobolus spekii*)
rudis (see *Darevskia*)
rupicola (see *Vhembelacerta*)
ruthveni (see *Podarcis erhardii*)

S

sabinae (see *Podarcis pityusensis*)
sachalinensis (see *Zootoca vivipara*)
salfii (see *Podarcis siculus*)
samharica (see *Latastia longicaudata*)
sammichelii (see *Podarcis muralis*)
samogradi (see *Podarcis siculus*)
sanctinicolai (see *Podarcis siculus*)
sanctistephani (see *Podarcis siculus*)
sapphirina (see *Darevskia*)
sargantanae (see *Podarcis lilfordi*)
saturata (see *Eremias multiocellata*)
sauteri (see *Takydromus*)
savagei (see *Pseuderemias*)
savignyi (see *Acanthodactylus*)
saxicola (see *Darevskia*)
scalaris (see *Nucras*)
schaekeli (see *Darevskia*)
schiebeli (see *Podacris cretensis*)
schlueteri (see *Ophisops elegans*)
schmidti (see *Acanthodactylus*)
schmidtii (see *Mesalina olivieri*)
schmidtlerorum (see *Apathya cappadocica*)
schreiberi (see *Acanthodactylus*)
schreiberi (see *Lacerta*)
schreibersiana (see *Zootoca vivipara*)
schreitmuelleri (see *Podarcis pityusensis*)
scopelicus (see *Podarcis siculus*)
scorteccii (see *Heliobolus spekii*)
scorteccii (see *Latastia doriai*)
scorteccii (see *Podarcis siculus*)

scripta (see *Eremias*)
scutellatus (see *Acanthodactylus*)
sebastiani (see *Podarcis liolepis*)
selvagensis (see *Teira dugesii*)
senegalensis (see *Acanthodactylus*)
septemstriata (see *Pseuderemias*)
septentrionalis (see *Takydromus*)
serripes (see *Meroles reticulatus*)
sexlineatus (see *Takydromus*)
sextaeniata (see *Heliobolus spekii*)
siculus (see *Podarcis*)
siebenrocki (see *Latastia*)
sikkimensis (see *Takydromus*)
silvatica (see *Darevskia derjugini*)
simoni (see *Mesalina*)
simonyi (see *Gallotia*)
smaragdinus (see *Takydromus*)
smithii (see *Pseuderemias*)
spekii (see *Heliobolus*)
spinalis (see *Philochortus*)
spinicauda (see *Acanthodactylus*)
spitzenbergerae (see *Darevskia valentini*)
squamulosus (see *Meroles*)
stehlini (see *Gallotia*)
steineri (see *Darevskia*)
stejnegeri (see *Takydromus*)
stenolepis (see *Darevskia praticola*)
stirpium (see *Lacerta agilis*)
strauchi (see *Eremias*)
striatus (see *Pseuderemias*)
strigata (see *Lacerta*)
strombolensis (see *Podarcis siculus*)
stummeri (see *Eremias*)
sturti (see *Latastia longicaudata*)
subformenterae (see *Podarcis pityusensis*)
subobscura (see *Podarcis erhardii*)
suborbitalis (see *Meroles*)
suphani (see *Eremias*)
susana (see *Mesalina olivieri*)
svanetica (see *Darevskia rudis*)
sylvaticus (see *Takydromus*)
syriacus (see *Acanthodactylus schreiberi*)
syrinae (see *Podarcis erhardii*)
szczerbaki (see *Darevskia braueri*)
szczerbaki (see *Eremias*)

T

taeniolata (see *Nucras*)
tachydromoides (see *Takydromus*)
taghitensis (see *Acanthodactylus*)
tagomagensis (see *Podarcis pityusensis*)
tanganicana (see *Ichnotropis*)
tangitanus (see *Timon*)
tauridica (see *Lacerta agilis*)
tauricus (see *Podarcis*)
taylori (see *Latastia*)
tenuis (see *Darevskia raddei*)
terentjevi (see *Darevskia valentini*)
tessellata (see *Nucras*)
thais (see *Podarcis peloponnesiacus*)
thasopulae (see *Podarcis tauricus*)
theobaldi (see *Ophisops jerdonii*)
thermiensis (see *Podarcis erhardii*)
thessalica (see *Podarcis erhardii*)
thetidis (see *Podarcis melisellensis*)
tigrina (see *Eremias velox*)
tilburyi (see *Acanthodactylus*)
tiliguerta (see *Podarcis*)
tinettoi (see *Podarcis muralis*)
tingitanus (see *Acanthodactylus erythrurus*)
tomasinii (see *Dalmatolacerta oxycephala*)
toro (see *Podarcis tiliguerta*)
toronis (see *Podarcis lilfordi*)
torretensis (see *Podarcis pityusensis*)
toyamai (see *Takydromus*)
tracensis (see *Acanthodactylus orientalis*)
traguriana (see *Podarcis melisellensis*)
transcaspica (see *Eremias intermedia*)
transcaucasica (see *Eremias arguta*)
trilineata (see *Lacerta*)
trischittai (see *Podarcis siculus*)
tristis (see *Darevskia bithynica*)
tristrami (see *Acanthodactylus*)
troodica (see *Phoenicolacerta*)
tropidopholis (see *Gastropholis*)
tsaganbogdensis (see *Eremias multiocellata*)
tuvensis (see *Eremias przewalskii*)
typus (see *Takydromus sexlineatus*)
tyrrhenicus (see *Podarcis siculus*)

U

undata (see *Pedioplanis*)
unicolor (see *Zootoca vivipara*)
unisexualis (see *Darevskia*)

urmiana (see *Apathya cappadocica*)
uzbekistanica (see *Eremias arguta*)
uzzelli (see *Darevskia*)

V

vaillanti (see *Acanthodactylus savignyi*)
valentini (see *Darevskia*)
vanensis (see *Darevskia raddei*)
variabilis (see *Eremias arguta*)
vaucheri (see *Podacris*)
vauereselli (see *Congolacerta*)
vedenica (see *Darevskia caucasica*)
vedrae (see *Podacris pityusensis*)
vedranellensis (see *Podacris pityusensis*)
velox (see *Eremias*)
ventotenensis (see *Podacris siculus*)
ventralis (see *Latastia caeruleopunctata*)
vermiculata (see *Eremias*)
vesseljuchi (see *Podarcis siculus*)
vietnamensis (see *Takydromus kuehnei*)
vinciguerrai (see *Podacris muralis*)
virescens (see *Podacris*)
viridipunctatus (see *Takydromus*)
viridis (see *Lacerta*)
vittata (see *Eremias velox*)
vittata (see *Gastropholis*)
vivipara (see *Zootoca*)
vulgaris (see *Acanthodactylus erythrurus*)

W

wachei (see *Latastia boscai*)
waglerianus (see *Podacris*)
watsonana (see *Mesalina*)
weigandi (see *Podacris gaigeae*)
werneriana (see *Podacris erhardii*)
wolteri (see *Apathya cappadocica*)
wolteri (see *Takydromus*)
wolterstorffi (see *Lacerta media*)
woosnami (see *Lacerta strigata*)

X

xapaticola (see *Podacris lilfordi*)

Y

yarkandensis (see *Eremias*)
yassujica (see *Apathya*)
yemenicus (see *Acanthodactylus*)

Z

zafranae (see *Podarcis erhardii*)
zagrosica (see *Iranolacerta*)

zaratinus (see *Podarcis siculus*)
zarudnyi (see *Eremias grammica*)
zeii (see *Podarcis siculus*)
zenonis (see *Podarcis pityusensis*)
zoliei (see *Philochortus*)